



## Donor 6761

### Genetic Testing Summary

Fairfax Cryobank recommends reviewing this genetic testing summary with your healthcare provider to determine suitability.

Last Updated: 8/29/22

Donor Reported Ancestry: Japanese, German, English, Scottish

Jewish Ancestry: No

Genetic Test*	Result	Comments/Donor's Residual Risk**
Chromosome analysis (karyotype)	Normal male karyotype	No evidence of clinically significant chromosome abnormalities
Hemoglobin evaluation	Normal hemoglobin fractionation and MCV/MCH results	Reduced risk to be a carrier for sickle cell anemia, beta thalassemia, alpha thalassemia trait (aa/-- and a-/a-) and other hemoglobinopathies
Expanded Genetic Disease Carrier Screening Panel attached- 502 diseases by gene sequencing.  Personalized residual risk by gene is on attached report.	<b>Carrier: Congenital Adrenal Hyperplasia due to 21-Hydroxylase Deficiency (CYP21A2)</b>  Negative for other genes sequenced	Partner testing recommended before using this donor.

\*No single test can screen for all genetic disorders. A negative screening result significantly reduces, but cannot eliminate, the risk for these conditions in a pregnancy.

\*\*Donor residual risk is the chance the donor is still a carrier after testing negative.

**Patient Information**

Name: Donor 6761  
 Date of Birth: [REDACTED]  
 Sema4 ID: [REDACTED]  
 Client ID: [REDACTED]  
 Indication: Carrier Screening

**Specimen Information**

Specimen Type: Blood  
 Date Collected: 01/25/2022  
 Date Received: 01/26/2022  
 Final Report: 02/19/2022

**Referring Provider**

[REDACTED]  
 Fairfax Cryobank, Inc.  
 [REDACTED]  
 [REDACTED]

## Expanded Carrier Screen (502 genes) with Personalized Residual Risk

### SUMMARY OF RESULTS AND RECOMMENDATIONS

⊕ Positive	⊖ Negative
<p><b>Carrier of Congenital Adrenal Hyperplasia due to 21-Hydroxylase Deficiency (AR)</b>            Associated gene(s): <i>CYP21A2</i>            Variant(s) Detected: c.841G&gt;T, p.V281L, Pathogenic, Heterozygous (one copy)</p>	<p><b>Negative for all other genes tested</b>            To view a full list of genes and diseases tested please see Table 1 in this report</p>

*AR=Autosomal recessive; XL=X-linked*

**Recommendations**

- Testing the partner for the above positive disorder(s) and genetic counseling are recommended.
- Please note that for female carriers of X-linked diseases, follow-up testing of a male partner is not indicated.
- CGG repeat analysis of *FMR1* for fragile X syndrome is not performed on males as repeat expansion of premutation alleles is not expected in the male germline.
- Individuals of Asian, African, Hispanic and Mediterranean ancestry should also be screened for hemoglobinopathies by CBC and hemoglobin electrophoresis.
- Consideration of residual risk by ethnicity after a negative carrier screen is recommended for the other diseases on the panel, especially in the case of a positive family history for a specific disorder.

## Interpretation of positive results

**Congenital Adrenal Hyperplasia due to 21-Hydroxylase Deficiency (AR)**

**Results and Interpretation**

*CYP21A2* copy number: 2  
 No pathogenic copy number variants detected  
*CYP21A2* sequencing: c.841G>T, p.V281L, Pathogenic, Heterozygous (one copy)

**Genes analyzed:** *CYP21A2* (NM\_000500.6)

**Inheritance:** Autosomal Recessive

A heterozygous (one copy) pathogenic missense variant, c.841G>T, p.V281L, was detected in the *CYP21A2* gene (NM\_000500.6). Please note that this variant is typically causative for the non-classic form of congenital adrenal hyperplasia (PMID: 29450859). Variants associated with the non-classic form usually cause non-classic congenital adrenal hyperplasia when found in trans with a pathogenic allele, regardless of whether the second variant is associated with classic or non-classic disease (PMID: 29450859). Therefore, this individual is expected to be at least a carrier for non-classic congenital adrenal hyperplasia. Heterozygous carriers are not expected to exhibit symptoms of this disease.

**What is congenital adrenal hyperplasia (due to 21-hydroxylase deficiency)?**

Congenital adrenal hyperplasia (CAH) is a group of autosomal recessive disorders resulting from deficiency in the enzymes involved in cortisol biosynthesis. The majority (95%) of CAH cases are due to 21-hydroxylase deficiency (21-OHD CAH), which is caused by homozygous or compound heterozygous pathogenic variants in the gene *CYP21A2*. Approximately 20% of mutant alleles have deletions of 30 kb that have been generated by unequal meiotic crossing-over between the two genes. Another 75% of mutant alleles are due to gene conversion events, where an inactivating mutation from the *CYP21A1P* pseudogene is introduced into one copy of the *CYP21A2* gene, thus making the gene non-functional. Three different forms of 21-OHD CAH have been reported: a classic salt wasting form, a classic simple virilizing form, and a non-classic form.

- The classic salt wasting form results from a nonfunctional enzyme and is the most severe. The phenotype includes prenatal onset of virilization and inadequate adrenal aldosterone secretion that can result in fatal salt-wasting crises.
- The classic simple virilizing form results from low levels of functional enzyme and involves prenatal virilization but no salt-wasting.
- The non-classic form, which results from a mild enzyme deficiency, occurs postnatally and involves phenotypes associated with hyperandrogenism, such as hirsutism, delayed menarche, and infertility.

Treatment for the classic forms of the disorder include glucocorticoid and mineralocorticoid replacement therapy, as well as the possibility of feminizing genitoplasty, while patients with the non-classic form usually do not require treatment. The life expectancy for this disorder can be normal with treatment, however the occurrence of salt-wasting crises can be fatal.

## Test description

This patient was tested for a panel of diseases using a combination of sequencing, targeted genotyping and copy number analysis. Please note that negative results reduce but do not eliminate the possibility that this individual is a carrier for one or more of the disorders tested. Please see Table 1 for a list of genes and diseases tested with the patient's personalized residual risk. If personalized residual risk is not provided, please see the complete residual risk table at [go.sema4.com/residualrisk](https://go.sema4.com/residualrisk). Only variants determined to be pathogenic or likely pathogenic are reported in this carrier screening test.



**Lisa Edlmann, Ph.D., FACMG, Laboratory Director**

Laboratory Medical Consultant: George A. Diaz, M.D., Ph.D

## Genes and diseases tested

The personalized residual risks listed below are specific to this individual. The complete residual risk table is available at [go.sema4.com/residualrisk](https://go.sema4.com/residualrisk)

Table 1: List of genes and diseases tested with detailed results

Disease	Gene	Inheritance Pattern	Status	Detailed Summary
<b>Positive</b>				
Congenital Adrenal Hyperplasia due to 21-Hydroxylase Deficiency	<i>CYP21A2</i>	AR	Carrier	<i>CYP21A2</i> copy number: 2 No pathogenic copy number variants detected <i>CYP21A2</i> sequencing: c.841G>T, p.V281L, Pathogenic, Heterozygous (one copy)
<b>Negative</b>				
2-Methylbutyrylglycinuria	<i>ACADSB</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 410
3-Beta-Hydroxysteroid Dehydrogenase Type II Deficiency	<i>HSD3B2</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 3,300
3-Methylcrotonyl-CoA Carboxylase Deficiency (MCCC1-Related)	<i>MCCC1</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 930
3-Methylcrotonyl-CoA Carboxylase Deficiency (MCCC2-Related)	<i>MCCC2</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 500
3-Methylglutaconic Aciduria, Type III	<i>OPA3</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 29,000
3-Phosphoglycerate Dehydrogenase Deficiency	<i>PHGDH</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 63,000
6-Pyruvoyl-Tetrahydropterin Synthase Deficiency	<i>PTS</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 1,800
CD59-Mediated Hemolytic Anemia	<i>CD59</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 513,000
Abetalipoproteinemia	<i>MTTP</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 3,200
Achalasia-Addisonianism-Alacrimia Syndrome	<i>AAAS</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 4,500
Achromatopsia (CNGA3-Related)	<i>CNGA3</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 320
Achromatopsia (CNGB3-related)	<i>CNGB3</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 8,600
Acrodermatitis Enteropathica	<i>SLC39A4</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 12,000
Acute Infantile Liver Failure	<i>TRMU</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 9,400
Acyl-CoA Oxidase I Deficiency	<i>ACOX1</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 39,000
Adams-Oliver Syndrome 4	<i>EOGT</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 44,000
Adenosine Deaminase Deficiency	<i>ADA</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 5,100
Adrenocorticotrophic Hormone Deficiency	<i>TBX19</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 8,500
Adrenoleukodystrophy, X-Linked	<i>ABCD1</i>	XL	Reduced Risk	Personalized Residual Risk: 1 in 19,000
Agammaglobulinemia	<i>BTK</i>	XL	Reduced Risk	Personalized Residual Risk: 1 in 250,000
Agenesis of the Corpus Callosum	<i>FRMD4A</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 348,000
Aicardi-Goutieres Syndrome (RNASEH2C-Related)	<i>RNASEH2C</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 11,000
Aicardi-Goutieres Syndrome (SAMHD1-Related)	<i>SAMHD1</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 2,700
Aicardi-Goutieres Syndrome (TREX1-Related)	<i>TREX1</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 3,200
Albinism, Oculocutaneous, Type III	<i>TYRP1</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 430
Alkaptonuria	<i>HGD</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 1,100
Alpha-Mannosidosis	<i>MAN2B1</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 6,200
Alpha-Thalassemia	<i>HBA1/HBA2</i>	AR	Reduced Risk	<i>HBA1</i> Copy Number: 2 <i>HBA2</i> Copy Number: 2 No pathogenic copy number variants detected <i>HBA1/HBA2</i> Sequencing: Negative Personalized Residual Risk: 1 in 380

Alpha-Thalassemia Intellectual Disability Syndrome	<i>ATRX</i>	XL	Reduced Risk	Personalized Residual Risk: 1 in 48,000
Alport Syndrome ( <i>COL4A3</i> -Related)	<i>COL4A3</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 1,700
Alport Syndrome ( <i>COL4A4</i> -Related)	<i>COL4A4</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 510
Alport Syndrome ( <i>COL4A5</i> -Related)	<i>COL4A5</i>	XL	Reduced Risk	Personalized Residual Risk: 1 in 150,000
Alstrom Syndrome	<i>ALMS1</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 3,100
Andermann Syndrome	<i>SLC12A6</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 151,000
Antley-Bixler Syndrome ( <i>POR</i> -Related)	<i>POR</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 650
Argininemia	<i>ARG1</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 2,900
Argininosuccinic Aciduria	<i>ASL</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 1,200
Aromatase Deficiency	<i>CYP19A1</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 1,200
Arthrogryposis, Intellectual Disability, and Seizures	<i>SLC35A3</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 240,000
Asparagine Synthetase Deficiency	<i>ASNS</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 178,000
Aspartylglycosaminuria	<i>AGA</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 13,000
Ataxia With Isolated Vitamin E Deficiency	<i>TTPA</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 20,000
Ataxia-Telangiectasia	<i>ATM</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 540
Ataxia-Telangiectasia-Like Disorder 1	<i>MRE11</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 6,700
Autosomal Recessive Spastic Ataxia of Charlevoix-Saguenay	<i>SACS</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 2,100
BH4-Deficient Hyperphenylalaninemia C	<i>QDPR</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 3,100
BH4-Deficient Hyperphenylalaninemia D	<i>PCBD1</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 8,000
Bardet-Biedl Syndrome ( <i>ARL6</i> -Related)	<i>ARL6</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 4,300
Bardet-Biedl Syndrome ( <i>BBS10</i> -Related)	<i>BBS10</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 2,100
Bardet-Biedl Syndrome ( <i>BBS12</i> -Related)	<i>BBS12</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 9,900
Bardet-Biedl Syndrome ( <i>BBS1</i> -Related)	<i>BBS1</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 6,400
Bardet-Biedl Syndrome ( <i>BBS2</i> -Related)	<i>BBS2</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 1,200
Bardet-Biedl Syndrome ( <i>BBS4</i> -Related)	<i>BBS4</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 22,000
Bare Lymphocyte Syndrome, Type II	<i>CIITA</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 35,000
Barth Syndrome	<i>TAZ</i>	XL	Reduced Risk	Personalized Residual Risk: 1 in 183,000
Bartter Syndrome, Type 3	<i>CLCNKB</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 710
Bartter Syndrome, Type 4A	<i>BSND</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 69,000
Bernard-Soulier Syndrome, Type A1	<i>GP1BA</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 42,000
Bernard-Soulier Syndrome, Type C	<i>GP9</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 1,100
Beta-Globin-Related Hemoglobinopathies	<i>HBB</i>	AR	Reduced Risk	Personalized Residual Risk (Beta-Globin-Related Hemoglobinopathies): 1 in 1,200 Personalized Residual Risk (Beta-Globin-Related Hemoglobinopathies: HbS Variant): 1 in 11,000 Personalized Residual Risk (Beta-Globin-Related Hemoglobinopathies: HbC Variant): 1 in 42,000
Beta-Ketothiolase Deficiency	<i>ACAT1</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 3,800
Beta-Mannosidosis	<i>MANBA</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 9,100
Bilateral Frontoparietal Polymicrogyria	<i>GPR56</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 143,000
Biotinidase Deficiency	<i>BTD</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 500
Bloom Syndrome	<i>BLM</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 7,400
Canavan Disease	<i>ASPA</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 4,000
Carbamoylphosphate Synthetase I Deficiency	<i>CPS1</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 690
Carnitine Acylcarnitine Translocase Deficiency	<i>SLC25A20</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 4,100
Carnitine Palmitoyltransferase IA Deficiency	<i>CPT1A</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 24,000
Carnitine Palmitoyltransferase II Deficiency	<i>CPT2</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 670
Carpenter Syndrome	<i>RAB23</i>	AR	Reduced Risk	Personalized Residual Risk: 1 in 21,000

Cartilage-Hair Hypoplasia	<i>RMRP</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 450</b>
Catecholaminergic Polymorphic Ventricular Tachycardia	<i>CASQ2</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 5,900</b>
Central Hypothyroidism and Testicular Enlargement	<i>IGSF1</i>	XL	Reduced Risk	<b>Personalized Residual Risk: 1 in 781,000</b>
Cerebral Creatine Deficiency Syndrome 1	<i>SLC6A8</i>	XL	Reduced Risk	<b>Personalized Residual Risk: 1 in 208,000</b>
Cerebral Creatine Deficiency Syndrome 2	<i>GAMT</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 2,300</b>
Cerebral Creatine Deficiency Syndrome 3	<i>GATM</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 4,000</b>
Cerebral Dysgenesis, Neuropathy, Ichthyosis, and Palmoplantar Keratoderma Syndrome	<i>SNAP29</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 383,000</b>
Cerebrotendinous Xanthomatosis	<i>CYP27A1</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 750</b>
Charcot-Marie-Tooth Disease, Type 4D	<i>NDRG1</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 225,000</b>
Charcot-Marie-Tooth Disease, Type 5 / Arts Syndrome	<i>PRPS1</i>	XL	Reduced Risk	<b>Personalized Residual Risk: 1 in 114,000</b>
Charcot-Marie-Tooth Disease, X-Linked	<i>GJB1</i>	XL	Reduced Risk	<b>Personalized Residual Risk: 1 in 11,000</b>
Chediak-Higashi Syndrome	<i>LYST</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 7,100</b>
Chondrodysplasia Punctata	<i>ARSE</i>	XL	Reduced Risk	<b>Personalized Residual Risk: 1 in 862,000</b>
Choreoacanthocytosis	<i>VPS13A</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 4,700</b>
Choroideremia	<i>CHM</i>	XL	Reduced Risk	<b>Personalized Residual Risk: 1 in 125,000</b>
Chronic Granulomatous Disease (CYBA-Related)	<i>CYBA</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 3,700</b>
Chronic Granulomatous Disease (CYBB-Related)	<i>CYBB</i>	XL	Reduced Risk	<b>Personalized Residual Risk: 1 in 294,000</b>
Citrin Deficiency	<i>SLC25A13</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 2,200</b>
Citrullinemia, Type 1	<i>ASS1</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 2,500</b>
Cockayne Syndrome, Type A	<i>ERCC8</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 8,900</b>
Cockayne Syndrome, Type B and other ERCC6-Related Disorders	<i>ERCC6</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 5,300</b>
Cohen Syndrome	<i>VPS13B</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 6,400</b>
Combined Factor V and VIII Deficiency	<i>LMAN1</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 68,000</b>
Combined Malonic and Methylmalonic Aciduria	<i>ACSF3</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 2,400</b>
Combined Oxidative Phosphorylation Deficiency 1	<i>GFM1</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 8,100</b>
Combined Oxidative Phosphorylation Deficiency 3	<i>TSMF</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 21,000</b>
Combined Pituitary Hormone Deficiency 1	<i>POU1F1</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 1,900</b>
Combined Pituitary Hormone Deficiency 2	<i>PROP1</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 2,800</b>
Combined Pituitary Hormone Deficiency 3	<i>LHX3</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 121,000</b>
Combined SAP Deficiency	<i>PSAP</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 44,000</b>
Cone-Rod Dystrophy 6 / Leber Congenital Amaurosis 1	<i>GUCY2D</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 720</b>
Congenital Adrenal Hyperplasia due to 11-Beta-Hydroxylase Deficiency	<i>CYP11B1</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 520</b>
Congenital Adrenal Hyperplasia due to 17-Alpha-Hydroxylase Deficiency	<i>CYP17A1</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 840</b>
Congenital Adrenal Hypoplasia (NR0B1-Related)	<i>NR0B1</i>	XL	Reduced Risk	<b>Personalized Residual Risk: 1 in 353,000</b>
Congenital Adrenal Insufficiency (CYP11A1-Related)	<i>CYP11A1</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 6,100</b>
Congenital Amegakaryocytic Thrombocytopenia	<i>MPL</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 3,100</b>
Congenital Bile Acid Synthesis Defect (AKR1D1-Related)	<i>AKR1D1</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 6,900</b>
Congenital Bile Acid Synthesis Defect (HSD3B7-Related)	<i>HSD3B7</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 4,700</b>
Congenital Disorder of Deglycosylation	<i>NGLY1</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 2,000</b>
Congenital Disorder of Glycosylation, Type Ia	<i>PMM2</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 550</b>
Congenital Disorder of Glycosylation, Type Ib	<i>MPL</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 2,100</b>
Congenital Disorder of Glycosylation, Type Ic	<i>ALG6</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 2,300</b>
Congenital Disorder of Glycosylation, Type Im	<i>DOLK</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 134,000</b>

<b>Congenital Dyserythropoietic Anemia Type 2</b>	<i>SEC23B</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 1,000</b>
<b>Congenital Dyserythropoietic Anemia, Type Ia</b>	<i>CDAN1</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 470</b>
<b>Congenital Ichthyosis 4A and 4B</b>	<i>ABCA12</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 1,500</b>
<b>Congenital Insensitivity to Pain with Anhidrosis</b>	<i>NTRK1</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 1,000</b>
<b>Congenital Muscular Dystrophy (LAMA2-Related)</b>	<i>LAMA2</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 690</b>
<b>Congenital Myasthenic Syndrome (CHAT-Related)</b>	<i>CHAT</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 1,500</b>
<b>Congenital Myasthenic Syndrome (CHRNE-Related)</b>	<i>CHRNE</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 4,100</b>
<b>Congenital Myasthenic Syndrome (DOK7-Related)</b>	<i>DOK7</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 470</b>
<b>Congenital Myasthenic Syndrome (RAPSN-Related)</b>	<i>RAPSN</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 2,900</b>
<b>Congenital Neutropenia (HAX1-Related)</b>	<i>HAX1</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 82,000</b>
<b>Congenital Neutropenia (VPS45-Related)</b>	<i>VPS45</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 110,000</b>
<b>Congenital Nongoitrous Hypothyroidism 1</b>	<i>TSHR</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 230</b>
<b>Congenital Nongoitrous Hypothyroidism 4</b>	<i>TSHB</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 118,000</b>
<b>Congenital Secretory Chloride Diarrhea 1</b>	<i>SLC26A3</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 2,400</b>
<b>Corneal Dystrophy and Perceptive Deafness</b>	<i>SLC4A11</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 2,200</b>
<b>Corticosterone Methyloxidase Deficiency</b>	<i>CYP11B2</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 1,700</b>
<b>Cystic Fibrosis</b>	<i>CFTR</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 440</b>
<b>Cystinosis</b>	<i>CTNS</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 7,100</b>
<b>Cystinuria (SLC3A1-Related)</b>	<i>SLC3A1</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 530</b>
<b>Cytochrome C Oxidase Deficiency / Leigh Syndrome (COX15-Related)</b>	<i>COX15</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 3,300</b>
<b>D-Bifunctional Protein Deficiency</b>	<i>HSD17B4</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 2,700</b>
<b>Deafness, Autosomal Recessive 3</b>	<i>MYO15A</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 100</b>
<b>Deafness, Autosomal Recessive 59</b>	<i>PJVK</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 57,000</b>
<b>Deafness, Autosomal Recessive 7</b>	<i>TMC1</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 1,100</b>
<b>Deafness, Autosomal Recessive 76</b>	<i>SYNE4</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 43,000</b>
<b>Deafness, Autosomal Recessive 77</b>	<i>LOXHD1</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 2,800</b>
<b>Deafness, Autosomal Recessive 8/10</b>	<i>TMPRSS3</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 330</b>
<b>Deafness, Autosomal Recessive 9</b>	<i>OTOF</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 370</b>
<b>Desbuquois Dysplasia 1</b>	<i>CANT1</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 7,800</b>
<b>Desmosterolosis</b>	<i>DHCR24</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 28,000</b>
<b>Diaphanospondylodysostosis</b>	<i>BMPER</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 18,000</b>
<b>Distal Renal Tubular Acidosis and other SLC4A1-related Disorders</b>	<i>SLC4A1</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 910</b>
<b>Duchenne Muscular Dystrophy / Becker Muscular Dystrophy</b>	<i>DMD</i>	XL	Reduced Risk	<b>Personalized Residual Risk: 1 in 10,000</b>
<b>Dyskeratosis Congenita (DKC1-related)</b>	<i>DKC1</i>	XL	Reduced Risk	<b>Personalized Residual Risk: 1 in 9,259,000</b>
<b>Dyskeratosis Congenita (RTEL1-Related)</b>	<i>RTEL1</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 3,900</b>
<b>Dystrophic Epidermolysis Bullosa</b>	<i>COL7A1</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 900</b>
<b>Ehlers-Danlos Syndrome, Type VI</b>	<i>PLOD1</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 8,700</b>
<b>Ehlers-Danlos Syndrome, Type VIIC</b>	<i>ADAMTS2</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 63,000</b>
<b>Ellis-Van Creveld Syndrome (EVC2-Related)</b>	<i>EVC2</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 6,100</b>
<b>Ellis-van Creveld Syndrome (EVC-Related)</b>	<i>EVC</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 4,200</b>
<b>Emery-Dreifuss Myopathy 1</b>	<i>EMD</i>	XL	Reduced Risk	<b>Personalized Residual Risk: 1 in 833,000</b>
<b>Enhanced S-Cone Syndrome</b>	<i>NR2E3</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 1,700</b>
<b>Ethylmalonic Encephalopathy</b>	<i>ETHE1</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 3,400</b>
<b>Fabry Disease</b>	<i>GLA</i>	XL	Reduced Risk	<b>Personalized Residual Risk: 1 in 7,700</b>
<b>Factor IX Deficiency</b>	<i>F9</i>	XL	Reduced Risk	<b>Personalized Residual Risk: 1 in 5,100</b>
<b>Factor VII Deficiency</b>	<i>F7</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 300</b>

Factor XI Deficiency	<i>F11</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 440</b>
Familial Autosomal Recessive Hypercholesterolemia	<i>LDLRAP1</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 136,000</b>
Familial Dysautonomia	<i>IKBKAP</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 51,000</b>
Familial Hypercholesterolemia	<i>LDLR</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 260</b>
Familial Hyperinsulinemic Hypoglycemia 4 / 3-Hydroxyacyl-CoA Dehydrogenase Deficiency	<i>HADH</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 6,000</b>
Familial Hyperinsulinism (ABCC8-Related)	<i>ABCC8</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 240</b>
Familial Hyperinsulinism (KCNJ11-Related)	<i>KCNJ11</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 4,300</b>
Familial Hyperphosphatemic Tumoral Calcinosis	<i>GALNT3</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 4,100</b>
Familial Mediterranean Fever	<i>MEFV</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 1,200</b>
Fanconi Anemia, Group A	<i>FANCA</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 1,100</b>
Fanconi Anemia, Group C	<i>FANCC</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 12,000</b>
Fanconi Anemia, Group G	<i>FANCG</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 1,200</b>
Fanconi-Bickel Syndrome	<i>SLC2A2</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 4,000</b>
Fragile X Syndrome	<i>FMR1</i>	XL	Reduced Risk	<i>FMR1</i> CGG repeat sizes: Not Performed <i>FMR1</i> Sequencing: Negative Fragile X CGG triplet repeat expansion testing was not performed at this time, as the patient has either been previously tested or is a male. <b>Personalized Residual Risk: 1 in 19,000</b>
Fructose-1,6-Bisphosphatase Deficiency	<i>FBP1</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 2,600</b>
Fucosidosis	<i>FUCA1</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 9,200</b>
Fumarase Deficiency	<i>FH</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 2,900</b>
Fundus Albipunctatus	<i>RDH5</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 810</b>
GRACILE Syndrome and Other <i>BCS1L</i> -Related Disorders	<i>BCS1L</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 3,900</b>
Galactokinase Deficiency	<i>GALK1</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 1,600</b>
Galactose Epimerase Deficiency	<i>GALE</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 850</b>
Galactosemia	<i>GALT</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 390</b>
Galactosialidosis	<i>CTSA</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 7,900</b>
Gaucher Disease	<i>GBA</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 1,200</b>
Generalized Thyrotropin-Releasing Hormone Resistance	<i>TRHR</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 104,000</b>
Geroderma Osteodysplasticum	<i>GORAB</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 76,000</b>
Gitelman Syndrome	<i>SLC12A3</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 230</b>
Glanzmann Thrombasthenia ( <i>ITGA2B</i> -Related)	<i>ITGA2B</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 1,200</b>
Glanzmann Thrombasthenia ( <i>ITGB3</i> -Related)	<i>ITGB3</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 1,200</b>
Glutaric Acidemia, Type I	<i>GCDH</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 2,700</b>
Glutaric Acidemia, Type IIa	<i>ETFA</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 2,100</b>
Glutaric Acidemia, Type IIb	<i>ETFB</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 5,900</b>
Glutaric Acidemia, Type IIc	<i>ETFDH</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 260</b>
Glutathione Synthetase Deficiency	<i>GSS</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 3,500</b>
Glycine Encephalopathy ( <i>AMT</i> -Related)	<i>AMT</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 5,700</b>
Glycine Encephalopathy ( <i>GLDC</i> -Related)	<i>GLDC</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 240</b>
Glycogen Storage Disease, Type 0	<i>GYS2</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 1,200</b>
Glycogen Storage Disease, Type II	<i>GAA</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 280</b>
Glycogen Storage Disease, Type III	<i>AGL</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 5,600</b>
Glycogen Storage Disease, Type IV / Adult Polyglucosan Body Disease	<i>GBE1</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 2,400</b>
Glycogen Storage Disease, Type IXb	<i>PHKB</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 2,600</b>
Glycogen Storage Disease, Type Ia	<i>G6PC</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 410</b>
Glycogen Storage Disease, Type Ib	<i>SLC37A4</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 3,900</b>
Glycogen Storage Disease, Type V	<i>PYGM</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 1,400</b>



Glycogen Storage Disease, Type VI	<i>PYGL</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,600
Glycogen Storage Disease, Type VII	<i>PFKM</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 4,300
Gray Platelet Syndrome	<i>NBEAL2</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 7,400
Growth Hormone Deficiency, Type IB	<i>GHRHR</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 3,900
HMG-CoA Lyase Deficiency	<i>HMGCL</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 2,700
Hemochromatosis, Type 2A	<i>HFE2</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 740
Hemochromatosis, Type 3	<i>TFR2</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 11,000
Hereditary Fructose Intolerance	<i>ALDOB</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,900
Hereditary Spastic Paraparesis 49	<i>TECPR2</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 116,000
Hermansky-Pudlak Syndrome, Type 1	<i>HPS1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 3,500
Hermansky-Pudlak Syndrome, Type 3	<i>HPS3</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 22,000
Hermansky-Pudlak Syndrome, Type 4	<i>HPS4</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 35,000
Hermansky-Pudlak Syndrome, Type 6	<i>HPS6</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 680
Hmg-CoA Synthase 2 Deficiency	<i>HMGCS2</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 2,000
Holocarboxylase Synthetase Deficiency	<i>HLCS</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 5,500
Homocystinuria (CBS-Related)	<i>CBS</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,400
Homocystinuria due to <i>MTHFR</i> Deficiency	<i>MTHFR</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,300
Homocystinuria, cblE Type	<i>MTRR</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 9,600
Homocystinuria-Megaloblastic Anemia, Cobalamin G Type	<i>MTR</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 2,100
Hydrocephalus	<i>L1CAM</i>	XL	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 40,000
Hydroletharus Syndrome	<i>HYLS1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 52,000
Hyper-Igm Syndrome	<i>CD40LG</i>	XL	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,167,000
Hyperornithinemia-Hyperammonemia-Homocitrullinuria Syndrome	<i>SLC25A15</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 5,700
Hyperuricemia, Pulmonary Hypertension, Renal Failure, and Alkalosis	<i>SARS2</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 23,000
Hypohidrotic Ectodermal Dysplasia 1	<i>EDA</i>	XL	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 22,000
Hypomagnesemia 1	<i>TRPM6</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 11,000
Hypomyelinating Leukodystrophy 3	<i>AIMP1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 273,000
Hypomyelinating Leukodystrophy 12	<i>VPS11</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 72,000
Hypoparathyroidism-Retardation-Dysmorphic Syndrome	<i>TBCE</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 21,000
Hypophosphatasia	<i>ALPL</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 790
Hypophosphatemic Rickets with Hypercalciuria	<i>SLC34A3</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,000
Hypotrichosis 8 / Autosomal Recessive Woolly Hair 1	<i>LPAR6</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 17,000
Immunodeficiency 18	<i>CD3E</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 73,000
Immunodeficiency 19	<i>CD3D</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 46,000
Inclusion Body Myopathy 2	<i>GNE</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 2,000
Infantile Cerebral and Cerebellar Atrophy	<i>MED17</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 130,000
Infantile Neuroaxonal Dystrophy 1 and other <i>PLA2G6</i> -Related Disorders	<i>PLA2G6</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 380
Intellectual Disability, Autosomal Recessive 3	<i>CC2D1A</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 108,000
Intrahepatic Cholestasis	<i>ATP8B1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 580
Isovaleric Acidemia	<i>IVD</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 2,000
Joubert Syndrome 2	<i>TMEM216</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 133,000
Joubert Syndrome 4 / Senior-Loken Syndrome 1 / Juvenile Nephronophthisis 1	<i>NPHP1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 2,000
Joubert Syndrome 7 / Meckel Syndrome 5 / COACH Syndrome	<i>RPGRIP1L</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,100
Junctional Epidermolysis Bullosa ( <i>COL17A1</i> -Related)	<i>COL17A1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 4,200
Junctional Epidermolysis Bullosa ( <i>ITGA6</i> -Related)	<i>ITGA6</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 125,000



Junctional Epidermolysis Bullosa ( <i>ITGB4</i> -Related)	<i>ITGB4</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 2,400
Junctional Epidermolysis Bullosa ( <i>LAMA3</i> -Related)	<i>LAMA3</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 21,000
Junctional Epidermolysis Bullosa ( <i>LAMB3</i> -Related)	<i>LAMB3</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,900
Junctional Epidermolysis Bullosa ( <i>LAMC2</i> -Related)	<i>LAMC2</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 28,000
Kohlschutter-Tonz Syndrome	<i>ROGDI</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 2,300
Krabbe Disease	<i>GALC</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 340
Lamellar Ichthyosis, Type 1	<i>TGM1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,500
Laron Dwarfism	<i>GHR</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 2,100
Leber Congenital Amaurosis 10 and Other CEP290-Related Ciliopathies	<i>CEP290</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,100
Leber Congenital Amaurosis 13	<i>RDH12</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 5,500
Leber Congenital Amaurosis 15 / Retinitis Pigmentosa 14	<i>TULP1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,600
Leber Congenital Amaurosis 2 / Retinitis Pigmentosa 20	<i>RPE65</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 2,100
Leber Congenital Amaurosis 4	<i>AIP1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 2,100
Leber Congenital Amaurosis 5	<i>LCA5</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 4,200
Leber Congenital Amaurosis 8 / Retinitis Pigmentosa 12 / Pigmented Paravenous Chorioretinal Atrophy	<i>CRB1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 960
Leigh Syndrome ( <i>NDUFS7</i> -Related)	<i>NDUFS7</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 26,000
Leigh Syndrome ( <i>SURF1</i> -Related)	<i>SURF1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 4,000
Leigh Syndrome, French-Canadian Type	<i>LRPPRC</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 22,000
Lethal Congenital Contracture Syndrome 1 / Lethal Arthrogyrosis with Anterior Horn Cell Disease	<i>GLE1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 5,900
Lethal Congenital Contracture Syndrome 2	<i>ERBB3</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 52,000
Lethal Congenital Contracture Syndrome 3	<i>PIP5K1C</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 304,000
Leukoencephalopathy with Vanishing White Matter	<i>EIF2B5</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 2,300
Limb-Girdle Muscular Dystrophy, Type 2A	<i>CAPN3</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 960
Limb-Girdle Muscular Dystrophy, Type 2B	<i>DYSF</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,100
Limb-Girdle Muscular Dystrophy, Type 2C	<i>SGCG</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 4,900
Limb-Girdle Muscular Dystrophy, Type 2D	<i>SGCA</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 3,500
Limb-Girdle Muscular Dystrophy, Type 2E	<i>SGCB</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 31,000
Limb-Girdle Muscular Dystrophy, Type 2F	<i>SGCD</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 36,000
Limb-Girdle Muscular Dystrophy, Type 2H	<i>TRIM32</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 10,000
Limb-Girdle Muscular Dystrophy, Type 2I	<i>FKRP</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 460
Limb-Girdle Muscular Dystrophy, Type 2L	<i>ANO5</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 660
Lipoamide Dehydrogenase Deficiency	<i>DLA</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 14,000
Lipoid Adrenal Hyperplasia	<i>STAR</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 3,600
Lipoprotein Lipase Deficiency	<i>LPL</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 800
Long-Chain 3-Hydroxyacyl-CoA Dehydrogenase Deficiency	<i>HADHA</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 4,500
Lowe Syndrome	<i>OCRL</i>	XL	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,375,000
Lysinuric Protein Intolerance	<i>SLC7A7</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 3,000
MEDNIK Syndrome	<i>API51</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 211,000
Malonyl-CoA Decarboxylase Deficiency	<i>MLYCD</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 2,800
Maple Syrup Urine Disease, Type 1a	<i>BCKDHA</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 4,000
Maple Syrup Urine Disease, Type 1b	<i>BCKDHB</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,100
Maple Syrup Urine Disease, Type 2	<i>DBT</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 790
Meckel Syndrome 1 / Bardet-Biedl Syndrome 13	<i>MKS1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,700

Medium Chain Acyl-CoA Dehydrogenase Deficiency	<i>ACADM</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,600
Megalencephalic Leukoencephalopathy with Subcortical Cysts	<i>MLC1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 4,300
Megaloblastic Anemia 1	<i>AMN</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 6,300
Menkes Disease	<i>ATP7A</i>	XL	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 172,000
Metachromatic Leukodystrophy	<i>ARSA</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,000
Methionine Adenosyltransferase I/III Deficiency	<i>MAT1A</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,700
Methylmalonic Acidemia (MMAA-Related)	<i>MMAA</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 15,000
Methylmalonic Acidemia (MMAB-Related)	<i>MMAB</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 3,700
Methylmalonic Acidemia (MUT-Related)	<i>MUT</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 830
Methylmalonic Aciduria and Homocystinuria, Cobalamin C Type	<i>MMACHC</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,300
Methylmalonic Aciduria and Homocystinuria, Cobalamin D Type	<i>MMADHC</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 172,000
Methylmalonic Aciduria and Homocystinuria, Cobalamin F Type	<i>LMBRD1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 6,600
Methylmalonyl-CoA Epimerase Deficiency	<i>MCEE</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 98,000
Microphthalmia / Anophthalmia	<i>VSX2</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 40,000
Mitochondrial Complex I Deficiency (ACAD9-Related)	<i>ACAD9</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,800
Mitochondrial Complex I Deficiency (NDUFA11-Related)	<i>NDUFA11</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 414,000
Mitochondrial Complex I Deficiency (NDUFAF5-Related)	<i>NDUFAF5</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 770
Mitochondrial Complex I Deficiency (NDUFS6-Related)	<i>NDUFS6</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 211,000
Mitochondrial Complex I Deficiency (NDUFV1-Related)	<i>NDUFV1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 870
Mitochondrial Complex I Deficiency / Leigh Syndrome (FOXRED1-Related)	<i>FOXRED1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 6,900
Mitochondrial Complex I Deficiency / Leigh Syndrome (NDUFAF2-Related)	<i>NDUFAF2</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 114,000
Mitochondrial Complex I Deficiency / Leigh Syndrome (NDUFS4-Related)	<i>NDUFS4</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 31,000
Mitochondrial Complex IV Deficiency (COX20-related)	<i>COX20</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 42,000
Mitochondrial Complex IV Deficiency (COX6B1-related)	<i>COX6B1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,231,000
Mitochondrial Complex IV Deficiency (APOPT1-Related)	<i>APOPT1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 9,200
Mitochondrial Complex IV Deficiency (PET100-Related)	<i>PET100</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 546,000
Mitochondrial Complex IV Deficiency (SCO1-related)	<i>SCO1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 13,000
Mitochondrial Complex IV Deficiency / Leigh Syndrome (COX10-Related)	<i>COX10</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 6,900
Mitochondrial DNA Depletion Syndrome 2	<i>TK2</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 5,700
Mitochondrial DNA Depletion Syndrome 3	<i>DGUOK</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 5,200
Mitochondrial DNA Depletion Syndrome 4A and 4B and other POLG-Related Disorders	<i>POLG</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 180
Mitochondrial DNA Depletion Syndrome 5	<i>SUCLA2</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 78,000
Mitochondrial DNA Depletion Syndrome 6 / Navajo Neurohepatopathy	<i>MPV17</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 4,400
Mitochondrial Myopathy and Sideroblastic Anemia 1	<i>PUS1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 333,000
Mitochondrial Trifunctional Protein Deficiency (HADHB-Related)	<i>HADHB</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,300
Molybdenum Cofactor Deficiency A	<i>MOCS1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 4,700
Mucopolipidosis II / IIIA	<i>GNPTAB</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,100
Mucopolipidosis III Gamma	<i>GNPTG</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 68,000

Mucopolipidosis IV	<i>MCOLN1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 6,500
Mucopolysaccharidosis Type I	<i>IDUA</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 630
Mucopolysaccharidosis Type II	<i>IDS</i>	XL	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 76,000
Mucopolysaccharidosis Type IIIA	<i>SGSH</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 700
Mucopolysaccharidosis Type IIIB	<i>NAGLU</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 900
Mucopolysaccharidosis Type IIIC	<i>HGSNAT</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 3,200
Mucopolysaccharidosis Type IIID	<i>GNS</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 137,000
Mucopolysaccharidosis Type IVa	<i>GALNS</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 440
Mucopolysaccharidosis Type IVb / GM1 Gangliosidosis	<i>GLB1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,200
Mucopolysaccharidosis VII	<i>GUSB</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,600
Mucopolysaccharidosis type IX	<i>HYAL1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 63,000
Mucopolysaccharidosis type VI	<i>ARSB</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,300
Mulibrey Nanism	<i>TRIM37</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 36,000
Multiple Congenital Anomalies-Hypotonia-Seizures Syndrome 1	<i>PIGN</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 2,800
Multiple Pterygium Syndrome	<i>CHRNA3</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 4,000
Multiple Sulfatase Deficiency	<i>SUMF1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 69,000
Muscle-Eye-Brain Disease and Other <i>POMGNT1</i> -Related Congenital Muscular Dystrophy-Dystroglycanopathies	<i>POMGNT1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 4,200
Myoneurogastrointestinal Encephalopathy	<i>TYMP</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 2,100
Myotubular Myopathy 1	<i>MTM1</i>	XL	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 192,000
N-Acetylglutamate Synthase Deficiency	<i>NAGS</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 3,200
Nemaline Myopathy 2	<i>NEB</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 300
Nephrogenic Diabetes Insipidus, Type II	<i>AQP2</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 3,400
Nephrogenic Diabetes insipidus ( <i>AVPR2</i> -related)/ Nephrogenic Syndrome of Inappropriate Antidiuresis	<i>AVPR2</i>	XL	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 471,000
Nephronophthisis 2	<i>INVS</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 24,000
Nephrotic Syndrome ( <i>NPHS1</i> -Related) / Congenital Finnish Nephrosis	<i>NPHS1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 980
Nephrotic Syndrome ( <i>NPHS2</i> -Related) / Steroid-Resistant Nephrotic Syndrome	<i>NPHS2</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 780
Neurodegeneration due to Cerebral Folate Transport Deficiency	<i>FOLR1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 3,700
Neurodevelopmental Disorder with Progressive Microcephaly, Spasticity, and Brain Anomalies	<i>PLAA</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 217,000
Neuronal Ceroid-Lipofuscinosis ( <i>CLN3</i> -Related)	<i>CLN3</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 9,200
Neuronal Ceroid-Lipofuscinosis ( <i>CLN5</i> -Related)	<i>CLN5</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 4,300
Neuronal Ceroid-Lipofuscinosis ( <i>CLN6</i> -Related)	<i>CLN6</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 8,600
Neuronal Ceroid-Lipofuscinosis ( <i>CLN8</i> -Related)	<i>CLN8</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 3,100
Neuronal Ceroid-Lipofuscinosis ( <i>MFSD8</i> -Related)	<i>MFSD8</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 6,200
Neuronal Ceroid-Lipofuscinosis ( <i>PPT1</i> -Related)	<i>PPT1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 2,300
Neuronal Ceroid-Lipofuscinosis ( <i>TPP1</i> -Related)	<i>TPP1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 3,000
Niemann-Pick Disease ( <i>SMPD1</i> -Related)	<i>SMPD1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,300
Niemann-Pick Disease, Type C ( <i>NPC1</i> -Related)	<i>NPC1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 690
Niemann-Pick Disease, Type C ( <i>NPC2</i> -Related)	<i>NPC2</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 6,600
Nijmegen Breakage Syndrome	<i>NBN</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 14,000
Non-Syndromic Hearing Loss ( <i>GJB2</i> -Related)	<i>GJB2</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 280
Oculocutaneous Albinism, Type IA / IB	<i>TYR</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 220
Oculocutaneous Albinism, Type IV	<i>SLC45A2</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 980
Odonto-Onycho-Dermal Dysplasia / Schopf-Schulz-Passarge Syndrome	<i>WNT10A</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 900

Omenn Syndrome ( <i>RAG2</i> -Related)	<i>RAG2</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 17,000
Omenn Syndrome / Severe Combined Immunodeficiency, Athabaskan-Type	<i>DCLRE1C</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 5,500
Omenn Syndrome and other <i>RAG1</i> -Related Disorders	<i>RAG1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 180
Ornithine Aminotransferase Deficiency	<i>OAT</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 6,900
Ornithine Transcarbamylase Deficiency	<i>OTC</i>	XL	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 103,000
Osteogenesis Imperfecta, Type XI	<i>FKBP10</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 8,100
Osteopetrosis 1	<i>TCIRG1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 5,700
Osteopetrosis 8	<i>SNX10</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 16,000
Otospondylomegaepiphyseal Dysplasia / Deafness / Fibrochondrogenesis 2	<i>COL11A2</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 2,700
Papillon-Lefevre Syndrome	<i>CTSC</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 5,000
Pendred Syndrome	<i>SLC26A4</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 72
Peroxisome Biogenesis Disorder 3A and 3B	<i>PEX12</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 30,000
Peroxisome Biogenesis Disorder 7A and 7B	<i>PEX26</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 23,000
Phenylalanine Hydroxylase Deficiency	<i>PAH</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 150
Polycystic Kidney Disease, Autosomal Recessive	<i>PKHD1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 350
Polyglandular Autoimmune Syndrome, Type 1	<i>AIRE</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 4,100
Pontocerebellar Hypoplasia, Type 1A	<i>VRK1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 25,000
Pontocerebellar Hypoplasia, Type 1B	<i>EXOSC3</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 10,000
Pontocerebellar Hypoplasia, Type 2A and Type 4	<i>TSEN54</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 4,700
Pontocerebellar Hypoplasia, Type 2E	<i>VPS53</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 146,000
Pontocerebellar Hypoplasia, Type 6	<i>RARS2</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 8,600
Primary Carnitine Deficiency	<i>SLC22A5</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 600
Primary Ciliary Dyskinesia ( <i>CCDC103</i> -Related)	<i>CCDC103</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 33,000
Primary Ciliary Dyskinesia ( <i>CCDC151</i> -Related)	<i>CCDC151</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 59,000
Primary Ciliary Dyskinesia ( <i>CCDC39</i> -Related)	<i>CCDC39</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 12,000
Primary Ciliary Dyskinesia ( <i>DNAH5</i> -Related)	<i>DNAH5</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,500
Primary Ciliary Dyskinesia ( <i>DNAI1</i> -Related)	<i>DNAI1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 5,000
Primary Ciliary Dyskinesia ( <i>DNAI2</i> -Related)	<i>DNAI2</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 76,000
Primary Ciliary Dyskinesia ( <i>RSPH9</i> -Related)	<i>RSPH9</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 73,000
Primary Coenzyme Q10 Deficiency 7	<i>COQ4</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 12,000
Primary Congenital Glaucoma 3A	<i>CYP1B1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 130
Primary Hyperoxaluria, Type 1	<i>AGXT</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,900
Primary Hyperoxaluria, Type 2	<i>GRHPR</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 11,000
Primary Hyperoxaluria, Type 3	<i>HOGA1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 2,400
Progressive Cerebello-Cerebral Atrophy	<i>SEPSECS</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 6,400
Progressive Familial Intrahepatic Cholestasis, Type 2	<i>ABCB11</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 390
Progressive Myoclonic Epilepsy, Type 1B	<i>PRICKLE1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 98,000
Progressive Pseudorheumatoid Dysplasia	<i>WISP3</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 5,600
Prolidase Deficiency	<i>PEPD</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 4,900
Propionic Acidemia ( <i>PCCA</i> -Related)	<i>PCCA</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 2,600
Propionic Acidemia ( <i>PCCB</i> -Related)	<i>PCCB</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 920
Pulmonary Surfactant Dysfunction	<i>ABCA3</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,200
Pycnodysostosis	<i>CTSK</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 5,100
Pyridoxamine 5'-Phosphate Oxidase Deficiency	<i>PNPO</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 6,600
Pyridoxine-Dependent Epilepsy	<i>ALDH7A1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,100
Pyruvate Carboxylase Deficiency	<i>PC</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 8,000
Pyruvate Dehydrogenase E1-Alpha Deficiency	<i>PDHA1</i>	XL	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 139,000

Pyruvate Dehydrogenase E1-Beta Deficiency	<i>PDHB</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 8,300
Renal Tubular Acidosis and Deafness	<i>ATP6V1B1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 7,800
Retinitis Pigmentosa 25	<i>EYS</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 580
Retinitis Pigmentosa 26	<i>CERKL</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 4,000
Retinitis Pigmentosa 28	<i>FAM161A</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 34,000
Retinitis Pigmentosa 36	<i>PRCD</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 304,000
Retinitis Pigmentosa 59	<i>DHDDS</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 201,000
Retinitis Pigmentosa 64 / Bardet-Biedl Syndrome 21 / Cone-Rod Dystrophy 16	<i>C8ORF37</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 4,300
Rh Deficiency Syndrome	<i>RHAG</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 46,000
Rhizomelic Chondrodysplasia Punctata, Type 1	<i>PEX7</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 10,000
Rhizomelic Chondrodysplasia Punctata, Type 3	<i>AGPS</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 620,000
Roberts Syndrome	<i>ESCO2</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 95,000
Salla Disease	<i>SLC17A5</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 8,400
Salt and Pepper Developmental Regression Syndrome	<i>ST3GAL5</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 25,000
Sandhoff Disease	<i>HEXB</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 680
Schimke Immunoosseous Dysplasia	<i>SMARCAL1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 3,800
Seckel Syndrome 5 / Microcephaly 9	<i>CEP152</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,500
Segawa Syndrome	<i>TH</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 3,000
Sepiapterin Reductase Deficiency	<i>SPR</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 43,000
Severe Combined Immunodeficiency ( <i>IL7R</i> -Related)	<i>IL7R</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 20,000
Severe Combined Immunodeficiency ( <i>JAK3</i> -Related)	<i>JAK3</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 2,100
Severe Combined Immunodeficiency ( <i>PTPRC</i> -Related)	<i>PTPRC</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 2,300
Severe Congenital Neutropenia 4	<i>G6PC3</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 10,000
Severe Neonatal Hyperparathyroidism	<i>CASR</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 2,700
Short Stature, Onychodysplasia, Facial Dysmorphism, and Hypotrichosis	<i>POC1A</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 98,000
Short-Chain Acyl-CoA Dehydrogenase Deficiency	<i>ACADS</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 340
Shwachman-Diamond Syndrome	<i>SBDS</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,200
Sialidosis, Type I and Type II	<i>NEU1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 2,000
Sjogren-Larsson Syndrome	<i>ALDH3A2</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 4100
Smith-Lemli-Opitz Syndrome	<i>DHCR7</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 750
Spastic Paraplegia 15	<i>ZFYVE26</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,600
Spastic Tetraplegia, Thin Corpus Callosum, and Progressive Microcephaly	<i>SLC1A4</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 136,000
Spherocytosis, Type 5	<i>EPB42</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,300
Spinal Muscular Atrophy	<i>SMN1</i>	AR	Reduced Risk	SMN1 copy number: 2 SMN2 copy number: 2 c.73+80T>G: Negative SMN1 Sequencing: Negative <b>Personalized Residual Risk:</b> 1 in 1,115
Spinal Muscular Atrophy with Respiratory Distress 1 / Charcot-Marie-Tooth Disease, Type 2S	<i>IGHMBP2</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,200
Spinocerebellar Ataxia with Axonal Neuropathy 3	<i>COA7</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 12,000
Spondylocostal Dysostosis 1	<i>DLL3</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 7,200
Spondylometaepiphyseal Dysplasia ( <i>DDR2</i> -Related)	<i>DDR2</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 220,000
Spondylothoracic Dysostosis	<i>MESP2</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 53,000
Steel Syndrome	<i>COL27A1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 93,000
Stuve-Wiedemann Syndrome	<i>LIFR</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 6,000

<b>Sulfate Transporter-Related Osteochondrodysplasia</b>	<i>SLC26A2</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,800 Tay-Sachs disease enzyme: Non-carrier White blood cells: Non-carrier <ul style="list-style-type: none"> <li>Hex A%: 57.9% (Non-carrier : 55.0 - 72.0%; Carrier: &lt;50%)</li> <li>Total hexosaminidase activity: 1844 nmol/hr/mg</li> </ul>
<b>Tay-Sachs Disease</b>	<i>HEXA</i>	AR	Reduced Risk	Plasma: Non-carrier <ul style="list-style-type: none"> <li>Hex A%: 71.7 (Non-carrier : 58.0 - 72.0%; Carrier: &lt;54%)</li> <li>Total hexosaminidase activity: 416 nmol/hr/ml</li> </ul> <i>HEXA</i> Sequencing: Negative <b>Personalized Residual Risk:</b> 1 in 1,400
<b>Thiamine-Responsive Megaloblastic Anemia Syndrome</b>	<i>SLC19A2</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 11,000
<b>Thyroid Dysmorphogenesis 1</b>	<i>SLC5A5</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 2,300
<b>Thyroid Dysmorphogenesis 2A</b>	<i>TPO</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 350
<b>Thyroid Dysmorphogenesis 3</b>	<i>TG</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 130
<b>Thyroid Dysmorphogenesis 4</b>	<i>IYD</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,800
<b>Thyroid Dysmorphogenesis 5</b>	<i>DUOXA2</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,300
<b>Thyroid Dysmorphogenesis 6</b>	<i>DUOX2</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 32
<b>Trichohepatoenteric Syndrome 1</b>	<i>TTC37</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 11,000
<b>Tyrosinemia, Type I</b>	<i>FAH</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,900
<b>Tyrosinemia, Type II</b>	<i>TAT</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 4,200
<b>Tyrosinemia, Type III</b>	<i>HPD</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 15,000
<b>Usher Syndrome, Type IB</b>	<i>MYO7A</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 180
<b>Usher Syndrome, Type IC</b>	<i>USH1C</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 400
<b>Usher Syndrome, Type ID</b>	<i>CDH23</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 880
<b>Usher Syndrome, Type IF</b>	<i>PCDH15</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,100
<b>Usher Syndrome, Type IIA</b>	<i>USH2A</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 54
<b>Usher Syndrome, Type III</b>	<i>CLRN1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,300
<b>Very Long Chain Acyl-CoA Dehydrogenase Deficiency</b>	<i>ACADVL</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 380
<b>Vitamin D-Dependent Rickets, Type I</b>	<i>CYP27B1</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,600
<b>Vitamin D-Resistant Rickets, Type IIA</b>	<i>VDR</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 21,000
<b>Walker-Warburg Syndrome and Other <i>FKTN</i>-Related Dystrophies</b>	<i>FKTN</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 390
<b>Werner Syndrome</b>	<i>WRN</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 2,000
<b>Wilson Disease</b>	<i>ATP7B</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 150
<b>Wiskott-Aldrich Syndrome (<i>WAS</i>-Related)</b>	<i>WAS</i>	XL	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 1,203,000
<b>Wolcott-Rallison Syndrome</b>	<i>EIF2AK3</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 22,000
<b>Wolman Disease / Cholesteryl Ester Storage Disease</b>	<i>LIPA</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 3,200
<b>Woodhouse-Sakati Syndrome</b>	<i>DCAF17</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 59,000
<b>X-Linked Juvenile Retinoschisis</b>	<i>RS1</i>	XL	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 40,000
<b>X-Linked Severe Combined Immunodeficiency</b>	<i>IL2RG</i>	XL	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 250,000
<b>Xeroderma Pigmentosum (<i>POLH</i>-Related)</b>	<i>POLH</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 4,300
<b>Xeroderma Pigmentosum, Group A</b>	<i>XPA</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 11,000
<b>Xeroderma Pigmentosum, Group C</b>	<i>XPC</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 12,000
<b>Xeroderma Pigmentosum, Group G</b>	<i>ERCC5</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 3,000
<b>Zellweger Syndrome Spectrum (<i>PEX10</i>-Related)</b>	<i>PEX10</i>	AR	Reduced Risk	<b>Personalized Residual Risk:</b> 1 in 6,300

<b>Zellweger Syndrome Spectrum (PEX1-Related)</b>	<i>PEX1</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 740</b>
<b>Zellweger Syndrome Spectrum (PEX2-Related)</b>	<i>PEX2</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 77,000</b>
<b>Zellweger Syndrome Spectrum (PEX6-Related)</b>	<i>PEX6</i>	AR	Reduced Risk	<b>Personalized Residual Risk: 1 in 1,500</b>

AR=Autosomal recessive; XL=X-linked

## Test methods and comments

Genomic DNA isolated from this patient was analyzed by one or more of the following methodologies, as applicable:

### Fragile X CGG Repeat Analysis (Analytical Detection Rate >99%)

PCR amplification using Asuragen, Inc. AmpliX<sup>®</sup> *FMR1* PCR reagents followed by capillary electrophoresis for allele sizing was performed. Samples positive for *FMR1* CGG repeats in the premutation and full mutation size range were further analyzed by Southern blot analysis to assess the size and methylation status of the *FMR1* CGG repeat.

### Genotyping (Analytical Detection Rate >99%)

Multiplex PCR amplification and allele specific primer extension analyses using the MassARRAY<sup>®</sup> System were used to identify certain recurrent variants that are complex in nature or are present in low copy repeats. Rare sequence variants may interfere with assay performance.

### Multiplex Ligation-Dependent Probe Amplification (MLPA) (Analytical Detection Rate >99%)

MLPA<sup>®</sup> probe sets and reagents from MRC-Holland were used for copy number analysis of specific targets versus known control samples. False positive or negative results may occur due to rare sequence variants in target regions detected by MLPA probes. Analytical sensitivity and specificity of the MLPA method are both 99%.

For alpha thalassemia, the copy numbers of the *HBA1* and *HBA2* genes were analyzed. Alpha-globin gene deletions, triplications, and the Constant Spring (CS) mutation are assessed. This test is expected to detect approximately 90% of all alpha-thalassemia mutations, varying by ethnicity, carriers of alpha-thalassemia with three or more *HBA* copies on one chromosome, and one or no copies on the other chromosome, may not be detected. With the exception of triplications, other benign alpha-globin gene polymorphisms will not be reported. Analyses of *HBA1* and *HBA2* are performed in association with long-range PCR of the coding regions followed by short-read sequencing.

For Duchenne muscular dystrophy, the copy numbers of all *DMD* exons were analyzed. Potentially pathogenic single exon deletions and duplications are confirmed by a second method. Analysis of *DMD* is performed in association with sequencing of the coding regions.

For congenital adrenal hyperplasia, the copy number of the *CYP21A2* gene was analyzed. This analysis can detect large deletions typically due to unequal meiotic crossing-over between *CYP21A2* and the pseudogene *CYP21A1P*. Classic 30-kb deletions make up approximately 20% of *CYP21A2* pathogenic alleles. This test may also identify certain point mutations in *CYP21A2* caused by gene conversion events between *CYP21A2* and *CYP21A1P*. Some carriers may not be identified by dosage sensitive methods as this testing cannot detect individuals with two copies (duplication) of the *CYP21A2* gene on one chromosome and loss of *CYP21A2* (deletion) on the other chromosome. Analysis of *CYP21A2* is performed in association with long-range PCR of the coding regions followed by short-read sequencing.

For spinal muscular atrophy (SMA), the copy numbers of the *SMN1* and *SMN2* genes were analyzed. The individual dosage of exons 7 and 8 as well as the combined dosage of exons 1, 4, 6 and 8 of *SMN1* and *SMN2* were assessed. Copy number gains and losses can be detected with this assay. Depending on ethnicity, 6 - 29 % of carriers will not be identified by dosage sensitive methods as this testing cannot detect individuals with two copies (duplication) of the *SMN1* gene on one chromosome and loss of *SMN1* (deletion) on the other chromosome (silent 2+0 carrier) or individuals that carry an intragenic mutation in *SMN1*. Please also note that 2% of individuals diagnosed with SMA have a causative *SMN1* variant that occurred *de novo*, and therefore cannot be picked up by carrier screening in the parents. Analysis of *SMN1* is performed in association with short-read sequencing of exons 2a-7, followed by confirmation using long-range PCR (described below).

The presence of the c.\*3+80T>G (chr5:70,247,901T>G) variant allele in an individual with Ashkenazi Jewish or Asian ancestry is typically indicative of a duplication of *SMN1*. When present in an Ashkenazi Jewish or Asian individual with two copies of *SMN1*, c.\*3+80T>G is likely indicative of a silent (2+0) carrier. In individuals with two copies of *SMN1* with African American, Hispanic or Caucasian ancestry, the presence or absence of c.\*3+80T>G significantly increases or decreases, respectively, the likelihood of being a silent 2+0 carrier.

MLPA for Gaucher disease (*GBA*), cystic fibrosis (*CFTR*), and non-syndromic hearing loss (*GJB2/GJB6*) will only be performed if indicated for confirmation of detected CNVs. If *GBA* analysis was performed, the copy numbers of exons 1, 3, 4, and 6 - 10 of the *GBA* gene (of 11 exons total) were analyzed. If *CFTR* analysis was performed, the copy numbers of all 27 *CFTR* exons were analyzed. If *GJB2/GJB6* analysis was performed, the copy number of the two *GJB2* exons were analyzed, as well as the presence or absence of the two upstream deletions of the *GJB2* regulatory region, del(*GJB6*-D13S1830) and del(*GJB6*-D13S1854).

### Next Generation Sequencing (NGS) (Analytical Detection Rate >95%)



NGS was performed on a panel of genes for the purpose of identifying pathogenic or likely pathogenic variants.

Agilent SureSelect<sup>TM</sup>XT Low Input technology was used with a custom capture library to target the exonic regions and intron/exon splice junctions of the relevant genes, as well as a number of UTR, intronic or promoter regions that contain previously reported mutations. Libraries were pooled and sequenced on the Illumina NovaSeq 9000 platform, using paired-end 100 bp reads. The sequencing data was analyzed using a custom bioinformatics algorithm designed and validated in house.

The coding exons and splice junctions of the known protein-coding RefSeq genes were assessed for the average depth of coverage (minimum of 20X) and data quality threshold values. Most exons not meeting a minimum of >20X read depth across the exon are further analyzed by Sanger sequencing. Please note that several genomic regions present difficulties in mapping or obtaining read depth >20X. These regions, which are described below, will not be reflexed to Sanger sequencing if the mapping quality or coverage is poor. Any variants identified during testing in these regions are confirmed by a second method and reported if determined to be pathogenic or likely pathogenic. However, as there is a possibility of false negative results within these regions, detection rates and residual risks for these genes have been calculated with the presumption that variants in these exons will not be detected, unless included in the MassARRAY<sup>®</sup> genotyping platform.

**Exceptions:** *ABCD1* (NM\_000033.3) exons 8 and 9; *ACADSB* (NM\_001609.3) chr10:124,810,695-124,810,707 (partial exon 9); *ADA* (NM\_000022.2) exon 1; *ADAMTS2* (NM\_014244.4) exon 1; *AGPS* (NM\_003659.3) chr2:178,257,512-178,257,649 (partial exon 1); *ALDH7A1* (NM\_001182.4) chr5:125,911,150-125,911,163 (partial exon 7) and chr5:125,896,807-125,896,821 (partial exon 10); *ALMS1* (NM\_015120.4) chr2:73,612,990-73,613,041 (partial exon 1); *APOPT1* (NM\_032374.4) chr14:104,040,437-104,040,455 (partial exon 3); *CDAN1* (NM\_138477.2) exon 2; *CEP152* (NM\_014985.3) chr15:49,061,146-49,061,165 (partial exon 14) and exon 22; *CEP290* (NM\_025114.3) exon 5, exon 7, chr12:88,519,017-88,519,039 (partial exon 13), chr12:88,514,049-88,514,058 (partial exon 15), chr12:88,502,837-88,502,841 (partial exon 23), chr12:88,481,551-88,481,589 (partial exon 32), chr12:88,471,605-88,471,700 (partial exon 40); *CFTR* (NM\_000492.3) exon 10; *COL4A4* (NM\_000092.4) chr2:227,942,604-227,942,619 (partial exon 25); *COX10* (NM\_001303.3) exon 6; *CYP11B1* (NM\_000497.3) exons 3-7; *CYP11B2* (NM\_000498.3) exons 3-7; *DNAI2* (NM\_023036.4) chr17:72,308,136-72,308,147 (partial exon 12); *DOK7* (NM\_173660.4) chr4:3,465,131-3,465,161 (partial exon 1) and exon 2; *DUOX2* (NM\_014080.4) exons 6-8; *EIF2AK3* (NM\_004836.5) exon 8; *EVC* (NM\_153717.2) exon 1; *FH* (NM\_000143.3) exon 1; *GAMT* (NM\_000156.5) exon 1; *GLDC* (NM\_000170.2) exon 1; *GNPTAB* (NM\_024312.4) chr17:4,837,000-4,837,400 (partial exon 2); *GNPTG* (NM\_032520.4) exon 1; *GHR* (NM\_000163.4) exon 3; *GYS2* (NM\_021957.3) chr12:21,699,370-21,699,409 (partial exon 12); *HGSNAT* (NM\_152419.2) exon 1; *IDS* (NM\_000202.6) exon 3; *ITGB4* (NM\_000213.4) chr17:73,749,976-73,750,060 (partial exon 33); *JAK3* (NM\_000215.3) chr19:17,950,462-17,950,483 (partial exon 10); *LIFR* (NM\_002310.5) exon 19; *LMBRD1* (NM\_018368.3) chr6:70,459,226-70,459,257 (partial exon 5), chr6:70,447,828-70,447,836 (partial exon 7) and exon 12; *LYST* (NM\_000081.3) chr1:235,944,158-235,944,176 (partial exon 16) and chr1:235,875,350-235,875,362 (partial exon 43); *MLYCD* (NM\_012213.2) chr16:83,933,242-83,933,282 (partial exon 1); *MTR* (NM\_000254.2) chr1:237,024,418-237,024,439 (partial exon 20) and chr1:237,038,019-237,038,029 (partial exon 24); *NBEAL2* (NM\_015175.2) chr3:47,021,385-47,021,407 (partial exon 1); *NEB* (NM\_001271208.1) exons 82-105; *NPC1* (NM\_000271.4) chr18:21,123,519-21,123,538 (partial exon 14); *NPHP1* (NM\_000272.3) chr2:110,937,251-110,937,263 (partial exon 3); *OCRL* (NM\_000276.3) chrX:128,674,450-128,674,460 (partial exon 1); *PHKB* (NM\_000293.2) exon 1 and chr16:47,732,498-47,732,504 (partial exon 30); *PIGN* (NM\_176787.4) chr18:59,815,547-59,815,576 (partial exon 8); *PIP5K1C* (NM\_012398.2) exon 1 and chr19:3637602-3637616 (partial exon 17); *POU1F1* (NM\_000306.3) exon 5; *PTPRC* (NM\_002838.4) exons 11 and 23; *PUS1* (NM\_025215.5) chr12:132,414,446-132,414,532 (partial exon 2); *RPGRIP1L* (NM\_015272.2) exon 23; *SGSH* (NM\_000199.3) chr17:78,194,022-78,194,072 (partial exon 1); *SLC6A8* (NM\_005629.3) exons 3 and 4; *ST3GAL5* (NM\_003896.3) exon 1; *SURF1* (NM\_003172.3) chr9:136,223,269-136,223,307 (partial exon 1); *TRPM6* (NM\_017662.4) chr9:77,362,800-77,362,811 (partial exon 31); *TSEN54* (NM\_207346.2) exon 1; *TYR* (NM\_000372.4) exon 5; *VWF* (NM\_000552.3) exons 24-26, chr12:6,125,675-6,125,684 (partial exon 30), chr12:6,121,244-6,121,265 (partial exon 33), and exon 34.

This test will detect variants within the exons and the intron-exon boundaries of the target regions. Variants outside these regions may not be detected, including, but not limited to, UTRs, promoters, and deep intronic areas, or regions that fall into the Exceptions mentioned above. This technology may not detect all small insertion/deletions and is not diagnostic for repeat expansions and structural genomic variation. In addition, a mutation(s) in a gene not included on the panel could be present in this patient.

Variant interpretation and classification was performed based on the American College of Medical Genetics Standards and Guidelines for the Interpretation of Sequence Variants (Richards et al, 2015). All potentially pathogenic variants may be confirmed by either a specific genotyping assay or Sanger sequencing, if indicated. Any benign variants, likely benign variants or variants of uncertain significance identified during this analysis will not be reported.

#### Next Generation Sequencing for SMN1

Exonic regions and intron/exon splice junctions of *SMN1* and *SMN2* were captured, sequenced, and analyzed as described above. Any variants located within exons 2a-7 and classified as pathogenic or likely pathogenic were confirmed to be in either *SMN1* or *SMN2* using gene-specific long-range PCR analysis followed by Sanger sequencing. Variants located in exon 1 cannot be accurately assigned to either *SMN1* or *SMN2* using our current methodology, and so these variants are considered to be of uncertain significance and are not reported.

#### Copy Number Variant Analysis (Analytical Detection Rate >95%)

Large duplications and deletions were called from the relative read depths on an exon-by-exon basis using a custom exome hidden Markov model (XHMM) algorithm. Deletions or duplications determined to be pathogenic or likely pathogenic were confirmed by either a custom arrayCGH platform, quantitative PCR, or MLPA (depending on CNV size and gene content). While this algorithm is designed to pick up deletions and duplications of 2 or more exons in length, potentially pathogenic single-exon CNVs will be confirmed and reported, if detected.

#### **Exon Array (Confirmation method) (Accuracy >99%)**

The customized oligonucleotide microarray (Oxford Gene Technology) is a highly-targeted exon-focused array capable of detecting medically relevant microdeletions and microduplications at a much higher resolution than traditional aCGH methods. Each array matrix has approximately 180,000 60-mer oligonucleotide probes that cover the entire genome. This platform is designed based on human genome NCBI Build 37 (hg19) and the CGH probes are enriched to target the exonic regions of the genes in this panel.

#### **Quantitative PCR (Confirmation method) (Accuracy >99%)**

Th relative quantification PCR is utilized on a Roche Universal Library Probe (UPL) system, which relates the PCR signal of the target region in one group to another. To test for genomic imbalances, both sample DNA and reference DNA is amplified with primer/probe sets that specific to the target region and a control region with known genomic copy number. Relative genomic copy numbers are calculated based on the standard  $\Delta\Delta C_t$  formula.

#### **Long-Range PCR (Analytical Detection Rate >99%)**

Long-range PCR was performed to generate locus-specific amplicons for *CYP21A2*, *HBA1* and *HBA2* and *GBA*. The PCR products were then prepared for short-read NGS sequencing and sequenced. Sequenced reads were mapped back to the original genomic locus and run through the bioinformatics pipeline. If indicated, copy number from MLPA was correlated with the sequencing output to analyze the results. For *CYP21A2*, a certain percentage of healthy individuals carry a duplication of the *CYP21A2* gene, which has no clinical consequences. In cases where two copies of a gene are located on the same chromosome in tandem, only the second copy will be amplified and assessed for potentially pathogenic variants, due to size limitations of the PCR reaction. However, because these alleles contain at least two copies of the *CYP21A2* gene in tandem, it is expected that this patient has at least one functional gene in the tandem allele and this patient is therefore less likely to be a carrier. When an individual carries both a duplication allele and a pathogenic variant, or multiple pathogenic variants, the current analysis may not be able to determine the phase (cis/trans configuration) of the *CYP21A2* alleles identified. Family studies may be required in certain scenarios where phasing is required to determine the carrier status.

#### **Residual Risk Calculations**

Carrier frequencies and detection rates for each ethnicity were calculated through the combination of internal curations of >30,000 variants and genomic frequency data from >138,000 individuals across seven ethnic groups in the gnomAD database. Additional variants in HGMD and novel deleterious variants were also incorporated into the calculation. Residual risk values are calculated using a Bayesian analysis combining the *a priori* risk of being a pathogenic mutation carrier (carrier frequency) and the detection rate. They are provided only as a guide for assessing approximate risk given a negative result, and values will vary based on the exact ethnic background of an individual. This report does not represent medical advice but should be interpreted by a genetic counselor, medical geneticist or physician skilled in genetic result interpretation and the relevant medical literature.

#### **Personalized Residual Risk Calculations**

Agilent SureSelect<sup>TM</sup>XT Low-Input technology was utilized in order to create whole-genome libraries for each patient sample. Libraries were then pooled and sequenced on the Illumina NovaSeq platform. Each sequencing lane was multiplexed to achieve 0.4-2x genome coverage, using paired-end 100 bp reads. The sequencing data underwent ancestral analysis using a customized, licensed bioinformatics algorithm that was validated in house. Identified sub-ethnic groupings were binned into one of 7 continental-level groups (African, East Asian, South Asian, Non-Finnish European, Finnish, Native American, and Ashkenazi Jewish) or, for those ethnicities that matched poorly to the continental-level groups, an 8<sup>th</sup> "unassigned" group, which were then used to select residual risk values for each gene. For individuals belonging to multiple high-level ethnic groupings, a weighting strategy was used to select the most appropriate residual risk. For genes that had insufficient data to calculate ethnic-specific residual risk values, or for sub-ethnic groupings that fell into the "unassigned" group, a "worldwide" residual risk was used. This "worldwide" residual risk was calculated using data from all available continental-level groups.

#### **Sanger Sequencing (Confirmation method) (Accuracy >99%)**

Sanger sequencing, as indicated, was performed using BigDye Terminator chemistry with the ABI 3730 DNA analyzer with target specific amplicons. It also may be used to supplement specific guaranteed target regions that fail NGS sequencing due to poor quality or low depth of coverage (<20 reads) or as a confirmatory method for NGS positive results. False negative results may occur if rare variants interfere with amplification or annealing.

#### **Tay-Sachs Disease (TSD) Enzyme Analysis (Analytical Detection Rate $\geq$ 98%)**

Hexosaminidase activity and Hex A% activity were measured by a standard heat-inactivation, fluorometric method using artificial 4-MU- $\beta$ -N-acetyl glucosaminide (4-MUG) substrate. This assay is highly sensitive and accurate in detecting Tay-Sachs carriers and individuals affected with TSD. Normal ranges of Hex A% activity are 55.0-72.0 for white blood cells and 58.0-72.0 for plasma. It is estimated that less than 0.5% of Tay-Sachs carriers have non-carrier levels of percent Hex A activity, and therefore may not be identified by this assay. In addition, this assay may detect individuals that are carriers of or are affected with Sandhoff disease. False positive results may occur if benign variants, such as pseudodeficiency alleles, interfere with the enzymatic assay. False negative results may occur if both *HEXA* and *HEXB* pathogenic or pseudodeficiency variants are present in the same individual.

Please note these tests were developed and their performance characteristics were determined by Sema4 Opco, Inc. They have not been cleared or approved by the FDA. These analyses generally provide highly accurate information regarding the patient's carrier or affected status. Despite this high level of accuracy, it should be kept in mind that there are many potential sources of diagnostic error, including misidentification of samples, polymorphisms, or other rare genetic variants that interfere with analysis. Families should understand that rare diagnostic errors may occur for these reasons.

## SELECTED REFERENCES

### Carrier Screening

Grody W et al. ACMG position statement on prenatal/preconception expanded carrier screening. *Genet Med*. 2013 15:482-3.

### Fragile X syndrome:

Chen L et al. An information-rich CGG repeat primed PCR that detects the full range of Fragile X expanded alleles and minimizes the need for Southern blot analysis. *J Mol Diag* 2010 12:589-600.

### Spinal Muscular Atrophy:

Luo M et al. An Ashkenazi Jewish SMN1 haplotype specific to duplication alleles improves pan-ethnic carrier screening for spinal muscular atrophy. *Genet Med*. 2014 16:149-56.

### Ashkenazi Jewish Disorders:

Scott SA et al. Experience with carrier screening and prenatal diagnosis for sixteen Ashkenazi Jewish Genetic Diseases. *Hum. Mutat*. 2010 31:1-11.

### Duchenne Muscular Dystrophy:

Flanigan KM et al. Mutational spectrum of DMD mutations in dystrophinopathy patients: application of modern diagnostic techniques to a large cohort. *Hum Mutat*. 2009 30:1657-66.

### Variant Classification:

Richards S et al. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med*. 2015 May;17(5):405-24

Additional disease-specific references available upon request.