



Donor 7414

Genetic Testing Summary

Fairfax Cryobank recommends reviewing this genetic testing summary with your healthcare provider to determine suitability.

Last Updated: 11/5/2025

Donor Reported Ancestry: Scottish

Jewish Ancestry: No

Genetic Test*	Result	Comments/ Donor's Residual Risk**
Chromosome analysis (karyotype)	Normal male karyotype	No evidence of clinically significant chromosome abnormalities
Hemoglobin evaluation	Normal hemoglobin fractionation and MCV/MCH results	Reduced risk to be a carrier for sickle cell anemia, beta thalassemia, alpha thalassemia trait (aa/-- and a-/a-) and other hemoglobinopathies
Expanded Genetic Disease Carrier Screening Panel attached- 549 diseases by gene sequencing.	<p>Carrier: Non-Syndromic Hearing Loss, GJB2-Related (GJB2)</p> <p>Carrier: Steroid-Resistant Nephrotic Syndrome (NPHS2)</p> <p>Carrier: Hereditary Hemochromatosis Type 1 (HFE)</p> <p>Negative for other genes sequenced.</p>	Partner testing is recommended before using this donor.

*No single test can screen for all genetic disorders. A negative screening result significantly reduces, but cannot eliminate, the risk for these conditions in a pregnancy.

**Donor residual risk is the chance the donor is still a carrier after testing negative.

Patient Information

Patient Name: Donor 7414
 Date Of Birth: [REDACTED]
 Gender: Male
 Ethnicity: Other
 Patient ID: [REDACTED]
 Medical Record #: N/A
 Collection Kit: [REDACTED]
 Accession ID: N/A
 Case File ID: [REDACTED]

Test Information

Ordering Physician: [REDACTED]
 Clinic Information: Fairfax Cryobank
 Phone: N/A
 Report Date: 04/01/2024
 Sample Collected: 03/18/2024
 Sample Received: 03/19/2024
 Sample Type: Blood

CARRIER SCREENING REPORT

ABOUT THIS SCREEN: Horizon™ is a carrier screen for specific autosomal recessive and X-linked diseases. This information can help patients learn their risk of having a child with specific genetic conditions.

ORDER SELECTED: The Horizon Custom panel was ordered for this patient. Males are not screened for X-linked diseases

FINAL RESULTS SUMMARY:

CARRIER for Hereditary Hemochromatosis Type 1

Positive for the pathogenic variant c.845G>A (p.C282Y) in the HFE gene. If this individual's partner is a carrier for HEREDITARY HEMOCHROMATOSIS TYPE 1, their chance to have a child with this condition is 1 in 4 (25%). Carrier screening for this individual's partner is suggested.

CARRIER for Non-Syndromic Hearing Loss, GJB2-Related

Positive for the pathogenic variant c.269T>C (p.L90P) in the GJB2 gene. Although most variants in this gene are associated with an autosomal recessive form of NON-SYNDROMIC HEARING LOSS, GJB2-RELATED, some rare GJB2 variants may cause an autosomal dominant form of the condition. To our knowledge, there is insufficient evidence that this variant causes an autosomal dominant form of this condition. If this individual's partner is a carrier for NON-SYNDROMIC HEARING LOSS, GJB2-RELATED, their chance to have a child with this condition is likely 1 in 4 (25%). Carrier screening for this individual's partner is suggested.

CARRIER for Steroid-Resistant Nephrotic Syndrome

Positive for the likely pathogenic variant c.686G>A (p.R229Q) in the NPHS2 gene. Please note the likely pathogenic variant c.686G>A, p.R229Q is a mild variant that is only expected to cause disease when found in trans with one of a specific set of variants that occurs in exons 7 or 8 (PMID: 30260545, 30241959). Homozygotes are not expected to be affected, unless this variant is part of a more complex allele (PMID: 24509478 and 29660491). If this individual's partner is a carrier for Steroid-Resistant Nephrotic Syndrome, their chance to have a child with this condition may be as high as 1 in 4 (25%). Carrier screening for this individual's partner is suggested.


Negative for 546 out of 549 diseases

No other pathogenic variants were detected in the genes that were screened. The patient's remaining carrier risk after the negative screening results is listed for each disease/gene on the Horizon website at <https://www.natera.com/panel-option/h-all/>. Please see the following pages of this report for a comprehensive list of all conditions included on this individual's screen.


Carrier screening is not diagnostic and may not detect all possible pathogenic variants in a given gene.

RECOMMENDATIONS


Individuals who would like to review their Horizon report with a Natera Laboratory Genetic Counselor may schedule a telephone genetic information session by calling 650-249-9090 or visiting naterasession.com. Clinicians with questions may contact Natera at 650-249-9090 or email support@natera.com. Individuals with positive results may wish to discuss these results with family members to allow them the option to be screened. Comprehensive genetic counseling to discuss the implications of these test results and possible associated reproductive risk is recommended.




Christine M. Eng, M.D.
 Medical Director, Baylor Genetics



Linyan Meng, Ph.D.
 Laboratory Director, Baylor Genetics



J. Dianne Keen-Kim, Ph.D., FACMG
 Senior Laboratory Director, Natera



Yang Wang, Ph.D., FACMG
 Laboratory Director, Natera

Patient Information

Patient Name: Donor 7414

Test Information

Ordering Physician: [REDACTED]



Date Of Birth: [REDACTED]

Case File ID: [REDACTED]

Clinic Information: Fairfax Cryobank

Report Date: 04/01/2024

HEREDITARY HEMOCHROMATOSIS TYPE 1

Understanding Your Horizon Carrier Screen Results

What does being a carrier mean?

Your results show that you are a carrier of hereditary hemochromatosis type 1. A carrier of a genetic condition does not have the condition. Carriers also are not certain to have a child with the condition. We are all carriers of one or more genetic conditions.

Your children are not at high risk for this condition unless your partner or donor is also a carrier of hereditary hemochromatosis type 1. Further testing can be done to see if your partner or donor is a carrier.

What is hereditary hemochromatosis type 1?

Hereditary hemochromatosis type 1 causes the body to absorb too much iron from food. Many people with this condition have iron levels that never get high enough to impact their health. A small number of people with hereditary hemochromatosis type 1 will have health problems. Their skin, heart, pancreas, joints, or liver can be damaged. People with hereditary hemochromatosis type 1 usually start having symptoms as adults. Males are more likely to have symptoms than females. Early symptoms can include joint pain, fatigue, weight loss, and abdominal pain. Symptoms can get worse over time as more iron is stored. Later symptoms can include discolored skin, arthritis, diabetes, liver disease, and heart disease. Symptoms can be impacted by diet and alcohol use. Hemochromatosis can be treated with regular phlebotomy (blood removal) to remove excess iron. Treatment can lead to fewer symptoms. People diagnosed and treated prior to getting liver disease have a normal life expectancy.^{1,2}

Clinical trials involving potential new treatments for this condition could be available (see clinicaltrials.gov).

What causes hereditary hemochromatosis type 1?

Hereditary hemochromatosis type 1 is caused by changes, or variants, in the HFE gene. These changes make the gene not work properly. Genes are a set of instructions inside the cells of our bodies that tell our bodies how to grow and function. Everyone has two copies of the HFE gene. Carriers of hereditary hemochromatosis type 1 have one working copy and one non-working copy of the gene. People with hereditary hemochromatosis type 1 have no working copies of the gene.

Hereditary hemochromatosis type 1 is usually passed down, or inherited, from both genetic parents. We inherit one copy of the HFE gene from each of our genetic parents. When both genetic parents are carriers, each child has a 1 in 4 (25%) chance of inheriting two non-working genes and having hereditary hemochromatosis type 1. Each child also has a 1 in 2 (50%) chance of being a carrier of hereditary hemochromatosis type 1 and a 1 in 4 (25%) chance of inheriting two working copies of the gene. This type of inheritance is called autosomal recessive inheritance.

Will my children have hereditary hemochromatosis type 1?

If your partner or donor also has a non-working copy of the HFE gene, your children could have hereditary hemochromatosis type 1. Each child you have together would have a 1 in 4 (25%) chance of having hereditary hemochromatosis type 1. Each child you have together would also have a 3 in 4 (75%) chance of **not** having the condition.

If your partner or donor has HFE carrier screening and no variants are found, the chance that your children would have hereditary hemochromatosis type 1 is very low. No further testing would usually be needed for you, your partner or donor, or your children related to hereditary hemochromatosis type 1.

What can I do next?

Carriers of hereditary hemochromatosis type 1 can have elevated iron levels in their blood, but they do not have symptoms of hemochromatosis.² Talk to your healthcare provider.

If you want to know if your children are at risk for hereditary hemochromatosis type 1, your partner or donor would need to have HFE carrier screening. If you have questions about this testing, please ask your healthcare provider or use the resources below. Many people find it helpful to speak with a genetic counselor.

If your partner or donor is found to be a hereditary hemochromatosis type 1 carrier, your children would be at risk for having hereditary hemochromatosis type 1. Some variants are more likely to cause symptoms than others. If your partner or donor has an HFE variant, genetic counseling to discuss how likely it is your children would have symptoms is recommended.

If you or your partner or surrogate are currently pregnant, tests called CVS (chorionic villus sampling) and amniocentesis can be done during pregnancy to find out if a baby has hereditary hemochromatosis type 1. These tests both have a small risk of miscarriage. Babies can also be tested for hereditary hemochromatosis type 1 after birth instead.

If you or your partner or surrogate are not yet pregnant, you could have these options:

- natural pregnancy with CVS or amniocentesis to test for hereditary hemochromatosis type 1 during pregnancy;
- natural pregnancy and testing the baby after birth for hereditary hemochromatosis type 1;
- preimplantation genetic testing (PGT-M) with in vitro fertilization (IVF) to test embryos for hereditary hemochromatosis type 1;
- adoption; or
- use of a sperm or egg donor who had no variants found in HFE carrier screening.

Where can I find more information?

- Genetic and Rare Diseases Information Center rarediseases.info.nih.gov/diseases/10746/hemochromatosis
- CVS marchofdimes.org/chorionic-villus-sampling

Patient Information

Patient Name: [REDACTED]

Test Information

Ordering Physician: [REDACTED]



Date Of Birth: [REDACTED]

Clinic Information:

Case File ID: [REDACTED]

Report Date:

- Amniocentesis marchofdimes.org/pregnancy/amniocentesis
- PGT-M natera.com/womens-health/spectrum-preimplantation-genetics

What does this mean for my family?

You likely got (inherited) this non-working gene from one of your genetic parents. Your genetic siblings and other family members could also carry it. You should tell your family members about your test results so they can decide if they want carrier screening for hereditary hemochromatosis type 1.

References

1. MedlinePlus [Internet]. Bethesda (MD): National Library of Medicine (US); [updated 2020 Jun 24]. Hereditary hemochromatosis; [updated 2019 Feb 1; cited 2024 Feb 1]; Available from: <https://medlineplus.gov/genetics/condition/hereditary-hemochromatosis/#synonyms>
2. Barton JC, Edwards CQ. HFE Hemochromatosis. 2000 Apr 3 [Updated 2018 Dec 6]. In: Adam MP et al, editors. GeneReviews® [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2024. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK1440/>

Patient Information

Patient Name: [REDACTED]

Test Information

Ordering Physician: [REDACTED]



Clinic Information: [REDACTED]

Date Of Birth: [REDACTED]

Case File ID: [REDACTED]

Report Date: [REDACTED]

NON-SYNDROMIC HEARING LOSS, GJB2-RELATED**Understanding Your Horizon Carrier Screen Results****What is Non-Syndromic Hearing Loss, GJB2-Related?**

Non-Syndromic Hearing Loss, GJB2-Related (also called DFNB1) is an inherited disorder that causes early-onset hearing loss. "Non-syndromic" means that no other parts of the body are affected, making hearing loss the only symptom of this condition. In Non-Syndromic Hearing Loss, GJB2-Related, hearing loss is typically present at birth (congenital). However, some children have normal hearing at birth and develop hearing loss during childhood. The severity varies from mild to profound sensorineural hearing loss. The treatment for hearing loss includes hearing aids and, in some cases, cochlear implants. Clinical trials involving potential new treatments for this condition may be available (see www.clinicaltrials.gov). Non-Syndromic Hearing Loss, GJB2-Related does not cause other health problems.

What causes Non-Syndromic Hearing Loss, GJB2-Related?

Non-Syndromic Hearing Loss, GJB2-Related is caused by a gene change, or mutation, in both copies of the GJB2 gene pair (also known as DFNB1). These mutations cause the genes to not work properly or not work at all. The function of the GJB2 genes is to make a protein that is important for hearing. When both copies of the GJB2 gene do not work correctly, it leads to Non-Syndromic Hearing Loss, GJB2-Related. Non-Syndromic Hearing Loss, GJB2-Related is inherited in an autosomal recessive manner. This means that, in most cases, both parents must be carriers of a mutation in one copy of the GJB2 gene to have a child with Non-Syndromic Hearing Loss, GJB2-Related. People who are carriers for Non-Syndromic Hearing Loss, GJB2-Related are usually healthy and usually do not have Non-Syndromic Hearing Loss themselves. Usually a child inherits two copies of each gene, one copy from the mother and one copy from the father. If the mother and father are both carriers for Non-Syndromic Hearing Loss, GJB2-Related, there is a 1 in 4, or 25%, chance in each pregnancy for both partners to pass on their GJB2 gene mutations to the child, who will then have Non-Syndromic Hearing Loss, GJB2-Related. Very rarely, carriers of a single GJB2 mutation will have inherited hearing loss with or without other symptoms. These individuals usually have one parent who is also affected. This type of inheritance, where having only one mutation causes symptoms, is called autosomal dominant. When a person with autosomal dominant hearing loss has a child, there is a 50%, or 1 in 2, chance with each pregnancy of having a child who will also develop this type of hearing loss. It is sometimes, but not always, possible to determine whether a specific mutation in the GJB2 gene will cause autosomal recessive Non-Syndromic Hearing Loss or an autosomal dominant type of hearing loss. Individuals found to carry more than one mutation for Non-Syndromic Hearing Loss, GJB2-Related should discuss their risk for having an affected child and any potential effects to their own hearing with their health care provider.

What can I do next?

You may wish to speak with a local genetic counselor about your carrier test results. A genetic counselor in your area can be located on the National Society of Genetic Counselors website (www.nsgc.org). You may wish to share your carrier screening results with your health care providers, especially if you have a family history of hearing loss or have concerns about your own hearing. Your siblings and other relatives are at increased risk to also have this mutation. You are encouraged to inform your family members of your test results as they may wish to consider being tested themselves. If you are pregnant, your partner can have carrier screening for Non-Syndromic Hearing Loss, GJB2-Related ordered by a health care professional. If your partner is not found to be a carrier for Non-Syndromic Hearing Loss, GJB2-Related, your risk of having a child with Non-Syndromic Hearing Loss, GJB2-Related is greatly reduced. Couples at risk of having a baby with Non-Syndromic Hearing Loss, GJB2-Related can opt to have prenatal diagnosis done through chorionic villus sampling (CVS) or amniocentesis during pregnancy or can choose to test the baby after birth for this condition. If you are not yet pregnant, your partner can have carrier screening for Non-Syndromic Hearing Loss, GJB2-Related ordered by a health care professional. If your partner is found to be a carrier for Non-Syndromic Hearing Loss, GJB2-Related, you have several reproductive options to consider:

- Natural pregnancy with or without prenatal diagnosis of the fetus or testing the baby after birth for Non-Syndromic Hearing Loss, GJB2-Related
- Preimplantation genetic diagnosis (PGD) with in vitro fertilization (IVF) to test embryos for Non-Syndromic Hearing Loss, GJB2-Related
- Adoption or use of a sperm or egg donor who is not a carrier for Non-Syndromic Hearing Loss, GJB2-Related

What resources are available?

- Genetics Home Reference: <http://ghr.nlm.nih.gov/condition/nonsyndromic-hearing-loss>
- Prenatal diagnosis done through CVS: <http://www.marchofdimes.org/chorionic-villus-sampling.aspx>
- Prenatal diagnosis done through Amniocentesis: <http://www.marchofdimes.org/amniocentesis.aspx>
- Preimplantation genetic diagnosis (PGD) with IVF: <http://www.natera.com/spectrum>

Patient Information

Patient Name: [REDACTED]

Test Information

Ordering Physician: [REDACTED]



Clinic Information: [REDACTED]

Date Of Birth: [REDACTED]

Case File ID: [REDACTED]

Report Date: [REDACTED]

STEROID-RESISTANT NEPHROTIC SYNDROME**Understanding Your Horizon Carrier Screen Results****What is Steroid-Resistant Nephrotic Syndrome?**

Steroid-Resistant Nephrotic Syndrome (also known as Familial Nephrotic Syndrome or Nephrotic Syndrome, Type 2) is an inherited disorder that causes abnormal kidney function. People with this condition have large amounts of protein in their urine, low amounts of albumin (a protein in the plasma of the blood), high levels of fat in the blood, and excess fluid in body tissues (edema). Symptoms vary from person to person and usually start in infancy or childhood, although some people do not show symptoms until early adulthood. The kidney problems worsen over time, often leading to kidney failure in childhood, the teenage years, or early adulthood. Once kidney failure occurs, dialysis and then kidney transplantation are needed. Currently there is no cure for this condition and treatment is based on symptoms. Clinical trials involving potential new treatments for this condition may be available (see www.clinicaltrials.gov).

What causes Steroid-Resistant Nephrotic Syndrome?

Steroid-Resistant Nephrotic Syndrome is caused by a gene change, or mutation, in both copies of the NPHS2 gene pair. These mutations cause the genes to not work properly or not work at all. When both copies of this gene do not work correctly, it leads to the symptoms described above. Steroid-Resistant Nephrotic Syndrome is inherited in an autosomal recessive manner. This means that, in most cases, both parents must be carriers of a mutation in one copy of the NPHS2 gene to have a child with Steroid-Resistant Nephrotic Syndrome. People who are carriers for Steroid-Resistant Nephrotic Syndrome are usually healthy and do not have symptoms nor do they have Steroid-Resistant Nephrotic Syndrome themselves. Usually a child inherits two copies of each gene, one copy from the mother and one copy from the father. If the mother and father are both carriers for Steroid-Resistant Nephrotic Syndrome, there is a 1 in 4, or 25%, chance in each pregnancy for both partners to pass on their NPHS2 gene mutations to the child, who would then have this condition. Individuals found to carry more than one mutation for Steroid-Resistant Nephrotic Syndrome should discuss any potential effects to their own health and their risk for having an affected child with their health care provider.

What can I do next?

You may wish to speak with a local genetic counselor about your carrier test results. A genetic counselor in your area can be located on the National Society of Genetic Counselors website (www.nsgc.org). Your siblings and other relatives are at increased risk to also have this mutation. You are encouraged to inform your family members of your test results as they may wish to consider being tested themselves. If you are pregnant, your partner can have carrier screening for Steroid-Resistant Nephrotic Syndrome ordered by a health care professional. If your partner is not found to be a carrier for Steroid-Resistant Nephrotic Syndrome, your risk of having a child with this condition is greatly reduced. Couples at risk of having a baby with Steroid-Resistant Nephrotic Syndrome can opt to have prenatal diagnosis done through chorionic villus sampling (CVS) or amniocentesis during pregnancy or can choose to have the baby tested after birth for this condition. If you are not yet pregnant, your partner can have carrier screening for Steroid-Resistant Nephrotic Syndrome ordered by a health care professional. If your partner is found to be a carrier for Steroid-Resistant Nephrotic Syndrome, you have several reproductive options to consider:

- Natural pregnancy with or without prenatal diagnosis of the fetus or testing the baby after birth for Steroid-Resistant Nephrotic Syndrome
- Preimplantation genetic diagnosis (PGD) with in vitro fertilization (IVF) to test embryos for Steroid-Resistant Nephrotic Syndrome
- Adoption or use of a sperm or egg donor who is not a carrier for Steroid-Resistant Nephrotic Syndrome

What resources are available?

- The Renal Association: www.rarerrenal.org
- Genetics Home Reference: <http://ghr.nlm.nih.gov/gene/NPHS2>
- Prenatal diagnosis done through CVS: <http://www.marchofdimes.org/chorionic-villus-sampling.aspx>
- Prenatal diagnosis done through Amniocentesis: <http://www.marchofdimes.org/amniocentesis.aspx>
- PGD with IVF: <http://www.natera.com/spectrum>

Patient Information

Patient Name:

Test Information

Ordering Physician:



Clinic Information:

Date Of Birth:

Case File ID:

Report Date:

DISEASES SCREENED

Below is a list of all diseases screened and the result. Certain conditions have unique patient-specific numerical values, therefore, results for those conditions are formatted differently.

Autosomal Recessive

- 1**
17-BETA HYDROXYSTEROID DEHYDROGENASE 3 DEFICIENCY (*HSD17B3*) **negative**
- 3**
3-BETA-HYDROXYSTEROID DEHYDROGENASE TYPE II DEFICIENCY (*HSD3B2*) **negative**
3-HYDROXY-3-METHYLGLUTARYL-COENZYME A LYASE DEFICIENCY (*HMGCL*) **negative**
3-HYDROXYACYL-COA DEHYDROGENASE DEFICIENCY (*HADH*) **negative**
3-METHYLCROTONYL-CoA CARBOXYLASE 2 DEFICIENCY (*MCCC2*) **negative**
3-PHOSPHOGLYCERATE DEHYDROGENASE DEFICIENCY (*PHGDH*) **negative**
- 5**
5-ALPHA-REDUCTASE DEFICIENCY (*SRD5A2*) **negative**
- 6**
6-PYRUVYL-TETRAHYDROPTERIN SYNTHASE (*PTPS*) DEFICIENCY (*PTS*) **negative**
- A**
ABCA4-RELATED CONDITIONS (*ABCA4*) **negative**
ABETALIPOPROTEINEMIA (*MTTP*) **negative**
ACHONDROGENESIS, TYPE 1B (*SLC26A2*) **negative**
ACHROMATOPSIA, CNGB3-RELATED (*CNGB3*) **negative**
ACRODERMATITIS ENTEROPATHICA (*SLC39A4*) **negative**
ACTION MYOCLONUS-RENAL FAILURE (AMRF) SYNDROME (*SCARB2*) **negative**
ACUTE INFANTILE LIVER FAILURE, TRMU-RELATED (*TRMU*) **negative**
ACYL-COA OXIDASE I DEFICIENCY (*ACOX1*) **negative**
AICARDI-GOUTIÈRES SYNDROME (*SAMHD1*) **negative**
AICARDI-GOUTIÈRES SYNDROME, RNASEH2A-RELATED (*RNASEH2A*) **negative**
AICARDI-GOUTIÈRES SYNDROME, RNASEH2B-RELATED (*RNASEH2B*) **negative**
AICARDI-GOUTIÈRES SYNDROME, RNASEH2C-RELATED (*RNASEH2C*) **negative**
AICARDI-GOUTIÈRES SYNDROME, TREX1-RELATED (*TREX1*) **negative**
ALPHA-MANNOSIDOSIS (*MAN2B1*) **negative**
ALPHA-THALASSEMIA (*HBA1/HBA2*) **negative**
ALPORT SYNDROME, COL4A3-RELATED (*COL4A3*) **negative**
ALPORT SYNDROME, COL4A4-RELATED (*COL4A4*) **negative**
ALSTROM SYNDROME (*ALMS1*) **negative**
AMISH INFANTILE EPILEPSY SYNDROME (*ST3GAL5*) **negative**
ANDERMANN SYNDROME (*SLC12A6*) **negative**
ARGININE:GLYCINE AMIDINOTRANSFERASE DEFICIENCY (AGAT DEFICIENCY) (*GATM*) **negative**
ARGININEMIA (*ARG1*) **negative**
ARGININOSUCCINATE LYASE DEFICIENCY (*ASL*) **negative**
AROMATASE DEFICIENCY (*CYP19A1*) **negative**
ASPARAGINE SYNTHETASE DEFICIENCY (*ASNS*) **negative**
ASPARTYLGLYCOSAMINURIA (*AGA*) **negative**
ATAXIA WITH VITAMIN E DEFICIENCY (*TTPA*) **negative**
ATAXIA-TELANGIECTASIA (*ATM*) **negative**
ATAXIA-TELANGIECTASIA-LIKE DISORDER 1 (*MRE11*) **negative**
ATANSFERRINEMIA (*TF*) **negative**
AUTISM SPECTRUM, EPILEPSY AND ARTHROGRYPOSIS (*SLC35A3*) **negative**
AUTOIMMUNE POLYGLANDULAR SYNDROME, TYPE 1 (*AIRE*) **negative**
AUTOSOMAL RECESSIVE CONGENITAL ICHTHYOSIS (*ARCI*), *SLC27A4*-RELATED (*SLC27A4*) **negative**
AUTOSOMAL RECESSIVE SPASTIC ATAXIA OF CHARLEVOIX-SAGUENAY (*SACS*) **negative**
- B**
BARDET-BIEDL SYNDROME, ARL6-RELATED (*ARL6*) **negative**
BARDET-BIEDL SYNDROME, BBS10-RELATED (*BBS10*) **negative**
BARDET-BIEDL SYNDROME, BBS12-RELATED (*BBS12*) **negative**
BARDET-BIEDL SYNDROME, BBS1-RELATED (*BBS1*) **negative**
BARDET-BIEDL SYNDROME, BBS2-RELATED (*BBS2*) **negative**
BARDET-BIEDL SYNDROME, BBS4-RELATED (*BBS4*) **negative**
BARDET-BIEDL SYNDROME, BBS5-RELATED (*BBS5*) **negative**
BARDET-BIEDL SYNDROME, BBS7-RELATED (*BBS7*) **negative**
BARDET-BIEDL SYNDROME, BBS9-RELATED (*BBS9*) **negative**
BARDET-BIEDL SYNDROME, TTC8-RELATED (*TTC8*) **negative**
BARE LYMPHOCYTE SYNDROME, CIITA-RELATED (*CIITA*) **negative**
BARTTER SYNDROME, BSND-RELATED (*BSND*) **negative**
BARTTER SYNDROME, KCNJ1-RELATED (*KCNJ1*) **negative**
BARTTER SYNDROME, SLC12A1-RELATED (*SLC12A1*) **negative**
BATTEN DISEASE, CLN3-RELATED (*CLN3*) **negative**
BETA-HEMOGLOBINOPATHIES (*HBB*) **negative**
BETA-KETOTHIOLASE DEFICIENCY (*ACAT1*) **negative**
BETA-MANNOSIDOSIS (*MANBA*) **negative**
BETA-UREIDOPROPIONASE DEFICIENCY (*UPB1*) **negative**
BILATERAL FRONTOPARIETAL POLYMICROGYRIA (*GPR56*) **negative**
- BIOTINIDASE DEFICIENCY (*BTD*) **negative**
BIOTIN-THIAMINE-RESPONSIVE BASAL GANGLIA DISEASE (BTBGD) (*SLC19A3*) **negative**
BLOOM SYNDROME (*BLM*) **negative**
BRITTLE CORNEA SYNDROME 1 (*ZNF469*) **negative**
BRITTLE CORNEA SYNDROME 2 (*PRDM5*) **negative**
- C**
CANAVAN DISEASE (*ASPA*) **negative**
CARBAMOYL PHOSPHATE SYNTHETASE I DEFICIENCY (*CPS1*) **negative**
CARNITINE DEFICIENCY (*SLC22A5*) **negative**
CARNITINE PALMITOYLTRANSFERASE IA DEFICIENCY (*CPT1A*) **negative**
CARNITINE PALMITOYLTRANSFERASE II DEFICIENCY (*CPT2*) **negative**
CARNITINE-ACYLCARNITINE TRANSLOCASE DEFICIENCY (*SLC25A20*) **negative**
CARPENTER SYNDROME (*RAB23*) **negative**
CARTILAGE-HAIR HYPOPLASIA (*RMRP*) **negative**
CATECHOLAMINERGIC POLYMORPHIC VENTRICULAR TACHYCARDIA (*CASQ2*) **negative**
CD59-MEDIATED HEMOLYTIC ANEMIA (*CD59*) **negative**
CEP152-RELATED MICROCEPHALY (*CEP152*) **negative**
CEREBRAL DYSGENESIS, NEUROPATHY, ICHTHYOSIS, AND PALMOPLANTAR KERATODERMA (CEDNIK) SYNDROME (*SNAP29*) **negative**
CEREBROTENDINOUS XANTHOMATOSIS (*CYP27A1*) **negative**
CHARCOT-MARIE-TOOTH DISEASE, RECESSIVE INTERMEDIATE C (*PLEKHG5*) **negative**
CHARCOT-MARIE-TOOTH-DISEASE, TYPE 4D (*NDRG1*) **negative**
CHEDIAK-HIGASHI SYNDROME (*LYST*) **negative**
CHOREOACANTHOCYTOSIS (*VPS13A*) **negative**
CHRONIC GRANULOMATOUS DISEASE, CYBA-RELATED (*CYBA*) **negative**
CHRONIC GRANULOMATOUS DISEASE, NCF2-RELATED (*NCF2*) **negative**
CILIOPATHIES, RPGRIP1L-RELATED (*RPGRIP1L*) **negative**
CITRIN DEFICIENCY (*SLC25A13*) **negative**
CITRULLINEMIA, TYPE 1 (*ASS1*) **negative**
CLN10 DISEASE (*CTSD*) **negative**
COHEN SYNDROME (*VPS13B*) **negative**
COL11A2-RELATED CONDITIONS (*COL11A2*) **negative**
COMBINED MALONIC AND METHYLMALONIC ACIDURIA (*ACSF3*) **negative**
COMBINED OXIDATIVE PHOSPHORYLATION DEFICIENCY 1 (*GMF1*) **negative**
COMBINED OXIDATIVE PHOSPHORYLATION DEFICIENCY 3 (*TSFM*) **negative**
COMBINED PITUITARY HORMONE DEFICIENCY 1 (*POU1F1*) **negative**
COMBINED PITUITARY HORMONE DEFICIENCY-2 (*PROP1*) **negative**
CONGENITAL ADRENAL HYPERPLASIA, 11-BETA-HYDROXYLASE DEFICIENCY (*CYP11B1*) **negative**
CONGENITAL ADRENAL HYPERPLASIA, 17-ALPHA-HYDROXYLASE DEFICIENCY (*CYP17A1*) **negative**
CONGENITAL ADRENAL HYPERPLASIA, 21-HYDROXYLASE DEFICIENCY (*CYP21A2*) **negative**
CONGENITAL ADRENAL INSUFFICIENCY, CYP11A1-RELATED (*CYP11A1*) **negative**
CONGENITAL AMEGAKARYOCYTIC THROMBOCYTOPENIA (*MPL*) **negative**
CONGENITAL CHRONIC DIARRHEA (*DGAT1*) **negative**
CONGENITAL DISORDER OF GLYCOSYLATION TYPE 1, ALG1-RELATED (*ALG1*) **negative**
CONGENITAL DISORDER OF GLYCOSYLATION, TYPE 1A, PMM2-Related (*PMM2*) **negative**
CONGENITAL DISORDER OF GLYCOSYLATION, TYPE 1B (*MPI*) **negative**
CONGENITAL DISORDER OF GLYCOSYLATION, TYPE 1C (*ALG6*) **negative**
CONGENITAL DYSERYTHROPOIETIC ANEMIA TYPE 2 (*SEC23B*) **negative**
CONGENITAL FINNISH NEPHROSIS (*NPHS1*) **negative**
CONGENITAL HYDROCEPHALUS 1 (*CCDC88C*) **negative**
CONGENITAL HYPERINSULINISM, KCNJ11-Related (*KCNJ11*) **negative**
CONGENITAL INSENSITIVITY TO PAIN WITH ANHIDROSIS (CIPA) (*NTRK1*) **negative**
CONGENITAL MYASTHENIC SYNDROME, CHAT-RELATED (*CHAT*) **negative**
CONGENITAL MYASTHENIC SYNDROME, CHRNE-RELATED (*CHRNE*) **negative**
CONGENITAL MYASTHENIC SYNDROME, COLQ-RELATED (*COLQ*) **negative**
CONGENITAL MYASTHENIC SYNDROME, DOK7-RELATED (*DOK7*) **negative**
CONGENITAL MYASTHENIC SYNDROME, RAPSN-RELATED (*RAPSN*) **negative**
CONGENITAL NEPHROTIC SYNDROME, PLCE1-RELATED (*PLCE1*) **negative**
CONGENITAL NEUTROPENIA, G6PC3-RELATED (*G6PC3*) **negative**
CONGENITAL NEUTROPENIA, HAX1-RELATED (*HAX1*) **negative**
CONGENITAL NEUTROPENIA, VPS45-RELATED (*VPS45*) **negative**
CONGENITAL SECRETORY CHLORIDE DIARRHEA 1 (*SLC26A3*) **negative**
CORNEAL DYSTROPHY AND PERCEPTIVE DEAFNESS (*SLC4A11*) **negative**
CORTICOSTERONE METHYLOXIDASE DEFICIENCY (*CYP11B2*) **negative**
COSTEFF SYNDROME (3-METHYLGLOUTACONIC ACIDURIA, TYPE 3) (*OPA3*) **negative**
CRB1-RELATED RETINAL DYSTROPHIES (*CRB1*) **negative**
CYSTIC FIBROSIS (*CFTR*) **negative**
CYSTINOSIS (*CTNS*) **negative**
CYTOCHROME C OXIDASE DEFICIENCY, PET100-RELATED (*PET100*) **negative**
CYTOCHROME P450 OXIOREDUCTASE DEFICIENCY (*POR*) **negative**
- D**
D-BIFUNCTIONAL PROTEIN DEFICIENCY (*HSD17B4*) **negative**

Patient Information

Patient Name:

Test Information

Ordering Physician:



Clinic Information:

Date Of Birth:

Case File ID:

Report Date:

D

DEAFNESS, AUTOSOMAL RECESSIVE 77 (LOXHD1) **negative**
 DIHYDROPTERIDINE REDUCTASE (DHPR) DEFICIENCY (QDPR) **negative**
 DONNAI-BARROW SYNDROME (LRP2) **negative**
 DUBIN-JOHNSON SYNDROME (ABCC2) **negative**
 DYSKERATOSIS CONGENITA SPECTRUM DISORDERS (TERT) **negative**
 DYSKERATOSIS CONGENITA, RTKL1-RELATED (RTKL1) **negative**
 DYSTROPHIC EPIDERMOLYSIS BULLOSA, COL7A1-Related (COL7A1) **negative**

E

EARLY INFANTILE EPILEPTIC ENCEPHALOPATHY, CAD-RELATED (CAD) **negative**
 EHLERS-DANLOS SYNDROME TYPE VI (PLOD1) **negative**
 EHLERS-DANLOS SYNDROME, CLASSIC-LIKE, TNXB-RELATED (TNXB) **negative**
 EHLERS-DANLOS SYNDROME, TYPE VII C (ADAMTS2) **negative**
 ELLIS-VAN CREVELD SYNDROME, EVC2-RELATED (EVC2) **negative**
 ELLIS-VAN CREVELD SYNDROME, EVC-RELATED (EVC) **negative**
 ENHANCED S-CONE SYNDROME (NR2E3) **negative**
 EPIMERASE DEFICIENCY (GALACTOSEMIA TYPE III) (GALE) **negative**
 EPIPHYSEAL DYSPLASIA, MULTIPLE, 7/DESBUQUOIS DYSPLASIA 1 (CANT1) **negative**
 ERCC6-RELATED DISORDERS (ERCC6) **negative**
 ERCC8-RELATED DISORDERS (ERCC8) **negative**
 ETHYLMALONIC ENCEPHALOPATHY (ETHE1) **negative**

F

FACTOR XI DEFICIENCY (F11) **negative**
 FAMILIAL DYSAUTONOMIA (IKBKAP) **negative**
 FAMILIAL HEMOPHAGOCYTIC LYMPHOHISTIOCYTOSIS, PRF1-RELATED (PRF1) **negative**
 FAMILIAL HEMOPHAGOCYTIC LYMPHOHISTIOCYTOSIS, STX11-RELATED (STX11) **negative**
 FAMILIAL HEMOPHAGOCYTIC LYMPHOHISTIOCYTOSIS, STXB2-RELATED (STXB2) **negative**
 FAMILIAL HEMOPHAGOCYTIC LYMPHOHISTIOCYTOSIS, UNC13D-RELATED (UNC13D) **negative**
 FAMILIAL HYPERCHOLESTEROLEMIA, LDLRAP1-RELATED (LDLRAP1) **negative**
 FAMILIAL HYPERCHOLESTEROLEMIA, LDLR-RELATED (LDLR) **negative**
 FAMILIAL HYPERINSULINISM, ABCC8-RELATED (ABCC8) **negative**
 FAMILIAL NEPHROGENIC DIABETES INSIPIDUS, AQP2-RELATED (AQP2) **negative**
 FANCONI ANEMIA, GROUP A (FANCA) **negative**
 FANCONI ANEMIA, GROUP C (FANCC) **negative**
 FANCONI ANEMIA, GROUP D2 (FANCD2) **negative**
 FANCONI ANEMIA, GROUP E (FANCE) **negative**
 FANCONI ANEMIA, GROUP F (FANCF) **negative**
 FANCONI ANEMIA, GROUP G (FANCG) **negative**
 FANCONI ANEMIA, GROUP I (FANCI) **negative**
 FANCONI ANEMIA, GROUP J (BRIP1) **negative**
 FANCONI ANEMIA, GROUP L (FANCL) **negative**
 FARBER LIPOGRANULOMATOSIS (ASH1) **negative**
 FOVEAL HYPOPLASIA (SLC38A8) **negative**
 FRASER SYNDROME 3, GRIP1-RELATED (GRIP1) **negative**
 FRASER SYNDROME, FRAS1-RELATED (FRAS1) **negative**
 FRASER SYNDROME, FREM2-RELATED (FREM2) **negative**
 FRIEDREICH ATAXIA (FXN) **negative**
 FRUCTOSE-1,6-BISPHOSPHATASE DEFICIENCY (FBP1) **negative**
 FUCOSIDOSIS, FUCA1-RELATED (FUCA1) **negative**
 FUMARASE DEFICIENCY (FH) **negative**

G

GABA-TRANSAMINASE DEFICIENCY (ABAT) **negative**
 GALACTOKINASE DEFICIENCY (GALACTOSEMIA, TYPE II) (GALK1) **negative**
 GALACTOSEMIA (GALT) **negative**
 GALACTOSIALIDOSIS (CTSA) **negative**
 GAUCHER DISEASE (GBA) **negative**
 GCH1-RELATED CONDITIONS (GCH1) **negative**
 GDF5-RELATED CONDITIONS (GDF5) **negative**
 GERODERMA OSTEODYSPLASTICA (GORAB) **negative**
 GITELMAN SYNDROME (SLC12A3) **negative**
 GLANZMANN THROMBASTHENIA (ITGB3) **negative**
 GLUTARIC ACIDEMIA, TYPE 1 (GCDH) **negative**
 GLUTARIC ACIDEMIA, TYPE 2A (ETFA) **negative**
 GLUTARIC ACIDEMIA, TYPE 2B (ETFB) **negative**
 GLUTARIC ACIDEMIA, TYPE 2C (ETFDH) **negative**
 GLUTATHIONE SYNTHETASE DEFICIENCY (GSS) **negative**
 GLYCINE ENCEPHALOPATHY, AMT-RELATED (AMT) **negative**
 GLYCINE ENCEPHALOPATHY, GLDC-RELATED (GLDC) **negative**
 GLYCOGEN STORAGE DISEASE TYPE 5 (McArdle Disease) (PYGM) **negative**
 GLYCOGEN STORAGE DISEASE TYPE IXB (PHKB) **negative**
 GLYCOGEN STORAGE DISEASE TYPE IXC (PHKG2) **negative**
 GLYCOGEN STORAGE DISEASE, TYPE 1a (G6PC) **negative**
 GLYCOGEN STORAGE DISEASE, TYPE 1b (SLC37A4) **negative**
 GLYCOGEN STORAGE DISEASE, TYPE 2 (POMPE DISEASE) (GAA) **negative**
 GLYCOGEN STORAGE DISEASE, TYPE 3 (AGL) **negative**
 GLYCOGEN STORAGE DISEASE, TYPE 4 (GBE1) **negative**
 GLYCOGEN STORAGE DISEASE, TYPE 7 (PFKM) **negative**

GRACILE SYNDROME (BCS1L) negativeGUANIDINOACETATE METHYLTRANSFERASE DEFICIENCY (GAMT) **negative****H**

HARLEQUIN ICHTHYOSIS (ABCA12) **negative**
 HEME OXYGENASE 1 DEFICIENCY (HMOX1) **negative**
 HEMOCHROMATOSIS, TYPE 3, TFR2-Related (TFR2) **negative**
 HEPATOCEREBRAL MITOCHONDRIAL DNA DEPLETION SYNDROME, MPV17-RELATED (MPV17) **negative**
 HEREDITARY FRUCTOSE INTOLERANCE (ALDOB) **negative**
 HEREDITARY HEMOCHROMATOSIS TYPE 1 (HFE) **see first page**
 HEREDITARY HEMOCHROMATOSIS TYPE 2B (HAMP) **negative**
 HEREDITARY SPASTIC PARAPARESIS, TYPE 49 (TECPR2) **negative**
 HEREDITARY SPASTIC PARAPLEGIA, CYP7B1-RELATED (CYP7B1) **negative**
 HERMANSKY-PUDLAK SYNDROME, AP3B1-RELATED (AP3B1) **negative**
 HERMANSKY-PUDLAK SYNDROME, BLOC1S3-RELATED (BLOC1S3) **negative**
 HERMANSKY-PUDLAK SYNDROME, BLOC1S6-RELATED (BLOC1S6) **negative**
 HERMANSKY-PUDLAK SYNDROME, HPS1-RELATED (HPS1) **negative**
 HERMANSKY-PUDLAK SYNDROME, HPS3-RELATED (HPS3) **negative**
 HERMANSKY-PUDLAK SYNDROME, HPS4-RELATED (HPS4) **negative**
 HERMANSKY-PUDLAK SYNDROME, HPS5-RELATED (HPS5) **negative**
 HERMANSKY-PUDLAK SYNDROME, HPS6-RELATED (HPS6) **negative**
 HOLOCARBOXYLASE SYNTHETASE DEFICIENCY (HLCS) **negative**
 HOMOCYSTEINURIA AND MEGALOBlastic ANEMIA TYPE CBLG (MTR) **negative**
 HOMOCYSTEINURIA DUE TO DEFICIENCY OF MTHFR (MTHFR) **negative**
 HOMOCYSTEINURIA, CBS-RELATED (CBS) **negative**
 HOMOCYSTEINURIA, Type cblE (MTRR) **negative**
 HYDROLETHALUS SYNDROME (HYLS1) **negative**
 HYPER-IGM IMMUNODEFICIENCY (CD40) **negative**
 HYPERORNITHINEMIA-HYPERAMMONEMIA-HOMOCITRULLINURIA (HHH SYNDROME) (SLC25A15) **negative**
 HYPERPHOSPHATEMIC FAMILIAL TUMORAL CALCINOSIS, GALNT3-RELATED (GALNT3) **negative**
 HYPOMYELINATING LEUKODYSTROPHY 12 (VPS11) **negative**
 HYPOPHOSPHATASIA, ALPL-RELATED (ALPL) **negative**

I

IMERSLUND-GRÄSBECK SYNDROME 2 (AMN) **negative**
 IMMUNODEFICIENCY-CENTROMERIC INSTABILITY-FACIAL ANOMALIES (ICF) SYNDROME, DNMT3B-RELATED (DNMT3B) **negative**
 IMMUNODEFICIENCY-CENTROMERIC INSTABILITY-FACIAL ANOMALIES (ICF) SYNDROME, ZBTB24-RELATED (ZBTB24) **negative**
 INCLUSION BODY MYOPATHY 2 (GNE) **negative**
 INFANTILE CEREBRAL AND CEREBELLAR ATROPHY (MED17) **negative**
 INFANTILE NEPHRONOPHTHISIS (INVS) **negative**
 INFANTILE NEUROAXONAL DYSTROPHY (PLA2G6) **negative**
 ISOLATED ECTOPIA LENTIS (ADAMTSL4) **negative**
 ISOLATED SULFITE OXIDASE DEFICIENCY (SUOX) **negative**
 ISOLATED THYROID-STIMULATING HORMONE DEFICIENCY (TSHB) **negative**
 ISOVALERIC ACIDEMIA (IVD) **negative**

J

JOHANSON-BLIZZARD SYNDROME (UBR1) **negative**
 JOUBERT SYNDROME 2 / MECKEL SYNDROME 2 (TMEM216) **negative**
 JOUBERT SYNDROME AND RELATED DISORDERS (JSRD), TMEM67-RELATED (TMEM67) **negative**
 JOUBERT SYNDROME, AH1-RELATED (AH1) **negative**
 JOUBERT SYNDROME, ARL13B-RELATED (ARL13B) **negative**
 JOUBERT SYNDROME, B9D1-RELATED (B9D1) **negative**
 JOUBERT SYNDROME, B9D2-RELATED (B9D2) **negative**
 JOUBERT SYNDROME, C2CD3-RELATED/OROFACIODIGITAL SYNDROME 14 (C2CD3) **negative**
 JOUBERT SYNDROME, CC2D2A-RELATED/COACH SYNDROME (CC2D2A) **negative**
 JOUBERT SYNDROME, CEP104-RELATED (CEP104) **negative**
 JOUBERT SYNDROME, CEP120-RELATED/SHORT-RIB THORACIC DYSPLASIA 13 WITH OR WITHOUT POLYDACTYLY (CEP120) **negative**
 JOUBERT SYNDROME, CEP41-RELATED (CEP41) **negative**
 JOUBERT SYNDROME, CPLANE1-RELATED / OROFACIODIGITAL SYNDROME 6 (CPLANE1) **negative**
 JOUBERT SYNDROME, CSPP1-RELATED (CSPP1) **negative**
 JOUBERT SYNDROME, INPP5E-RELATED (INPP5E) **negative**
 JUNCTIONAL EPIDERMOLYSIS BULLOSA, COL17A1-RELATED (COL17A1) **negative**
 JUNCTIONAL EPIDERMOLYSIS BULLOSA, ITGA6-RELATED (ITGA6) **negative**
 JUNCTIONAL EPIDERMOLYSIS BULLOSA, ITGB4-RELATED (ITGB4) **negative**
 JUNCTIONAL EPIDERMOLYSIS BULLOSA, LAMB3-RELATED (LAMB3) **negative**
 JUNCTIONAL EPIDERMOLYSIS BULLOSA, LAMC2-RELATED (LAMC2) **negative**
 JUNCTIONAL EPIDERMOLYSIS BULLOSA/LARYNGOONYCHOCUTANEOUS SYNDROME, LAMA3-RELATED (LAMA3) **negative**

KKRABBE DISEASE (GALC) **negative****L**LAMELLAR ICHTHYOSIS, TYPE 1 (TGM1) **negative**

Patient Information

Patient Name:

Test Information

Ordering Physician:



Clinic Information:

Date Of Birth:



Case File ID:



Report Date:

L
 LARON SYNDROME (GHR) **negative**
 LEBER CONGENITAL AMAUROSIS 2 (RPE65) **negative**
 LEBER CONGENITAL AMAUROSIS TYPE AIPL1 (AIPL1) **negative**
 LEBER CONGENITAL AMAUROSIS TYPE GUCY2D (GUCY2D) **negative**
 LEBER CONGENITAL AMAUROSIS TYPE TULP1 (TULP1) **negative**
 LEBER CONGENITAL AMAUROSIS, IQCB1-RELATED/SENIOR-LOKEN SYNDROME 5 (IQCB1) **negative**
 LEBER CONGENITAL AMAUROSIS, TYPE CEP290 (CEP290) **negative**
 LEBER CONGENITAL AMAUROSIS, TYPE LCA5 (LCA5) **negative**
 LEBER CONGENITAL AMAUROSIS, TYPE RDH12 (RDH12) **negative**
 LEIGH SYNDROME, FRENCH-CANADIAN TYPE (LRPPRC) **negative**
 LETHAL CONGENITAL CONTRACTURE SYNDROME 1 (GLE1) **negative**
 LEUKOENCEPHALOPATHY WITH VANISHING WHITE MATTER (EIF2B5) **negative**
 LEUKOENCEPHALOPATHY WITH VANISHING WHITE MATTER, EIF2B1-RELATED (EIF2B1) **negative**
 LEUKOENCEPHALOPATHY WITH VANISHING WHITE MATTER, EIF2B2-RELATED (EIF2B2) **negative**
 LEUKOENCEPHALOPATHY WITH VANISHING WHITE MATTER, EIF2B3-RELATED (EIF2B3) **negative**
 LEUKOENCEPHALOPATHY WITH VANISHING WHITE MATTER, EIF2B4-RELATED (EIF2B4) **negative**
 LIG4 SYNDROME (LIG4) **negative**
 LIMB-GIRDLE MUSCULAR DYSTROPHY TYPE 8 (TRIM32) **negative**
 LIMB-GIRDLE MUSCULAR DYSTROPHY, TYPE 2A (CAPN3) **negative**
 LIMB-GIRDLE MUSCULAR DYSTROPHY, TYPE 2B (DYSF) **negative**
 LIMB-GIRDLE MUSCULAR DYSTROPHY, TYPE 2C (SGCG) **negative**
 LIMB-GIRDLE MUSCULAR DYSTROPHY, TYPE 2D (SGCA) **negative**
 LIMB-GIRDLE MUSCULAR DYSTROPHY, TYPE 2E (SGCB) **negative**
 LIMB-GIRDLE MUSCULAR DYSTROPHY, TYPE 2F (SGCD) **negative**
 LIMB-GIRDLE MUSCULAR DYSTROPHY, TYPE 2I (FKRP) **negative**
 LIPOAMIDE DEHYDROGENASE DEFICIENCY (DIHYDROLIPOAMIDE DEHYDROGENASE DEFICIENCY) (DLI) **negative**
 LIPOID ADRENAL HYPERPLASIA (STAR) **negative**
 LIPOPROTEIN LIPASE DEFICIENCY (LPL) **negative**
 LONG CHAIN 3-HYDROXYACYL-CoA DEHYDROGENASE DEFICIENCY (HADHA) **negative**
 LRAT-RELATED CONDITIONS (LRAT) **negative**
 LUNG DISEASE, IMMUNODEFICIENCY, AND CHROMOSOME BREAKAGE SYNDROME (LICS) (NSMCE3) **negative**
 LYSINURIC PROTEIN INTOLERANCE (SLC7A7) **negative**

M
 MALONYL-CoA DECARBOXYLASE DEFICIENCY (MLYCD) **negative**
 MAPLE SYRUP URINE DISEASE, TYPE 1A (BCKDHA) **negative**
 MAPLE SYRUP URINE DISEASE, TYPE 1B (BCKDHB) **negative**
 MAPLE SYRUP URINE DISEASE, TYPE 2 (DBT) **negative**
 MCKUSICK-KAUFMAN SYNDROME (MKKS) **negative**
 MECKEL SYNDROME 7/NEPHRONOPHTHISIS 3 (NPHP3) **negative**
 MECKEL-GRUBER SYNDROME, TYPE 1 (MKS1) **negative**
 MECR-RELATED NEUROLOGIC DISORDER (MECR) **negative**
 MEDIUM CHAIN ACYL-CoA DEHYDROGENASE DEFICIENCY (ACADM) **negative**
 MEDNIK SYNDROME (AP1S1) **negative**
 MEGALENCEPHALIC LEUKOENCEPHALOPATHY WITH SUBCORTICAL CYSTS (MLC1) **negative**
 MEROSIN-DEFICIENT MUSCULAR DYSTROPHY (LAMA2) **negative**
 METABOLIC ENCEPHALOPATHY AND ARRHYTHMIAS, TANGO2-RELATED (TANGO2) **negative**
 METACHROMATIC LEUKODYSTROPHY, ARSA-RELATED (ARSA) **negative**
 METACHROMATIC LEUKODYSTROPHY, PSAP-RELATED (PSAP) **negative**
 METHYLMALONIC ACIDEMIA AND HOMOCYSTINURIA TYPE CBLF (LMBRD1) **negative**
 METHYLMALONIC ACIDEMIA, MCEE-RELATED (MCEE) **negative**
 METHYLMALONIC ACIDURIA AND HOMOCYSTINURIA, TYPE CBLF (MMACHC) **negative**
 METHYLMALONIC ACIDURIA AND HOMOCYSTINURIA, TYPE CblD (MMADHC) **negative**
 METHYLMALONIC ACIDURIA, MMAA-RELATED (MMAA) **negative**
 METHYLMALONIC ACIDURIA, MMAB-RELATED (MMAB) **negative**
 METHYLMALONIC ACIDURIA, TYPE MUT (0) (MUT) **negative**
 MEVALONIC KINASE DEFICIENCY (MVK) **negative**
 MICROCEPHALIC OSTEODYSPLASTIC PRIMORDIAL DWARFISM TYPE II (PCNT) **negative**
 MICROPHthalmia / ANOPHTHALMIA, VSX2-RELATED (VSX2) **negative**
 MITOCHONDRIAL COMPLEX 1 DEFICIENCY, ACAD9-RELATED (ACAD9) **negative**
 MITOCHONDRIAL COMPLEX 1 DEFICIENCY, NDUFAF5-RELATED (NDUFAF5) **negative**
 MITOCHONDRIAL COMPLEX 1 DEFICIENCY, NDUFS6-RELATED (NDUFS6) **negative**
 MITOCHONDRIAL COMPLEX I DEFICIENCY, NUCLEAR TYPE 1 (NDUFS4) **negative**
 MITOCHONDRIAL COMPLEX I DEFICIENCY, NUCLEAR TYPE 10 (NDUFAF2) **negative**
 MITOCHONDRIAL COMPLEX I DEFICIENCY, NUCLEAR TYPE 17 (NDUFAF6) **negative**
 MITOCHONDRIAL COMPLEX I DEFICIENCY, NUCLEAR TYPE 19 (FOXRED1) **negative**
 MITOCHONDRIAL COMPLEX I DEFICIENCY, NUCLEAR TYPE 3 (NDUFS7) **negative**
 MITOCHONDRIAL COMPLEX I DEFICIENCY, NUCLEAR TYPE 4 (NDUFV1) **negative**
 MITOCHONDRIAL COMPLEX IV DEFICIENCY, NUCLEAR TYPE 2, SCO2-RELATED (SCO2) **negative**
 MITOCHONDRIAL COMPLEX IV DEFICIENCY, NUCLEAR TYPE 6 (COX15) **negative**
 MITOCHONDRIAL DNA DEPLETION SYNDROME 2 (TK2) **negative**

MITOCHONDRIAL DNA DEPLETION SYNDROME 3 (DGUOK) **negative**
 MITOCHONDRIAL MYOPATHY AND SIDEROBLASTIC ANEMIA (MLASA1) (PUS1) **negative**
 MITOCHONDRIAL TRIFUNCTIONAL PROTEIN DEFICIENCY, HADHB-RELATED (HADHB) **negative**
 MOLYBDENUM COFACTOR DEFICIENCY TYPE B (MOCS2) **negative**
 MOLYBDENUM COFACTOR DEFICIENCY, TYPE A (MOCS1) **negative**
 MUCOLIPIDOSIS II/III A (GNPTAB) **negative**
 MUCOLIPIDOSIS III GAMMA (GNPTG) **negative**
 MUCOLIPIDOSIS, TYPE IV (MCOLN1) **negative**
 MUCOPOLYSACCHARIDOSIS, TYPE I (HURLER SYNDROME) (IDUA) **negative**
 MUCOPOLYSACCHARIDOSIS, TYPE III A (SANFILIPPO A) (SGSH) **negative**
 MUCOPOLYSACCHARIDOSIS, TYPE III B (SANFILIPPO B) (NAGLU) **negative**
 MUCOPOLYSACCHARIDOSIS, TYPE III C (SANFILIPPO C) (HGSNAT) **negative**
 MUCOPOLYSACCHARIDOSIS, TYPE III D (SANFILIPPO D) (GNS) **negative**
 MUCOPOLYSACCHARIDOSIS, TYPE IV A (MORQUIO SYNDROME) (GALNS) **negative**
 MUCOPOLYSACCHARIDOSIS, TYPE IV B/GM1 GANGLIOSIDOSIS (GLB1) **negative**
 MUCOPOLYSACCHARIDOSIS, TYPE IX (HYAL1) **negative**
 MUCOPOLYSACCHARIDOSIS, TYPE VI (MARQUET-LAMY) (ARSB) **negative**
 MUCOPOLYSACCHARIDOSIS, TYPE VII (GUSB) **negative**
 MULIBREY NANISM (TRIM37) **negative**
 MULTIPLE PTERYGIUM SYNDROME, CHRNG-RELATED/ESCOBAR SYNDROME (CHRNG) **negative**
 MULTIPLE SULFATASE DEFICIENCY (SUMF1) **negative**
 MUSCLE-EYE-BRAIN DISEASE, POMGNT1-RELATED (POMGNT1) **negative**
 MUSCULAR DYSTROPHY-DYSTROGLYCANOPATHY (RXYLT1) **negative**
 MUSK-RELATED CONGENITAL MYASTHENIC SYNDROME (MUSK) **negative**
 MYONEUROGASTROINTESTINAL ENCEPHALOPATHY (MNGIE) (TYMP) **negative**
 MYOTONIA CONGENITA (CLCN1) **negative**

N
 N-ACETYLGlutamate SYNTHASE DEFICIENCY (NAGS) **negative**
 NEMALINE MYOPATHY, NEB-RELATED (NEB) **negative**
 NEPHRONOPHTHISIS 1 (NPHP1) **negative**
 NEURONAL CEROID LIPOFUSCINOSIS, CLN5-RELATED (CLN5) **negative**
 NEURONAL CEROID LIPOFUSCINOSIS, CLN6-RELATED (CLN6) **negative**
 NEURONAL CEROID LIPOFUSCINOSIS, CLN8-RELATED (CLN8) **negative**
 NEURONAL CEROID LIPOFUSCINOSIS, MFSD8-RELATED (MFSD8) **negative**
 NEURONAL CEROID LIPOFUSCINOSIS, PPT1-RELATED (PPT1) **negative**
 NEURONAL CEROID LIPOFUSCINOSIS, TPP1-RELATED (TPP1) **negative**
 NGLY1-CONGENITAL DISORDER OF GLYCOSYLATION (NGLY1) **negative**
 NIEMANN-PICK DISEASE, TYPE C1 / D (NPC1) **negative**
 NIEMANN-PICK DISEASE, TYPE C2 (NPC2) **negative**
 NIEMANN-PICK DISEASE, TYPES A / B (SMPD1) **negative**
 NIJMEGEN BREAKAGE SYNDROME (NBN) **negative**
 NON-SYNDROMIC HEARING LOSS, GJB2-RELATED (GJB2) **see first page**
 NON-SYNDROMIC HEARING LOSS, MYO15A-RELATED (MYO15A) **negative**
 NONSYNDROMIC HEARING LOSS, OTOA-RELATED (OTOA) **negative**
 NONSYNDROMIC HEARING LOSS, OTOF-RELATED (OTOF) **negative**
 NONSYNDROMIC HEARING LOSS, PJK-RELATED (PJK) **negative**
 NONSYNDROMIC HEARING LOSS, SYNE4-RELATED (SYNE4) **negative**
 NONSYNDROMIC HEARING LOSS, TMC1-RELATED (TMC1) **negative**
 NONSYNDROMIC HEARING LOSS, TMRSS3-RELATED (TMRSS3) **negative**
 NONSYNDROMIC INTELLECTUAL DISABILITY (CC2D1A) **negative**
 NORMOPHOSPHATEMIC TUMORAL CALCINOSIS (SAMD9) **negative**

O
 OCULOCUTANEOUS ALBINISM TYPE IV (SLC45A2) **negative**
 OCULOCUTANEOUS ALBINISM TYPE, III (TYRP1) **negative**
 OCULOCUTANEOUS ALBINISM, OCA2-RELATED (OCA2) **negative**
 OCULOCUTANEOUS ALBINISM, TYPES 1A AND 1B (TYR) **negative**
 ODONTO-ONYCHO-DERMAL DYSPLASIA / SCHOPF-SCHULZ-PASSARGE SYNDROME (WNT10A) **negative**
 OMENN SYNDROME, RAG2-RELATED (RAG2) **negative**
 ORNITHINE AMINOTRANSFERASE DEFICIENCY (OAT) **negative**
 OSTEOGENESIS IMPERFECTA TYPE VII (CRTAP) **negative**
 OSTEOGENESIS IMPERFECTA TYPE VIII (P3H1) **negative**
 OSTEOGENESIS IMPERFECTA TYPE XI (FKBP10) **negative**
 OSTEOGENESIS IMPERFECTA TYPE XIII (BMP1) **negative**
 OSTEOPEETROSIS, INFANTILE MALIGNANT, TCIRG1-RELATED (TCIRG1) **negative**
 OSTEOPEETROSIS, OSTM1-RELATED (OSTM1) **negative**

P
 PANTOTHENATE KINASE-ASSOCIATED NEURODEGENERATION (PANK2) **negative**
 PAPILLON LEFÈVRE SYNDROME (CTSC) **negative**
 PARKINSON DISEASE 15 (FBXO7) **negative**
 PENDORF SYNDROME (SLC26A4) **negative**
 PERLMAN SYNDROME (DIS3L2) **negative**
 PGM3-CONGENITAL DISORDER OF GLYCOSYLATION (PGM3) **negative**
 PHENYLKETONURIA (PAH) **negative**
 PIGN-CONGENITAL DISORDER OF GLYCOSYLATION (PIGN) **negative**
 PITUITARY HORMONE DEFICIENCY, COMBINED 3 (LHX3) **negative**
 POLG-RELATED DISORDERS (POLG) **negative**

Patient Information

Patient Name: [REDACTED]

Test Information

Ordering Physician: [REDACTED]



Clinic Information: [REDACTED]

Date Of Birth: [REDACTED]

Case File ID: [REDACTED]

Report Date: [REDACTED]

P

POLYCYSTIC KIDNEY DISEASE, AUTOSOMAL RECESSIVE (*PKHD1*) **negative**
 PONTocerebellar Hypoplasia, *EXOSC3*-RELATED (*EXOSC3*) **negative**
 PONTocerebellar Hypoplasia, *RARS2*-RELATED (*RARS2*) **negative**
 PONTocerebellar Hypoplasia, *TSEN2*-RELATED (*TSEN2*) **negative**
 PONTocerebellar Hypoplasia, *TSEN54*-RELATED (*TSEN54*) **negative**
 PONTocerebellar Hypoplasia, TYPE 1A (*VRK1*) **negative**
 PONTocerebellar Hypoplasia, TYPE 2D (*SEPECS*) **negative**
 PONTocerebellar Hypoplasia, *VPS53*-RELATED (*VPS53*) **negative**
 PRIMARY CILIARY DYSKINESIA, *CCDC103*-RELATED (*CCDC103*) **negative**
 PRIMARY CILIARY DYSKINESIA, *CCDC39*-RELATED (*CCDC39*) **negative**
 PRIMARY CILIARY DYSKINESIA, *DNAH11*-RELATED (*DNAH11*) **negative**
 PRIMARY CILIARY DYSKINESIA, *DNAH5*-RELATED (*DNAH5*) **negative**
 PRIMARY CILIARY DYSKINESIA, *DNAI1*-RELATED (*DNAI1*) **negative**
 PRIMARY CILIARY DYSKINESIA, *DNAI2*-RELATED (*DNAI2*) **negative**
 PRIMARY CONGENITAL GLAUCOMA/PETERS ANOMALY (*CYP1B1*) **negative**
 PRIMARY HYPEROXALURIA, TYPE 1 (*AGXT*) **negative**
 PRIMARY HYPEROXALURIA, TYPE 2 (*GRHPR*) **negative**
 PRIMARY HYPEROXALURIA, TYPE 3 (*HOGA1*) **negative**
 PRIMARY MICROCEPHALY 1, AUTOSOMAL RECESSIVE (*MCPH1*) **negative**
 PROGRESSIVE EARLY-ONSET ENCEPHALOPATHY WITH BRAIN ATROPHY AND THIN CORPUS CALLOSUM (*TBCD*) **negative**
 PROGRESSIVE FAMILIAL INTRAHEPATIC CHOLESTASIS, *ABC4*-RELATED (*ABC4*) **negative**
 PROGRESSIVE FAMILIAL INTRAHEPATIC CHOLESTASIS, TYPE 1 (*PFIC1*) (*ATP8B1*) **negative**
 PROGRESSIVE FAMILIAL INTRAHEPATIC CHOLESTASIS, TYPE 2 (*ABC11*) **negative**
 PROGRESSIVE FAMILIAL INTRAHEPATIC CHOLESTASIS, TYPE 4 (*PFIC4*) (*TJP2*) **negative**
 PROGRESSIVE PSEUDORHEUMATOID DYSPLASIA (*CCN6*) **negative**
 PROLIDASE DEFICIENCY (*PEPD*) **negative**
 PROPIONIC ACIDEMIA, *PCCA*-RELATED (*PCCA*) **negative**
 PROPIONIC ACIDEMIA, *PCCB*-RELATED (*PCCB*) **negative**
 PSEUDOXANTHOMA ELASTICUM (*ABCC6*) **negative**
 PTERIN-4 ALPHA-CARBINOLAMINE DEHYDRATASE (*PCD*) DEFICIENCY (*PCBD1*) **negative**
 PYCNODYSTOSIS (*CTS1*) **negative**
 PYRIDOXAL 5'-PHOSPHATE-DEPENDENT EPILEPSY (*PNPO*) **negative**
 PYRIDOXINE-DEPENDENT EPILEPSY (*ALDH7A1*) **negative**
 PYRUVATE CARBOXYLASE DEFICIENCY (*PC*) **negative**
 PYRUVATE DEHYDROGENASE DEFICIENCY, *PDHB*-RELATED (*PDHB*) **negative**

R

REFSUM DISEASE, *PHYH*-RELATED (*PHYH*) **negative**
 RENAL TUBULAR ACIDOSIS AND DEAFNESS, *ATP6V1B1*-RELATED (*ATP6V1B1*) **negative**
 RENAL TUBULAR ACIDOSIS, PROXIMAL, WITH OCULAR ABNORMALITIES AND MENTAL RETARDATION (*SLC4A4*) **negative**
 RETINITIS PIGMENTOSA 25 (*EYS*) **negative**
 RETINITIS PIGMENTOSA 26 (*CERKL*) **negative**
 RETINITIS PIGMENTOSA 28 (*FAM161A*) **negative**
 RETINITIS PIGMENTOSA 36 (*PRCD*) **negative**
 RETINITIS PIGMENTOSA 59 (*DHDDS*) **negative**
 RETINITIS PIGMENTOSA 62 (*MAK*) **negative**
 RHIZOMELIC CHONDRODYSPLASIA PUNCTATA, TYPE 1 (*PEX7*) **negative**
 RHIZOMELIC CHONDRODYSPLASIA PUNCTATA, TYPE 2 (*GPNAT*) **negative**
 RHIZOMELIC CHONDRODYSPLASIA PUNCTATA, TYPE 3 (*AGPS*) **negative**
RLBP1-RELATED RETINOPATHY (*RLBP1*) **negative**
 ROBERTS SYNDROME (*ESCO2*) **negative**
RYR1-RELATED CONDITIONS (*RYR1*) **negative**

S

SALLA DISEASE (*SLC17A5*) **negative**
 SANDHOFF DISEASE (*HEXB*) **negative**
 SCHIMKE IMMUNODYSPLASIA (*SMARCA1*) **negative**
 SCHINDLER DISEASE (*NAGA*) **negative**
 SEGAWA SYNDROME, *TH*-RELATED (*TH*) **negative**
 SENIOR-LOKEN SYNDROME 4/NEPHRONOPHTHISIS 4 (*NPHP4*) **negative**
 SEPIAPTERIN REDUCTASE DEFICIENCY (*SPR*) **negative**
 SEVERE COMBINED IMMUNODEFICIENCY (*SCID*), *CD3D*-RELATED (*CD3D*) **negative**
 SEVERE COMBINED IMMUNODEFICIENCY (*SCID*), *CD3E*-RELATED (*CD3E*) **negative**
 SEVERE COMBINED IMMUNODEFICIENCY (*SCID*), *FOXN1*-RELATED (*FOXN1*) **negative**
 SEVERE COMBINED IMMUNODEFICIENCY (*SCID*), *IKBKB*-RELATED (*IKBKB*) **negative**
 SEVERE COMBINED IMMUNODEFICIENCY (*SCID*), *IL7R*-RELATED (*IL7R*) **negative**
 SEVERE COMBINED IMMUNODEFICIENCY (*SCID*), *JAK3*-RELATED (*JAK3*) **negative**
 SEVERE COMBINED IMMUNODEFICIENCY (*SCID*), *PTPRC*-RELATED (*PTPRC*) **negative**
 SEVERE COMBINED IMMUNODEFICIENCY (*SCID*), *RAG1*-RELATED (*RAG1*) **negative**
 SEVERE COMBINED IMMUNODEFICIENCY, *ADA*-Related (*ADA*) **negative**
 SEVERE COMBINED IMMUNODEFICIENCY, TYPE ATHABASKAN (*DCLRE1C*) **negative**
 SHORT-RIB THORACIC DYSPLASIA 3 WITH OR WITHOUT POLYDACTYLY (*DYNC2H1*) **negative**
 SHWACHMAN-DIAMOND SYNDROME, *SBDS*-RELATED (*SBDS*) **negative**
 SIALIDOSIS (*NEU1*) **negative**
 SJÖGREN-LARSSON SYNDROME (*ALDH3A2*) **negative**
 SMITH-LEMLI-OPITZ SYNDROME (*DHCR7*) **negative**
 SPASTIC PARAPLEGIA, TYPE 15 (*ZFYVE26*) **negative**

SPASTIC TETRAPLEGIA, THIN CORPUS CALLOSUM, AND PROGRESSIVE MICROCEPHALY (*SPATCCM*) (*SLC1A4*) **negative**
SPG11-RELATED CONDITIONS (*SPG11*) **negative**
 SPINAL MUSCULAR ATROPHY (*SMN1*) **negative** *SMN1*: Two copies; *g.27134T>G*: absent; the absence of the *g.27134T>G* variant decreases the chance to be a silent (2+0) carrier.
 SPINAL MUSCULAR ATROPHY WITH RESPIRATORY DISTRESS TYPE 1 (*IGHMBP2*) **negative**
 SPINOCEREBELLAR ATAXIA, AUTOSOMAL RECESSIVE 10 (*ANO10*) **negative**
 SPINOCEREBELLAR ATAXIA, AUTOSOMAL RECESSIVE 12 (*WWOX*) **negative**
 SPONDYLOCOSTAL DYSOSTOSIS 1 (*DLL3*) **negative**
 SPONDYLOTHORACIC DYSOSTOSIS, *MESP2*-Related (*MESP2*) **negative**
 STEEL SYNDROME (*COL27A1*) **negative**
 STEROID-RESISTANT NEPHROTIC SYNDROME (*NPHS2*) **see first page**
 STUVE-WIEDEMANN SYNDROME (*LIFR*) **negative**
SURF1-RELATED CONDITIONS (*SURF1*) **negative**
 SURFACTANT DYSFUNCTION, *ABCA3*-RELATED (*ABCA3*) **negative**

T

TAY-SACHS DISEASE (*HEXA*) **negative**
TBCE-RELATED CONDITIONS (*TBCE*) **negative**
 THIAMINE-RESPONSIVE MEGALOBlastic ANEMIA SYNDROME (*SLC19A2*) **negative**
 THYROID DYSHORMONOGENESIS 1 (*SLC5A5*) **negative**
 THYROID DYSHORMONOGENESIS 2A (*TPO*) **negative**
 THYROID DYSHORMONOGENESIS 3 (*TG*) **negative**
 THYROID DYSHORMONOGENESIS 6 (*DUOX2*) **negative**
 TRANSCOBALAMIN II DEFICIENCY (*TCN2*) **negative**
 TRICHOHEPATOENTERIC SYNDROME, *SKIC2*-RELATED (*SKIC2*) **negative**
 TRICHOHEPATOENTERIC SYNDROME, *TTC37*-RELATED (*TTC37*) **negative**
 TRICHOHYDROSTROPHY 1/XERODERMA PIGMENTOSUM, GROUP D (*ERCC2*) **negative**
 TRIMETHYLAMINURIA (*FMO3*) **negative**
 TRIPLE A SYNDROME (*AAAS*) **negative**
TSHR-RELATED CONDITIONS (*TSHR*) **negative**
 TYROSINEMIA TYPE III (*HPD*) **negative**
 TYROSINEMIA, TYPE 1 (*FAH*) **negative**
 TYROSINEMIA, TYPE 2 (*TAT*) **negative**

U

USHER SYNDROME, TYPE 1B (*MYO7A*) **negative**
 USHER SYNDROME, TYPE 1C (*USH1C*) **negative**
 USHER SYNDROME, TYPE 1D (*CDH23*) **negative**
 USHER SYNDROME, TYPE 1F (*PCDH15*) **negative**
 USHER SYNDROME, TYPE 1J/DEAFNESS, AUTOSOMAL RECESSIVE, 48 (*CIB2*) **negative**
 USHER SYNDROME, TYPE 2A (*USH2A*) **negative**
 USHER SYNDROME, TYPE 2C (*ADGRV1*) **negative**
 USHER SYNDROME, TYPE 3 (*CLRN1*) **negative**

V

VERY LONG-CHAIN ACYL-CoA DEHYDROGENASE DEFICIENCY (*ACADVL*) **negative**
 VICI SYNDROME (*EPG5*) **negative**
 VITAMIN D-DEPENDENT RICKETS, TYPE 1A (*CYP27B1*) **negative**
 VITAMIN D-RESISTANT RICKETS TYPE 2A (*VDR*) **negative**
VLDLR-ASSOCIATED CEREBELLAR HYPOPLASIA (*VLDLR*) **negative**

W

WALKER-WARBURG SYNDROME, *CRPPA*-RELATED (*CRPPA*) **negative**
 WALKER-WARBURG SYNDROME, *FKTN*-RELATED (*FKTN*) **negative**
 WALKER-WARBURG SYNDROME, *LARGE1*-RELATED (*LARGE1*) **negative**
 WALKER-WARBURG SYNDROME, *POMT1*-RELATED (*POMT1*) **negative**
 WALKER-WARBURG SYNDROME, *POMT2*-RELATED (*POMT2*) **negative**
 WARSAW BREAKAGE SYNDROME (*DDX11*) **negative**
 WERNER SYNDROME (*WRN*) **negative**
 WILSON DISEASE (*ATP7B*) **negative**
 WOLCOTT-RALLISON SYNDROME (*EIF2AK3*) **negative**
 WOLMAN DISEASE (*LIPA*) **negative**
 WOODHOUSE-SAKATI SYNDROME (*DCAF17*) **negative**

X

XERODERMA PIGMENTOSUM VARIANT TYPE (*POLH*) **negative**
 XERODERMA PIGMENTOSUM, GROUP A (*XPA*) **negative**
 XERODERMA PIGMENTOSUM, GROUP C (*XPC*) **negative**

Z

ZELLWEGER SPECTRUM DISORDER, *PEX13*-RELATED (*PEX13*) **negative**
 ZELLWEGER SPECTRUM DISORDER, *PEX16*-RELATED (*PEX16*) **negative**
 ZELLWEGER SPECTRUM DISORDER, *PEX5*-RELATED (*PEX5*) **negative**
 ZELLWEGER SPECTRUM DISORDERS, *PEX10*-RELATED (*PEX10*) **negative**
 ZELLWEGER SPECTRUM DISORDERS, *PEX12*-RELATED (*PEX12*) **negative**
 ZELLWEGER SPECTRUM DISORDERS, *PEX16*-RELATED (*PEX16*) **negative**
 ZELLWEGER SPECTRUM DISORDERS, *PEX26*-RELATED (*PEX26*) **negative**
 ZELLWEGER SPECTRUM DISORDERS, *PEX2*-RELATED (*PEX2*) **negative**

Patient Information

Patient Name:

Test Information

Ordering Physician:



Date Of Birth:



Clinic Information:

Case File ID:



Report Date:

Z

ZELLWEGER SPECTRUM DISORDERS, PEX6-RELATED (PEX6) **negative**

Patient Information

Patient Name: [REDACTED]

Test Information

Ordering Physician: [REDACTED]



Clinic Information:

Date Of Birth: [REDACTED]

Case File ID: [REDACTED]

Report Date:

Testing Methodology, Limitations, and Comments:**Next-generation sequencing (NGS)**

Sequencing library prepared from genomic DNA isolated from a patient sample is enriched for targets of interest using standard hybridization capture protocols and PCR amplification (for targets specified below). NGS is then performed to achieve the standards of quality control metrics, including a minimum coverage of 99% of targeted regions at 20X sequencing depth. Sequencing data is aligned to human reference sequence, followed by deduplication, metric collection and variant calling (coding region +/- 20bp). Variants are then classified according to ACMGG/AMP standards of interpretation using publicly available databases including but not limited to ENSEMBL, HGMD Pro, ClinGen, ClinVar, 1000G, ESP and gnomAD. Variants predicted to be pathogenic or likely pathogenic for the specified diseases are reported. It should be noted that the data interpretation is based on our current understanding of the genes and variants at the time of reporting. Putative positive sequencing variants that do not meet internal quality standards or are within highly homologous regions are confirmed by Sanger sequencing or gene-specific long-range PCR as needed prior to reporting.

Copy Number Variant (CNV) analysis is limited to deletions involving two or more exons for all genes on the panel, in addition to specific known recurrent single-exon deletions. CNVs of small size may have reduced detection rate. This method does not detect gene inversions, single-exonic and sub-exonic deletions (unless otherwise specified), and duplications of all sizes (unless otherwise specified). Additionally, this method does not define the exact breakpoints of detected CNV events. Confirmation testing for copy number variation is performed by specific PCR, Multiplex Ligation-dependent Probe Amplification (MLPA), next generation sequencing, or other methodology.

This test may not detect certain variants due to local sequence characteristics, high/low genomic complexity, homologous sequence, or allele dropout (PCR-based assays). Variants within noncoding regions (promoter, 5'UTR, 3'UTR, deep intronic regions, unless otherwise specified), small deletions or insertions larger than 25bp, low-level mosaic variants, structural variants such as inversions, and/or balanced translocations may not be detected with this technology.

SPECIAL NOTES

For ABCC6, variants in exons 1-9 are not detected due to the presence of regions of high homology.

For CFTR, when the CFTR R117H variant is detected, reflex analysis of the polythymidine variations (5T, 7T and 9T) at the intron 9 branch/acceptor site of the CFTR gene will be performed.

For CYP21A2, targets were enriched using long-range PCR amplification, followed by next generation sequencing. Duplication analysis will only be performed and reported when c.955C>T (p.Q319*) is detected. Sequencing and CNV analysis may have reduced sensitivity, if variants result from complex rearrangements, in trans with a gene deletion, or CYP21A2 gene duplication on one chromosome and deletion on the other chromosome. This analysis cannot detect sequencing variants located on the CYP21A2 duplicated copy.

For DDX11, only NM_030653.3:c.1763 - 1G > C variant will be analyzed and reported.

For HBA1/HBA2, CNV analysis is offered to detect common deletions of -alpha3.7, -alpha4.2, --MED, --SEA, --FIL, --THAI, --alpha20.5, and/or HS-40.

For OTOA, variants in exons 20 - 28 are not analyzed due to high sequence homology.

For RPGRIP1L, variants in exon 23 are not detected due to assay limitation.

For SAMD9, only p.K1495E variant will be analyzed and reported.

Friedreich Ataxia (FXN)

The GAA repeat region of the FXN gene is assessed by trinucleotide PCR assay and capillary electrophoresis. Variances of +/-1 repeat for normal alleles and up to +/-3 repeats for premutation alleles may occur. For fully penetrant expanded alleles, the precise repeat size cannot be determined, therefore the approximate allele size is reported. This analysis does not detect deletions or point mutations, which comprise about four percent of the FXN pathogenic variants.

Friedreich Ataxia Repeat Categories

Categories	GAA Repeat Sizes
Normal	<34
Premutation	34 - 65
Full	>65

Patient Information

Patient Name: [REDACTED]

Test Information

Ordering Physician: [REDACTED]



Date Of Birth: [REDACTED]

Clinic Information:

Case File ID: [REDACTED]

Report Date:

Spinal Muscular Atrophy (SMN1)

The total combined copy number of SMN1 and SMN2 exon 7 is quantified based on NGS read depth. The ratio of SMN1 to SMN2 is calculated based on the read depth of a single nucleotide that distinguishes these two genes in exon 7. In addition to copy number analysis, testing for the presence or absence of a single nucleotide polymorphism (g.27134T>G in intron 7 of SMN1) associated with the presence of a SMN1 duplication allele is performed using NGS.

Ethnicity	Two SMN1 copies carrier risk before g.27134T>G testing	Carrier risk after g.27134T>G testing	
		g.27134T>G ABSENT	g.27134T>G PRESENT
Caucasian	1 in 632	1 in 769	1 in 29
Ashkenazi Jewish	1 in 350	1 in 580	LIKELY CARRIER
Asian	1 in 628	1 in 702	LIKELY CARRIER
African-American	1 in 121	1 in 396	1 in 34
Hispanic	1 in 1061	1 in 1762	1 in 140

Variant Classification

Only pathogenic or likely pathogenic variants are reported. Other variants including benign variants, likely benign variants, variants of uncertain significance, or inconclusive variants identified during this analysis may be reported in certain circumstances. Our laboratory's variant classification criteria are based on the ACMG and internal guidelines and our current understanding of the specific genes. This interpretation may change over time as more information about a gene and/or variant becomes available. Natera and its lab partner(s) may reclassify variants at certain intervals but may not release updated reports without a specific request made to Natera by the ordering provider. Natera may disclose incidental findings if deemed clinically pertinent to the test performed.

Negative Results

A negative carrier screening result reduces the risk for a patient to be a carrier of a specific disease but does not completely rule out carrier status. Please visit <https://www.natera.com/panel-option/h-all/> for a table of carrier rates, detection rates, residual risks and promised variants/exons per gene. Carrier rates before and after testing vary by ethnicity and assume a negative family history for each disease screened and the absence of clinical symptoms in the patient. Any patient with a family history for a specific genetic disease will have a higher carrier risk prior to testing and, if the disease-causing mutation in their family is not included on the test, their carrier risk would remain unchanged. Genetic counseling is recommended for patients with a family history of genetic disease so that risk figures based on actual family history can be determined and discussed along with potential implications for reproduction. Horizon carrier screening has been developed to identify the reproductive risks for monogenic inherited conditions. Even when one or both members of a couple screen negative for pathogenic variants in a specific gene, the disease risk for their offspring is not zero. There is still a low risk for the condition in their offspring due to a number of different mechanisms that are not detected by Horizon including, but not limited to, pathogenic variant(s) in the tested gene or in a different gene not included on Horizon, pathogenic variant(s) in an upstream regulator, uniparental disomy, de novo mutation(s), or digenic or polygenic inheritance.

Additional Comments

These analyses generally provide highly accurate information regarding the patient's carrier status. Despite this high level of accuracy, it should be kept in mind that there are many potential sources of diagnostic error, including misidentification of samples, polymorphisms, or other rare genetic variants that interfere with analysis. Families should understand that rare diagnostic errors may occur for these reasons.

Patient Information

Patient Name: Donor 7414
Date Of Birth: [REDACTED]
Gender: Male
Ethnicity: Other
Patient ID: [REDACTED]
Medical Record #: N/A
Collection Kit: [REDACTED]
Accession ID: N/A
Case File ID: [REDACTED]

Test Information

Ordering Physician: [REDACTED]
Clinic Information: Fairfax Cryobank
Phone: N/A
Report Date: 04/19/2024
Sample Collected: 03/18/2024
Sample Received: 03/19/2024
Sample Type: Blood

CARRIER SCREENING REPORT

ABOUT THIS SCREEN: Horizon™ is a carrier screen for specific autosomal recessive and X-linked diseases. This information can help patients learn their risk of having a child with specific genetic conditions.

ORDER SELECTED: The Horizon Custom panel was ordered for this patient. Males are not screened for X-linked diseases

AMENDED FINAL RESULTS SUMMARY:**CARRIER for Non-Syndromic Hearing Loss, GJB2-Related**

Positive for the pathogenic variant c.269T>C (p.L90P) in the GJB2 gene. Although most variants in this gene are associated with an autosomal recessive form of NON-SYNDROMIC HEARING LOSS, GJB2-RELATED, some rare GJB2 variants may cause an autosomal dominant form of the condition. To our knowledge, there is insufficient evidence that this variant causes an autosomal dominant form of this condition. If this individual's partner is a carrier for NON-SYNDROMIC HEARING LOSS, GJB2-RELATED, their chance to have a child with this condition is likely 1 in 4 (25%). Carrier screening for this individual's partner is suggested.

CARRIER for Steroid-Resistant Nephrotic Syndrome

Positive for the likely pathogenic variant c.686G>A (p.R229Q) in the NPHS2 gene. Please note the likely pathogenic variant c.686G>A, p.R229Q is a mild variant that is only expected to cause disease when found in trans with one of a specific set of variants that occurs in exons 7 or 8 (PMID: 30260545, 30241959). Homozygotes are not expected to be affected, unless this variant is part of a more complex allele (PMID: 24509478, 29660491). If this individual's partner is a carrier for Steroid-Resistant Nephrotic Syndrome, their chance to have a child with this condition may be as high as 1 in 4 (25%). Carrier screening for this individual's partner is suggested.

Negative for 547 out of 549 diseases


No other pathogenic variants were detected in the genes that were screened. The patient's remaining carrier risk after the negative screening results is listed for each disease/gene on the Horizon website at <https://www.natera.com/panel-option/h-all/>. Please see the following pages of this report for a comprehensive list of all conditions included on this individual's screen.

Carrier screening is not diagnostic and may not detect all possible pathogenic variants in a given gene.


AMENDED REPORT 04/19/2024: This report is an update to the previous report dated 04/01/2024 to include results for the updated panel.

RECOMMENDATIONS


Individuals who would like to review their Horizon report with a Natera Laboratory Genetic Counselor may schedule a telephone genetic information session by calling 650-249-9090 or visiting naterasession.com. Clinicians with questions may contact Natera at 650-249-9090 or email support@natera.com. Individuals with positive results may wish to discuss these results with family members to allow them the option to be screened. Comprehensive genetic counseling to discuss the implications of these test results and possible associated reproductive risk is recommended.




Christine M. Eng, M.D.
Medical Director, Baylor Genetics



Jun Chen, Ph.D.
Assistant Laboratory Director, Baylor Genetics



J. Dianne Keen-Kim, Ph.D., FACMG
Senior Laboratory Director, Natera



Yang Wang, Ph.D., FACMG
Laboratory Director, Natera

Patient Information

Patient Name: Donor 7414

Test Information

Ordering Physician: [REDACTED]



Clinic Information: Fairfax Cryobank

Date Of Birth: [REDACTED]

Case File ID: [REDACTED]

Report Date: 04/19/2024

NON-SYNDROMIC HEARING LOSS, GJB2-RELATED**Understanding Your Horizon Carrier Screen Results****What is Non-Syndromic Hearing Loss, GJB2-Related?**

Non-Syndromic Hearing Loss, GJB2-Related (also called DFNB1) is an inherited disorder that causes early-onset hearing loss. "Non-syndromic" means that no other parts of the body are affected, making hearing loss the only symptom of this condition. In Non-Syndromic Hearing Loss, GJB2-Related, hearing loss is typically present at birth (congenital). However, some children have normal hearing at birth and develop hearing loss during childhood. The severity varies from mild to profound sensorineural hearing loss. The treatment for hearing loss includes hearing aids and, in some cases, cochlear implants. Clinical trials involving potential new treatments for this condition may be available (see www.clinicaltrials.gov). Non-Syndromic Hearing Loss, GJB2-Related does not cause other health problems.

What causes Non-Syndromic Hearing Loss, GJB2-Related?

Non-Syndromic Hearing Loss, GJB2-Related is caused by a gene change, or mutation, in both copies of the GJB2 gene pair (also known as DFNB1). These mutations cause the genes to not work properly or not work at all. The function of the GJB2 genes is to make a protein that is important for hearing. When both copies of the GJB2 gene do not work correctly, it leads to Non-Syndromic Hearing Loss, GJB2-Related. Non-Syndromic Hearing Loss, GJB2-Related is inherited in an autosomal recessive manner. This means that, in most cases, both parents must be carriers of a mutation in one copy of the GJB2 gene to have a child with Non-Syndromic Hearing Loss, GJB2-Related. People who are carriers for Non-Syndromic Hearing Loss, GJB2-Related are usually healthy and usually do not have Non-Syndromic Hearing Loss themselves. Usually a child inherits two copies of each gene, one copy from the mother and one copy from the father. If the mother and father are both carriers for Non-Syndromic Hearing Loss, GJB2-Related, there is a 1 in 4, or 25%, chance in each pregnancy for both partners to pass on their GJB2 gene mutations to the child, who will then have Non-Syndromic Hearing Loss, GJB2-Related. Very rarely, carriers of a single GJB2 mutation will have inherited hearing loss with or without other symptoms. These individuals usually have one parent who is also affected. This type of inheritance, where having only one mutation causes symptoms, is called autosomal dominant. When a person with autosomal dominant hearing loss has a child, there is a 50%, or 1 in 2, chance with each pregnancy of having a child who will also develop this type of hearing loss. It is sometimes, but not always, possible to determine whether a specific mutation in the GJB2 gene will cause autosomal recessive Non-Syndromic Hearing Loss or an autosomal dominant type of hearing loss. Individuals found to carry more than one mutation for Non-Syndromic Hearing Loss, GJB2-Related should discuss their risk for having an affected child and any potential effects to their own hearing with their health care provider.

What can I do next?

You may wish to speak with a local genetic counselor about your carrier test results. A genetic counselor in your area can be located on the National Society of Genetic Counselors website (www.nsgc.org). You may wish to share your carrier screening results with your health care providers, especially if you have a family history of hearing loss or have concerns about your own hearing. Your siblings and other relatives are at increased risk to also have this mutation. You are encouraged to inform your family members of your test results as they may wish to consider being tested themselves. If you are pregnant, your partner can have carrier screening for Non-Syndromic Hearing Loss, GJB2-Related ordered by a health care professional. If your partner is not found to be a carrier for Non-Syndromic Hearing Loss, GJB2-Related, your risk of having a child with Non-Syndromic Hearing Loss, GJB2-Related is greatly reduced. Couples at risk of having a baby with Non-Syndromic Hearing Loss, GJB2-Related can opt to have prenatal diagnosis done through chorionic villus sampling (CVS) or amniocentesis during pregnancy or can choose to test the baby after birth for this condition. If you are not yet pregnant, your partner can have carrier screening for Non-Syndromic Hearing Loss, GJB2-Related ordered by a health care professional. If your partner is found to be a carrier for Non-Syndromic Hearing Loss, GJB2-Related, you have several reproductive options to consider:

- Natural pregnancy with or without prenatal diagnosis of the fetus or testing the baby after birth for Non-Syndromic Hearing Loss, GJB2-Related
- Preimplantation genetic diagnosis (PGD) with in vitro fertilization (IVF) to test embryos for Non-Syndromic Hearing Loss, GJB2-Related
- Adoption or use of a sperm or egg donor who is not a carrier for Non-Syndromic Hearing Loss, GJB2-Related

What resources are available?

- Genetics Home Reference: <http://ghr.nlm.nih.gov/condition/nonsyndromic-hearing-loss>
- Prenatal diagnosis done through CVS: <http://www.marchofdimes.org/chorionic-villus-sampling.aspx>
- Prenatal diagnosis done through Amniocentesis: <http://www.marchofdimes.org/amniocentesis.aspx>
- Preimplantation genetic diagnosis (PGD) with IVF: <http://www.natera.com/spectrum>

Patient Information

Patient Name: [REDACTED]

Test Information

Ordering Physician: [REDACTED]



Clinic Information: [REDACTED]

Date Of Birth: [REDACTED]

Case File ID: [REDACTED]

Report Date: [REDACTED]

STEROID-RESISTANT NEPHROTIC SYNDROME**Understanding Your Horizon Carrier Screen Results****What is Steroid-Resistant Nephrotic Syndrome?**

Steroid-Resistant Nephrotic Syndrome (also known as Familial Nephrotic Syndrome or Nephrotic Syndrome, Type 2) is an inherited disorder that causes abnormal kidney function. People with this condition have large amounts of protein in their urine, low amounts of albumin (a protein in the plasma of the blood), high levels of fat in the blood, and excess fluid in body tissues (edema). Symptoms vary from person to person and usually start in infancy or childhood, although some people do not show symptoms until early adulthood. The kidney problems worsen over time, often leading to kidney failure in childhood, the teenage years, or early adulthood. Once kidney failure occurs, dialysis and then kidney transplantation are needed. Currently there is no cure for this condition and treatment is based on symptoms. Clinical trials involving potential new treatments for this condition may be available (see www.clinicaltrials.gov).

What causes Steroid-Resistant Nephrotic Syndrome?

Steroid-Resistant Nephrotic Syndrome is caused by a gene change, or mutation, in both copies of the NPHS2 gene pair. These mutations cause the genes to not work properly or not work at all. When both copies of this gene do not work correctly, it leads to the symptoms described above. Steroid-Resistant Nephrotic Syndrome is inherited in an autosomal recessive manner. This means that, in most cases, both parents must be carriers of a mutation in one copy of the NPHS2 gene to have a child with Steroid-Resistant Nephrotic Syndrome. People who are carriers for Steroid-Resistant Nephrotic Syndrome are usually healthy and do not have symptoms nor do they have Steroid-Resistant Nephrotic Syndrome themselves. Usually a child inherits two copies of each gene, one copy from the mother and one copy from the father. If the mother and father are both carriers for Steroid-Resistant Nephrotic Syndrome, there is a 1 in 4, or 25%, chance in each pregnancy for both partners to pass on their NPHS2 gene mutations to the child, who would then have this condition. Individuals found to carry more than one mutation for Steroid-Resistant Nephrotic Syndrome should discuss any potential effects to their own health and their risk for having an affected child with their health care provider.

What can I do next?

You may wish to speak with a local genetic counselor about your carrier test results. A genetic counselor in your area can be located on the National Society of Genetic Counselors website (www.nsgc.org). Your siblings and other relatives are at increased risk to also have this mutation. You are encouraged to inform your family members of your test results as they may wish to consider being tested themselves. If you are pregnant, your partner can have carrier screening for Steroid-Resistant Nephrotic Syndrome ordered by a health care professional. If your partner is not found to be a carrier for Steroid-Resistant Nephrotic Syndrome, your risk of having a child with this condition is greatly reduced. Couples at risk of having a baby with Steroid-Resistant Nephrotic Syndrome can opt to have prenatal diagnosis done through chorionic villus sampling (CVS) or amniocentesis during pregnancy or can choose to have the baby tested after birth for this condition. If you are not yet pregnant, your partner can have carrier screening for Steroid-Resistant Nephrotic Syndrome ordered by a health care professional. If your partner is found to be a carrier for Steroid-Resistant Nephrotic Syndrome, you have several reproductive options to consider:

- Natural pregnancy with or without prenatal diagnosis of the fetus or testing the baby after birth for Steroid-Resistant Nephrotic Syndrome
- Preimplantation genetic diagnosis (PGD) with in vitro fertilization (IVF) to test embryos for Steroid-Resistant Nephrotic Syndrome
- Adoption or use of a sperm or egg donor who is not a carrier for Steroid-Resistant Nephrotic Syndrome

What resources are available?

- The Renal Association: www.rarerrenal.org
- Genetics Home Reference: <http://ghr.nlm.nih.gov/gene/NPHS2>
- Prenatal diagnosis done through CVS: <http://www.marchofdimes.org/chorionic-villus-sampling.aspx>
- Prenatal diagnosis done through Amniocentesis: <http://www.marchofdimes.org/amniocentesis.aspx>
- PGD with IVF: <http://www.natera.com/spectrum>

Patient Information

Patient Name:

Test Information

Ordering Physician:



Clinic Information:

Date Of Birth:

Case File ID:

Report Date:

DISEASES SCREENED

Below is a list of all diseases screened and the result. Certain conditions have unique patient-specific numerical values, therefore, results for those conditions are formatted differently.

Autosomal Recessive

1
17-BETA HYDROXYSTEROID DEHYDROGENASE 3 DEFICIENCY (*HSD17B3*) **negative**

3
3-BETA-HYDROXYSTEROID DEHYDROGENASE TYPE II DEFICIENCY (*HSD3B2*) **negative**
3-HYDROXY-3-METHYLGLUTARYL-COENZYME A LYASE DEFICIENCY (*HMGCL*) **negative**
3-HYDROXYACYL-COA DEHYDROGENASE DEFICIENCY (*HADH*) **negative**
3-METHYLCROTONYL-CoA CARBOXYLASE 2 DEFICIENCY (*MCC22*) **negative**
3-PHOSPHOGLYCERATE DEHYDROGENASE DEFICIENCY (*PHGDH*) **negative**

5
5-ALPHA-REDUCTASE DEFICIENCY (*SRD5A2*) **negative**

6
6-PYRUVOYL-TETRAHYDROPTERIN SYNTHASE (*PTPS*) DEFICIENCY (*PTS*) **negative**

A
ABCA4-RELATED CONDITIONS (*ABCA4*) **negative**
ABETALIPOPROTEINEMIA (*MTTP*) **negative**
ACHONDROGENESIS, TYPE 1B (*SLC26A2*) **negative**
ACHROMATOPSIA, CNGB3-RELATED (*CNGB3*) **negative**
ACRODERMATITIS ENTEROPATHICA (*SLC39A4*) **negative**
ACTION MYOCLONUS-RENAL FAILURE (AMRF) SYNDROME (*SCARB2*) **negative**
ACUTE INFANTILE LIVER FAILURE, TRMU-RELATED (*TRMU*) **negative**
ACYL-COA OXIDASE I DEFICIENCY (*ACOX1*) **negative**
AICARDI-GOUTIERES SYNDROME (*SAMHD1*) **negative**
AICARDI-GOUTIERES SYNDROME, RNASEH2A-RELATED (*RNASEH2A*) **negative**
AICARDI-GOUTIERES SYNDROME, RNASEH2B-RELATED (*RNASEH2B*) **negative**
AICARDI-GOUTIERES SYNDROME, RNASEH2C-RELATED (*RNASEH2C*) **negative**
AICARDI-GOUTIERES SYNDROME, TREX1-RELATED (*TREX1*) **negative**
ALPHA-MANNOSIDOSIS (*MAN2B1*) **negative**
ALPHA-THALASSEMIA (*HBA1/HBA2*) **negative**
ALPORT SYNDROME, COL4A3-RELATED (*COL4A3*) **negative**
ALPORT SYNDROME, COL4A4-RELATED (*COL4A4*) **negative**
ALSTROM SYNDROME (*ALMS1*) **negative**
AMISH INFANTILE EPILEPSY SYNDROME (*ST3GAL5*) **negative**
ANDERMANN SYNDROME (*SLC12A6*) **negative**
ARGININE:GLYCINE AMIDINOTRANSFERASE DEFICIENCY (AGAT DEFICIENCY) (*GATM*) **negative**
ARGININEMIA (*ARG1*) **negative**
ARGININOSUCCINATE LYASE DEFICIENCY (*ASL*) **negative**
AROMATASE DEFICIENCY (*CYP19A1*) **negative**
ASPARAGINE SYNTHETASE DEFICIENCY (*ASNS*) **negative**
ASPARTYLGLYCOSAMINURIA (*AGA*) **negative**
ATAXIA WITH VITAMIN E DEFICIENCY (*TTPA*) **negative**
ATAXIA-TELANGIECTASIA (*ATM*) **negative**
ATAXIA-TELANGIECTASIA-LIKE DISORDER 1 (*MRE11*) **negative**
ATANSFERRINEMIA (*TF*) **negative**
AUTISM SPECTRUM, EPILEPSY AND ARTHROGRYPOSIS (*SLC35A3*) **negative**
AUTOIMMUNE POLYGLANDULAR SYNDROME, TYPE 1 (*AIRE*) **negative**
AUTOSOMAL RECESSIVE CONGENITAL ICHTHYOSIS (ARCI), SLC27A4-RELATED (*SLC27A4*) **negative**
AUTOSOMAL RECESSIVE SPASTIC ATAXIA OF CHARLEVOIX-SAGUENAY (*SACS*) **negative**

B
BARDET-BIEDL SYNDROME, ARL6-RELATED (*ARL6*) **negative**
BARDET-BIEDL SYNDROME, BBS10-RELATED (*BBS10*) **negative**
BARDET-BIEDL SYNDROME, BBS12-RELATED (*BBS12*) **negative**
BARDET-BIEDL SYNDROME, BBS1-RELATED (*BBS1*) **negative**
BARDET-BIEDL SYNDROME, BBS2-RELATED (*BBS2*) **negative**
BARDET-BIEDL SYNDROME, BBS4-RELATED (*BBS4*) **negative**
BARDET-BIEDL SYNDROME, BBS5-RELATED (*BBS5*) **negative**
BARDET-BIEDL SYNDROME, BBS7-RELATED (*BBS7*) **negative**
BARDET-BIEDL SYNDROME, BBS9-RELATED (*BBS9*) **negative**
BARDET-BIEDL SYNDROME, TTC8-RELATED (*TTC8*) **negative**
BARE LYMPHOCYTE SYNDROME, CIITA-RELATED (*CIITA*) **negative**
BARTTER SYNDROME, BSND-RELATED (*BSND*) **negative**
BARTTER SYNDROME, KCNJ1-RELATED (*KCNJ1*) **negative**
BARTTER SYNDROME, SLC12A1-RELATED (*SLC12A1*) **negative**
BATTEN DISEASE, CLN3-RELATED (*CLN3*) **negative**
BETA-HEMOGLOBINOPATHIES (*HBB*) **negative**
BETA-KETOTHIOLASE DEFICIENCY (*ACAT1*) **negative**
BETA-MANNOSIDOSIS (*MANBA*) **negative**
BETA-UREIDOPROPIONASE DEFICIENCY (*UPB1*) **negative**
BILATERAL FRONTOPIRIETAL POLYMICROGYRIA (*GPR56*) **negative**

BIOTINIDASE DEFICIENCY (*BTD*) **negative**
BIOTIN-THIAMINE-RESPONSIVE BASAL GANGLIA DISEASE (BTBGD) (*SLC19A3*) **negative**
BLOOM SYNDROME (*BLM*) **negative**
BRITTLE CORNEA SYNDROME 1 (*ZNF469*) **negative**
BRITTLE CORNEA SYNDROME 2 (*PRDM5*) **negative**

C
CANAVAN DISEASE (*ASPA*) **negative**
CARBAMOYL PHOSPHATE SYNTHETASE I DEFICIENCY (*CPS1*) **negative**
CARNITINE DEFICIENCY (*SLC22A5*) **negative**
CARNITINE PALMITOYLTRANSFERASE IA DEFICIENCY (*CPT1A*) **negative**
CARNITINE PALMITOYLTRANSFERASE II DEFICIENCY (*CPT2*) **negative**
CARNITINE-ACYLCARNITINE TRANSLOCASE DEFICIENCY (*SLC25A20*) **negative**
CARPENTER SYNDROME (*RAB23*) **negative**
CARTILAGE-HAIR HYPOPLASIA (*RMRP*) **negative**
CATECHOLAMINERGIC POLYMORPHIC VENTRICULAR TACHYCARDIA (*CASQ2*) **negative**
CD59-MEDIATED HEMOLYTIC ANEMIA (*CD59*) **negative**
CEP152-RELATED MICROCEPHALY (*CEP152*) **negative**
CEREBRAL DYSGENESIS, NEUROPATHY, ICHTHYOSIS, AND PALMOPANTAR KERATODERMA (CEDNIK) SYNDROME (*SNAP29*) **negative**
CEREBROTENDINOUS XANTHOMATOSIS (*CYP27A1*) **negative**
CHARCOT-MARIE-TOOTH DISEASE, RECESSIVE INTERMEDIATE C (*PLEKHG5*) **negative**
CHARCOT-MARIE-TOOTH-DISEASE, TYPE 4D (*NDRG1*) **negative**
CHEDIAK-HIGASHI SYNDROME (*LYST*) **negative**
CHOREOACANTHOCTOSIS (*VPS13A*) **negative**
CHRONIC GRANULOMATOUS DISEASE, CYBA-RELATED (*CYBA*) **negative**
CHRONIC GRANULOMATOUS DISEASE, NCF2-RELATED (*NCF2*) **negative**
CILIOPATHIES, RPGRIP1L-RELATED (*RPGRIP1L*) **negative**
CITRIN DEFICIENCY (*SLC25A13*) **negative**
CITRULLINEMIA, TYPE 1 (*ASS1*) **negative**
CLN10 DISEASE (*CTSD*) **negative**
COHEN SYNDROME (*VPS13B*) **negative**
COL11A2-RELATED CONDITIONS (*COL11A2*) **negative**
COMBINED MALONIC AND METHYLMALONIC ACIDURIA (*ACSF3*) **negative**
COMBINED OXIDATIVE PHOSPHORYLATION DEFICIENCY 1 (*GFM1*) **negative**
COMBINED OXIDATIVE PHOSPHORYLATION DEFICIENCY 3 (*TSMF*) **negative**
COMBINED PITUITARY HORMONE DEFICIENCY 1 (*POU1F1*) **negative**
COMBINED PITUITARY HORMONE DEFICIENCY-2 (*PROP1*) **negative**
CONGENITAL ADRENAL HYPERPLASIA, 11-BETA-HYDROXYLASE DEFICIENCY (*CYP11B1*) **negative**
CONGENITAL ADRENAL HYPERPLASIA, 17-ALPHA-HYDROXYLASE DEFICIENCY (*CYP17A1*) **negative**
CONGENITAL ADRENAL HYPERPLASIA, 21-HYDROXYLASE DEFICIENCY (*CYP21A2*) **negative**
CONGENITAL ADRENAL INSUFFICIENCY, CYP11A1-RELATED (*CYP11A1*) **negative**
CONGENITAL AMEGAKARYOCYTIC THROMBOCYTOPENIA (*MPL*) **negative**
CONGENITAL CHRONIC DIARRHEA (*DGAT1*) **negative**
CONGENITAL DISORDER OF GLYCOSYLATION TYPE 1, ALG1-RELATED (*ALG1*) **negative**
CONGENITAL DISORDER OF GLYCOSYLATION, TYPE 1A, PMM2-Related (*PMM2*) **negative**
CONGENITAL DISORDER OF GLYCOSYLATION, TYPE 1B (*MPI*) **negative**
CONGENITAL DISORDER OF GLYCOSYLATION, TYPE 1C (*ALG6*) **negative**
CONGENITAL DYSERYTHROPOIETIC ANEMIA TYPE 2 (*SEC23B*) **negative**
CONGENITAL FINNISH NEPHROSIS (*NPHS1*) **negative**
CONGENITAL HYDROCEPHALUS 1 (*CCDC88C*) **negative**
CONGENITAL HYPERINSULINISM, KCNJ11-Related (*KCNJ11*) **negative**
CONGENITAL INSENSITIVITY TO PAIN WITH ANHIDROSIS (CIPA) (*NTRK1*) **negative**
CONGENITAL MYASTHENIC SYNDROME, CHAT-RELATED (*CHAT*) **negative**
CONGENITAL MYASTHENIC SYNDROME, CHRNE-RELATED (*CHRNE*) **negative**
CONGENITAL MYASTHENIC SYNDROME, COLQ-RELATED (*COLQ*) **negative**
CONGENITAL MYASTHENIC SYNDROME, DOK7-RELATED (*DOK7*) **negative**
CONGENITAL MYASTHENIC SYNDROME, RAPSIN-RELATED (*RAPSIN*) **negative**
CONGENITAL NEPHROTIC SYNDROME, PLCE1-RELATED (*PLCE1*) **negative**
CONGENITAL NEUTROPENIA, G6PC3-RELATED (*G6PC3*) **negative**
CONGENITAL NEUTROPENIA, HAX1-RELATED (*HAX1*) **negative**
CONGENITAL NEUTROPENIA, VPS45-RELATED (*VPS45*) **negative**
CONGENITAL SECRETORY CHLORIDE DIARRHEA 1 (*SLC26A3*) **negative**
CORNEAL DYSTROPHY AND PERCEPTIVE DEAFNESS (*SLC4A11*) **negative**
CORTICOSTERONE METHYLOXIDASE DEFICIENCY (*CYP11B2*) **negative**
COSTEFF SYNDROME (3-METHYLGLUTACONIC ACIDURIA, TYPE 3) (*OPA3*) **negative**
CRB1-RELATED RETINAL DYSTROPHIES (*CRB1*) **negative**
CYSTIC FIBROSIS (*CFTR*) **negative**
CYSTINOSIS (*CTNS*) **negative**
CYTOCHROME C OXIDASE DEFICIENCY, PET100-RELATED (*PET100*) **negative**
CYTOCHROME P450 OXIOREDUCTASE DEFICIENCY (*POR*) **negative**

D
D-BIFUNCTIONAL PROTEIN DEFICIENCY (*HSD17B4*) **negative**

Patient Information

Patient Name:

Test Information

Ordering Physician:



Clinic Information:

Date Of Birth:

Case File ID:

Report Date:

D

DEAFNESS, AUTOSOMAL RECESSIVE 77 (*LOXHD1*) **negative**
 DIHYDROPTERIDINE REDUCTASE (DHPR) DEFICIENCY (*QDPR*) **negative**
 DONNAI-BARROW SYNDROME (*LRP2*) **negative**
 DUBIN-JOHNSON SYNDROME (*ABCC2*) **negative**
 DYSKERATOSIS CONGENITA SPECTRUM DISORDERS (*TERT*) **negative**
 DYSKERATOSIS CONGENITA, RTTEL1-RELATED (*RTTEL1*) **negative**
 DYSTROPHIC EPIDERMOLYSIS BULLOSA, COL7A1-Related (*COL7A1*) **negative**

E

EARLY INFANTILE EPILEPTIC ENCEPHALOPATHY, CAD-RELATED (*CAD*) **negative**
 EHLERS-DANLOS SYNDROME TYPE VI (*PLOD1*) **negative**
 EHLERS-DANLOS SYNDROME, CLASSIC-LIKE, TNXB-RELATED (*TNXB*) **negative**
 EHLERS-DANLOS SYNDROME, TYPE VII C (*ADAMTS2*) **negative**
 ELLIS-VAN CREVELD SYNDROME, EVC2-RELATED (*EVC2*) **negative**
 ELLIS-VAN CREVELD SYNDROME, EVC-RELATED (*EVC*) **negative**
 ENHANCED S-CONE SYNDROME (*NR2E3*) **negative**
 EPIMERASE DEFICIENCY (GALACTOSEMIA TYPE III) (*GALE*) **negative**
 EPIPHYSEAL DYSPLASIA, MULTIPLE, 7/DESBUQUOIS DYSPLASIA 1 (*CANT1*) **negative**
 ERCC6-RELATED DISORDERS (*ERCC6*) **negative**
 ERCC8-RELATED DISORDERS (*ERCC8*) **negative**
 ETHYLMALONIC ENCEPHALOPATHY (*ETHE1*) **negative**

F

FACTOR XI DEFICIENCY (*F11*) **negative**
 FAMILIAL DYSAUTONOMIA (*IKBKAP*) **negative**
 FAMILIAL HEMOPHAGOCYtic LYMPHOHISTIOCYTOSIS, PRF1-RELATED (*PRF1*) **negative**
 FAMILIAL HEMOPHAGOCYtic LYMPHOHISTIOCYTOSIS, STX11-RELATED (*STX11*) **negative**
 FAMILIAL HEMOPHAGOCYtic LYMPHOHISTIOCYTOSIS, STXBP2-RELATED (*STXBP2*) **negative**
 FAMILIAL HEMOPHAGOCYtic LYMPHOHISTIOCYTOSIS, UNC13D-RELATED (*UNC13D*) **negative**
 FAMILIAL HYPERCHOLESTEROLEMIA, LDLRAP1-RELATED (*LDLRAP1*) **negative**
 FAMILIAL HYPERCHOLESTEROLEMIA, LDLR-RELATED (*LDLR*) **negative**
 FAMILIAL HYPERINSULINISM, ABCC8-RELATED (*ABCC8*) **negative**
 FAMILIAL NEPHROGENIC DIABETES INSIPIDUS, AQP2-RELATED (*AQP2*) **negative**
 FANCONI ANEMIA, GROUP A (*FANCA*) **negative**
 FANCONI ANEMIA, GROUP C (*FANCC*) **negative**
 FANCONI ANEMIA, GROUP D2 (*FANCD2*) **negative**
 FANCONI ANEMIA, GROUP E (*FANCE*) **negative**
 FANCONI ANEMIA, GROUP F (*FANCF*) **negative**
 FANCONI ANEMIA, GROUP G (*FANCG*) **negative**
 FANCONI ANEMIA, GROUP I (*FANCI*) **negative**
 FANCONI ANEMIA, GROUP J (*BRIP1*) **negative**
 FANCONI ANEMIA, GROUP L (*FANCL*) **negative**
 FARBER LIPOGRANULOMATOSIS (*ASAH1*) **negative**
 FOVEAL HYPOPLASIA (*SLC38A8*) **negative**
 FRASER SYNDROME 3, GRIP1-RELATED (*GRIP1*) **negative**
 FRASER SYNDROME, FRAS1-RELATED (*FRAS1*) **negative**
 FRASER SYNDROME, FREM2-RELATED (*FREM2*) **negative**
 FRIEDREICH ATAXIA (*FXN*) **negative**
 FRUCTOSE-1,6-BISPHOSPHATASE DEFICIENCY (*FBP1*) **negative**
 FUCOSIDOSIS, FUCA1-RELATED (*FUCA1*) **negative**
 FUMARASE DEFICIENCY (*FH*) **negative**

G

GABA-TRANSAMINASE DEFICIENCY (*ABAT*) **negative**
 GALACTOKINASE DEFICIENCY (GALACTOSEMIA, TYPE II) (*GALK1*) **negative**
 GALACTOSEMIA (*GALT*) **negative**
 GALACTOSIALIDOSIS (*CTSA*) **negative**
 GAUCHER DISEASE (*GBA*) **negative**
 GCH1-RELATED CONDITIONS (*GCH1*) **negative**
 GDF5-RELATED CONDITIONS (*GDF5*) **negative**
 GERODERMA OSTEODYSPLASTICA (*GORAB*) **negative**
 GITELMAN SYNDROME (*SLC12A3*) **negative**
 GLANZMANN THROMBASTHENIA (*ITGB3*) **negative**
 GLUTARIC ACIDEMIA, TYPE 1 (*GCDH*) **negative**
 GLUTARIC ACIDEMIA, TYPE 2A (*ETFA*) **negative**
 GLUTARIC ACIDEMIA, TYPE 2B (*ETFB*) **negative**
 GLUTARIC ACIDEMIA, TYPE 2C (*ETFDH*) **negative**
 GLUTATHIONE SYNTHETASE DEFICIENCY (*GSS*) **negative**
 GLYCINE ENCEPHALOPATHY, AMT-RELATED (*AMT*) **negative**
 GLYCINE ENCEPHALOPATHY, GLDC-RELATED (*GLDC*) **negative**
 GLYCOGEN STORAGE DISEASE TYPE 5 (McArdle Disease) (*PYGM*) **negative**
 GLYCOGEN STORAGE DISEASE TYPE IXB (*PHKB*) **negative**
 GLYCOGEN STORAGE DISEASE TYPE IXC (*PHKG2*) **negative**
 GLYCOGEN STORAGE DISEASE, TYPE 1a (*G6PC*) **negative**
 GLYCOGEN STORAGE DISEASE, TYPE 1b (*SLC37A4*) **negative**
 GLYCOGEN STORAGE DISEASE, TYPE 2 (POMPE DISEASE) (*GAA*) **negative**
 GLYCOGEN STORAGE DISEASE, TYPE 3 (*AGL*) **negative**
 GLYCOGEN STORAGE DISEASE, TYPE 4 (*GBE1*) **negative**
 GLYCOGEN STORAGE DISEASE, TYPE 7 (*PFKM*) **negative**

GRACILE SYNDROME (*BCS1L*) **negative**
 GUANIDINOACETATE METHYLTRANSFERASE DEFICIENCY (*GAMT*) **negative**

H

HARLEQUIN ICHTHYOSIS (*ABCA12*) **negative**
 HEME OXYGENASE 1 DEFICIENCY (*HMOX1*) **negative**
 HEMOCHROMATOSIS TYPE 2A (*HFE2*) **negative**
 HEMOCHROMATOSIS, TYPE 3, TFR2-Related (*TFR2*) **negative**
 HEPATOCEREBRAL MITOCHONDRIAL DNA DEPLETION SYNDROME, MPV17-RELATED (*MPV17*) **negative**
 HEREDITARY FRUCTOSE INTOLERANCE (*ALDOB*) **negative**
 HEREDITARY HEMOCHROMATOSIS TYPE 2B (*HAMP*) **negative**
 HEREDITARY SPASTIC PARAPARESIS, TYPE 49 (*TECPR2*) **negative**
 HEREDITARY SPASTIC PARAPLEGIA, CYP7B1-RELATED (*CYP7B1*) **negative**
 HERMANSKY-PUDLAK SYNDROME, AP3B1-RELATED (*AP3B1*) **negative**
 HERMANSKY-PUDLAK SYNDROME, BLOC1S3-RELATED (*BLOC1S3*) **negative**
 HERMANSKY-PUDLAK SYNDROME, BLOC1S6-RELATED (*BLOC1S6*) **negative**
 HERMANSKY-PUDLAK SYNDROME, HPS1-RELATED (*HPS1*) **negative**
 HERMANSKY-PUDLAK SYNDROME, HPS3-RELATED (*HPS3*) **negative**
 HERMANSKY-PUDLAK SYNDROME, HPS4-RELATED (*HPS4*) **negative**
 HERMANSKY-PUDLAK SYNDROME, HPS5-RELATED (*HPS5*) **negative**
 HERMANSKY-PUDLAK SYNDROME, HPS6-RELATED (*HPS6*) **negative**
 HOLOCARBOXYLASE SYNTHETASE DEFICIENCY (*HLC5*) **negative**
 HOMOCYSTINURIA AND MEGALOBlastic ANEMIA TYPE CBLG (*MTR*) **negative**
 HOMOCYSTINURIA DUE TO DEFICIENCY OF MTHFR (*MTHFR*) **negative**
 HOMOCYSTINURIA, CBS-RELATED (*CBS*) **negative**
 HOMOCYSTINURIA, Type cblE (*MTRR*) **negative**
 HYDROLETHALUS SYNDROME (*HYLS1*) **negative**
 HYPER-IGM IMMUNODEFICIENCY (*CD40*) **negative**
 HYPERORNITHINEMIA-HYPERAMMONEMIA-HOMOCITRULLINURIA (HHH SYNDROME) (*SLC25A15*) **negative**
 HYPERPHOSPHATEMIC FAMILIAL TUMORAL CALCINOSIS, GALNT3-RELATED (*GALNT3*) **negative**
 HYPOMYELINATING LEUKODYSTROPHY 12 (*VPS11*) **negative**
 HYPOPHOSPHATASIA, ALPL-RELATED (*ALPL*) **negative**

I

IMERSLUND-GRÄSBECK SYNDROME 2 (*AMN*) **negative**
 IMMUNODEFICIENCY-CENTROMERIC INSTABILITY-FACIAL ANOMALIES (ICF) SYNDROME, DNMT3B-RELATED (*DNMT3B*) **negative**
 IMMUNODEFICIENCY-CENTROMERIC INSTABILITY-FACIAL ANOMALIES (ICF) SYNDROME, ZBTB24-RELATED (*ZBTB24*) **negative**
 INCLUSION BODY MYOPATHY 2 (*GNE*) **negative**
 INFANTILE CEREBRAL AND CEREBELLAR ATROPHY (*MED17*) **negative**
 INFANTILE NEPHRONOPHTHISIS (*INVS*) **negative**
 INFANTILE NEUROAXONAL DYSTROPHY (*PLA2G6*) **negative**
 ISOLATED ECTOPIA LENTIS (*ADAMTSL4*) **negative**
 ISOLATED SULFITE OXIDASE DEFICIENCY (*SUOX*) **negative**
 ISOLATED THYROID-STIMULATING HORMONE DEFICIENCY (*TSHB*) **negative**
 ISOVALERIC ACIDEMIA (*IVD*) **negative**

J

JOHANSON-BLIZZARD SYNDROME (*UBR1*) **negative**
 JOUBERT SYNDROME 2 / MECKEL SYNDROME 2 (*TMEM216*) **negative**
 JOUBERT SYNDROME AND RELATED DISORDERS (JSRD), TMEM67-RELATED (*TMEM67*) **negative**
 JOUBERT SYNDROME, AH11-RELATED (*AH11*) **negative**
 JOUBERT SYNDROME, ARL13B-RELATED (*ARL13B*) **negative**
 JOUBERT SYNDROME, B9D1-RELATED (*B9D1*) **negative**
 JOUBERT SYNDROME, B9D2-RELATED (*B9D2*) **negative**
 JOUBERT SYNDROME, C2CD3-RELATED/OROFACIODIGITAL SYNDROME 14 (*C2CD3*) **negative**
 JOUBERT SYNDROME, CC2D2A-RELATED/COACH SYNDROME (*CC2D2A*) **negative**
 JOUBERT SYNDROME, CEP104-RELATED (*CEP104*) **negative**
 JOUBERT SYNDROME, CEP120-RELATED/SHORT-RIB THORACIC DYSPLASIA 13 WITH OR WITHOUT POLYDACTYLY (*CEP120*) **negative**
 JOUBERT SYNDROME, CEP41-RELATED (*CEP41*) **negative**
 JOUBERT SYNDROME, CPLANE1-RELATED / OROFACIODIGITAL SYNDROME 6 (*CPLANE1*) **negative**
 JOUBERT SYNDROME, CSPP1-RELATED (*CSPP1*) **negative**
 JOUBERT SYNDROME, INPP5E-RELATED (*INPP5E*) **negative**
 JUNCTIONAL EPIDERMOLYSIS BULLOSA, COL17A1-RELATED (*COL17A1*) **negative**
 JUNCTIONAL EPIDERMOLYSIS BULLOSA, ITGA6-RELATED (*ITGA6*) **negative**
 JUNCTIONAL EPIDERMOLYSIS BULLOSA, ITGB4-RELATED (*ITGB4*) **negative**
 JUNCTIONAL EPIDERMOLYSIS BULLOSA, LAMB3-RELATED (*LAMB3*) **negative**
 JUNCTIONAL EPIDERMOLYSIS BULLOSA, LAMC2-RELATED (*LAMC2*) **negative**
 JUNCTIONAL EPIDERMOLYSIS BULLOSA/LARYNGOOXYCHOCUTANEOUS SYNDROME, LAMA3-RELATED (*LAMA3*) **negative**

K

KRABBE DISEASE (*GALC*) **negative**

L

LAMELLAR ICHTHYOSIS, TYPE 1 (*TGM1*) **negative**

Patient Information

Patient Name:

Test Information

Ordering Physician:



Clinic Information:

Date Of Birth:



Case File ID:



Report Date:

L
 LARON SYNDROME (*GHR*) **negative**
 LEBER CONGENITAL AMAUROSIS 2 (*RPE65*) **negative**
 LEBER CONGENITAL AMAUROSIS TYPE AIPL1 (*AIPL1*) **negative**
 LEBER CONGENITAL AMAUROSIS TYPE GUCY2D (*GUCY2D*) **negative**
 LEBER CONGENITAL AMAUROSIS TYPE TULP1 (*TULP1*) **negative**
 LEBER CONGENITAL AMAUROSIS, IQCB1-RELATED/SENIOR-LOKEN SYNDROME 5 (*IQCB1*) **negative**
 LEBER CONGENITAL AMAUROSIS, TYPE CEP290 (*CEP290*) **negative**
 LEBER CONGENITAL AMAUROSIS, TYPE LCA5 (*LCA5*) **negative**
 LEBER CONGENITAL AMAUROSIS, TYPE RDH12 (*RDH12*) **negative**
 LEIGH SYNDROME, FRENCH-CANADIAN TYPE (*LRPPRC*) **negative**
 LETHAL CONGENITAL CONTRACTURE SYNDROME 1 (*GLE1*) **negative**
 LEUKOENCEPHALOPATHY WITH VANISHING WHITE MATTER (*EIF2B5*) **negative**
 LEUKOENCEPHALOPATHY WITH VANISHING WHITE MATTER, EIF2B1-RELATED (*EIF2B1*) **negative**
 LEUKOENCEPHALOPATHY WITH VANISHING WHITE MATTER, EIF2B2-RELATED (*EIF2B2*) **negative**
 LEUKOENCEPHALOPATHY WITH VANISHING WHITE MATTER, EIF2B3-RELATED (*EIF2B3*) **negative**
 LEUKOENCEPHALOPATHY WITH VANISHING WHITE MATTER, EIF2B4-RELATED (*EIF2B4*) **negative**
 LIG4 SYNDROME (*LIG4*) **negative**
 LIMB-GIRDLE MUSCULAR DYSTROPHY TYPE 8 (*TRIM32*) **negative**
 LIMB-GIRDLE MUSCULAR DYSTROPHY, TYPE 2A (*CAPN3*) **negative**
 LIMB-GIRDLE MUSCULAR DYSTROPHY, TYPE 2B (*DYSF*) **negative**
 LIMB-GIRDLE MUSCULAR DYSTROPHY, TYPE 2C (*SGCG*) **negative**
 LIMB-GIRDLE MUSCULAR DYSTROPHY, TYPE 2D (*SGCA*) **negative**
 LIMB-GIRDLE MUSCULAR DYSTROPHY, TYPE 2E (*SGCB*) **negative**
 LIMB-GIRDLE MUSCULAR DYSTROPHY, TYPE 2F (*SGCD*) **negative**
 LIMB-GIRDLE MUSCULAR DYSTROPHY, TYPE 2I (*FKRP*) **negative**
 LIPOAMIDE DEHYDROGENASE DEFICIENCY (DIHYDROLIPOAMIDE DEHYDROGENASE DEFICIENCY) (*DLI*) **negative**
 LIPOID ADRENAL HYPERPLASIA (*STAR*) **negative**
 LIPOPROTEIN LIPASE DEFICIENCY (*LPL*) **negative**
 LONG CHAIN 3-HYDROXYACYL-COA DEHYDROGENASE DEFICIENCY (*HADHA*) **negative**
 LRAT-RELATED CONDITIONS (*LRAT*) **negative**
 LUNG DISEASE, IMMUNODEFICIENCY, AND CHROMOSOME BREAKAGE SYNDROME (LICS) (*NSMCE3*) **negative**
 LYSINURIC PROTEIN INTOLERANCE (*SLC7A7*) **negative**

M
 MALONYL-COA DECARBOXYLASE DEFICIENCY (*MLYCD*) **negative**
 MAPLE SYRUP URINE DISEASE, TYPE 1A (*BCKDHA*) **negative**
 MAPLE SYRUP URINE DISEASE, TYPE 1B (*BCKDHB*) **negative**
 MAPLE SYRUP URINE DISEASE, TYPE 2 (*DBT*) **negative**
 MCKUSICK-KAUFMAN SYNDROME (*MKKS*) **negative**
 MECKEL SYNDROME 7/NEPHRONOPHTHISIS 3 (*NPHP3*) **negative**
 MECKEL-GRUBER SYNDROME, TYPE 1 (*MKS1*) **negative**
 MECR-RELATED NEUROLOGIC DISORDER (*MECR*) **negative**
 MEDIUM CHAIN ACYL-CoA DEHYDROGENASE DEFICIENCY (*ACADM*) **negative**
 MEDNIK SYNDROME (*AP1S1*) **negative**
 MEGALENCEPHALIC LEUKOENCEPHALOPATHY WITH SUBCORTICAL CYSTS (*MLC1*) **negative**
 MEROSIN-DEFICIENT MUSCULAR DYSTROPHY (*LAMA2*) **negative**
 METABOLIC ENCEPHALOPATHY AND ARRHYTHMIAS, TANGO2-RELATED (*TANGO2*) **negative**
 METACHROMATIC LEUKODYSTROPHY, ARSA-RELATED (*ARSA*) **negative**
 METACHROMATIC LEUKODYSTROPHY, PSAP-RELATED (*PSAP*) **negative**
 METHYLMALONIC ACIDEMIA AND HOMOCYSTINURIA TYPE CBLF (*LMBRD1*) **negative**
 METHYLMALONIC ACIDEMIA, MCEE-RELATED (*MCEE*) **negative**
 METHYLMALONIC ACIDURIA AND HOMOCYSTINURIA, TYPE CBLC (*MMACHC*) **negative**
 METHYLMALONIC ACIDURIA AND HOMOCYSTINURIA, TYPE CblD (*MMADHC*) **negative**
 METHYLMALONIC ACIDURIA, MMAA-RELATED (*MMAA*) **negative**
 METHYLMALONIC ACIDURIA, MMAB-RELATED (*MMAB*) **negative**
 METHYLMALONIC ACIDURIA, TYPE MUT (0) (*MUT*) **negative**
 MEVALONIC KINASE DEFICIENCY (*MVK*) **negative**
 MICROCEPHALIC OSTEOPLASTIC PRIMORDIAL DWARFISM TYPE II (*PCNT*) **negative**
 MICROPTHALMIA / ANOPHTHALMIA, VSX2-RELATED (*VSX2*) **negative**
 MITOCHONDRIAL COMPLEX 1 DEFICIENCY, ACAD9-RELATED (*ACAD9*) **negative**
 MITOCHONDRIAL COMPLEX 1 DEFICIENCY, NDUFAF5-RELATED (*NDUFAF5*) **negative**
 MITOCHONDRIAL COMPLEX 1 DEFICIENCY, NDUFS6-RELATED (*NDUFS6*) **negative**
 MITOCHONDRIAL COMPLEX I DEFICIENCY, NUCLEAR TYPE 1 (*NDUFS4*) **negative**
 MITOCHONDRIAL COMPLEX I DEFICIENCY, NUCLEAR TYPE 10 (*NDUFAF2*) **negative**
 MITOCHONDRIAL COMPLEX I DEFICIENCY, NUCLEAR TYPE 17 (*NDUFAF6*) **negative**
 MITOCHONDRIAL COMPLEX I DEFICIENCY, NUCLEAR TYPE 19 (*FOXRED1*) **negative**
 MITOCHONDRIAL COMPLEX I DEFICIENCY, NUCLEAR TYPE 3 (*NDUFS7*) **negative**
 MITOCHONDRIAL COMPLEX I DEFICIENCY, NUCLEAR TYPE 4 (*NDUFV1*) **negative**
 MITOCHONDRIAL COMPLEX IV DEFICIENCY, NUCLEAR TYPE 2, SCO2-RELATED (*SCO2*) **negative**
 MITOCHONDRIAL COMPLEX IV DEFICIENCY, NUCLEAR TYPE 6 (*COX15*) **negative**
 MITOCHONDRIAL DNA DEPLETION SYNDROME 2 (*TK2*) **negative**

MITOCHONDRIAL DNA DEPLETION SYNDROME 3 (*DGUOK*) **negative**
 MITOCHONDRIAL MYOPATHY AND SIDEROBLASTIC ANEMIA (MLASA1) (*PUS1*) **negative**
 MITOCHONDRIAL TRIFUNCTIONAL PROTEIN DEFICIENCY, HADHB-RELATED (*HADHB*) **negative**
 MOLYBDENUM COFACTOR DEFICIENCY TYPE B (*MOC52*) **negative**
 MOLYBDENUM COFACTOR DEFICIENCY, TYPE A (*MOC51*) **negative**
 MUCOLIPIDOSIS II/III A (*GNPTAB*) **negative**
 MUCOLIPIDOSIS III GAMMA (*GNPTG*) **negative**
 MUCOLIPIDOSIS, TYPE IV (*MCOLN1*) **negative**
 MUCOPOLYSACCHARIDOSIS, TYPE I (HURLER SYNDROME) (*IDUA*) **negative**
 MUCOPOLYSACCHARIDOSIS, TYPE III A (SANFILIPPO A) (*SGSH*) **negative**
 MUCOPOLYSACCHARIDOSIS, TYPE III B (SANFILIPPO B) (*NAGLU*) **negative**
 MUCOPOLYSACCHARIDOSIS, TYPE III C (SANFILIPPO C) (*HGSNAT*) **negative**
 MUCOPOLYSACCHARIDOSIS, TYPE III D (SANFILIPPO D) (*GNS*) **negative**
 MUCOPOLYSACCHARIDOSIS, TYPE VI (MORQUIO SYNDROME) (*GALNS*) **negative**
 MUCOPOLYSACCHARIDOSIS, TYPE IV B/GM1 GANGLIOSIDOSIS (*GLB1*) **negative**
 MUCOPOLYSACCHARIDOSIS, TYPE IX (*HYAL1*) **negative**
 MUCOPOLYSACCHARIDOSIS, TYPE VI (MAROTEAUX-LAMY) (*ARSB*) **negative**
 MUCOPOLYSACCHARIDOSIS, TYPE VII (*GUSB*) **negative**
 MULIBREY NANISM (*TRIM37*) **negative**
 MULTIPLE PTERYGIUM SYNDROME, CHRNG-RELATED/ESCOBAR SYNDROME (*CHRNG*) **negative**
 MULTIPLE SULFATASE DEFICIENCY (*SUMF1*) **negative**
 MUSCLE-EYE-BRAIN DISEASE, POMGNT1-RELATED (*POMGNT1*) **negative**
 MUSCULAR DYSTROPHY-DYSTROGLYCANOPATHY (*RXYLT1*) **negative**
 MUSK-RELATED CONGENITAL MYASTHENIC SYNDROME (*MUSK*) **negative**
 MYONEUROGASTROINTESTINAL ENCEPHALOPATHY (MNGIE) (*TYMP*) **negative**
 MYOTONIA CONGENITA (*CLCN1*) **negative**

N
 N-ACETYLGUTAMATE SYNTHASE DEFICIENCY (*NAGS*) **negative**
 NEMALINE MYOPATHY, NEB-RELATED (*NEB*) **negative**
 NEPHRONOPHTHISIS 1 (*NPHP1*) **negative**
 NEURONAL CEROID LIPOFUSCINOSIS, CLN5-RELATED (*CLN5*) **negative**
 NEURONAL CEROID LIPOFUSCINOSIS, CLN6-RELATED (*CLN6*) **negative**
 NEURONAL CEROID LIPOFUSCINOSIS, CLN8-RELATED (*CLN8*) **negative**
 NEURONAL CEROID LIPOFUSCINOSIS, MFSD8-RELATED (*MFSD8*) **negative**
 NEURONAL CEROID LIPOFUSCINOSIS, PPT1-RELATED (*PPT1*) **negative**
 NEURONAL CEROID LIPOFUSCINOSIS, TPP1-RELATED (*TPP1*) **negative**
 NGLY1-CONGENITAL DISORDER OF GLYCOSYLATION (*NGLY1*) **negative**
 NIEMANN-PICK DISEASE, TYPE C1 / D (*NPC1*) **negative**
 NIEMANN-PICK DISEASE, TYPE C2 (*NPC2*) **negative**
 NIEMANN-PICK DISEASE, TYPES A / B (*SMPD1*) **negative**
 NIJMEGEN BREAKAGE SYNDROME (*NBN*) **negative**
 NON-SYNDROMIC HEARING LOSS, GJB2-RELATED (*GJB2*) **see first page**
 NON-SYNDROMIC HEARING LOSS, MYO15A-RELATED (*MYO15A*) **negative**
 NONSYNDROMIC HEARING LOSS, OTOA-RELATED (*OTOA*) **negative**
 NONSYNDROMIC HEARING LOSS, OTOF-RELATED (*OTOF*) **negative**
 NONSYNDROMIC HEARING LOSS, PJK-RELATED (*PJK*) **negative**
 NONSYNDROMIC HEARING LOSS, SYNE4-RELATED (*SYNE4*) **negative**
 NONSYNDROMIC HEARING LOSS, TMC1-RELATED (*TMC1*) **negative**
 NONSYNDROMIC HEARING LOSS, TMPRSS3-RELATED (*TMPRSS3*) **negative**
 NONSYNDROMIC INTELLECTUAL DISABILITY (*CC2D1A*) **negative**
 NORMOPHOSPHATEMIC TUMORAL CALCINOSIS (*SAMD9*) **negative**

O
 OCULOCUTANEOUS ALBINISM TYPE IV (*SLC45A2*) **negative**
 OCULOCUTANEOUS ALBINISM TYPE, III (*TYRP1*) **negative**
 OCULOCUTANEOUS ALBINISM, OCA2-RELATED (*OCA2*) **negative**
 OCULOCUTANEOUS ALBINISM, TYPES 1A AND 1B (*TYR*) **negative**
 ODONTO-ONYCHO-DERMAL DYSPLASIA / SCHOPF-SCHULZ-PASSARGE SYNDROME (*WNT10A*) **negative**
 OMENN SYNDROME, RAG2-RELATED (*RAG2*) **negative**
 ORNITHINE AMINOTRANSFERASE DEFICIENCY (*OAT*) **negative**
 OSTEOGENESIS IMPERFECTA TYPE VII (*CRTAP*) **negative**
 OSTEOGENESIS IMPERFECTA TYPE VIII (*P3H1*) **negative**
 OSTEOGENESIS IMPERFECTA TYPE XI (*FKBP10*) **negative**
 OSTEOGENESIS IMPERFECTA TYPE XIII (*BMP1*) **negative**
 OSTEOPECTOSIS, INFANTILE MALIGNANT, TCIRG1-RELATED (*TCIRG1*) **negative**
 OSTEOPECTOSIS, OSTM1-RELATED (*OSTM1*) **negative**

P
 PANTOTHENATE KINASE-ASSOCIATED NEURODEGENERATION (*PANK2*) **negative**
 PAPILLON LEFÈVRE SYNDROME (*CTSC*) **negative**
 PARKINSON DISEASE 15 (*FBXO7*) **negative**
 PENDRED SYNDROME (*SLC26A4*) **negative**
 PERLMAN SYNDROME (*DIS3L2*) **negative**
 PGM3-CONGENITAL DISORDER OF GLYCOSYLATION (*PGM3*) **negative**
 PHENYLKETONURIA (*PAH*) **negative**
 PIGN-CONGENITAL DISORDER OF GLYCOSYLATION (*PIGN*) **negative**
 PITUITARY HORMONE DEFICIENCY, COMBINED 3 (*LHX3*) **negative**
 POLG-RELATED DISORDERS (*POLG*) **negative**

Patient Information

Patient Name:

Test Information

Ordering Physician:



Clinic Information:

Date Of Birth:

Case File ID:

Report Date:

P

POLYCYSTIC KIDNEY DISEASE, AUTOSOMAL RECESSIVE (*PKHD1*) **negative**
 PONTOCEREBELLAR HYPOPLASIA, EXOSC3-RELATED (*EXOSC3*) **negative**
 PONTOCEREBELLAR HYPOPLASIA, RARS2-RELATED (*RARS2*) **negative**
 PONTOCEREBELLAR HYPOPLASIA, TSEN2-RELATED (*TSEN2*) **negative**
 PONTOCEREBELLAR HYPOPLASIA, TSEN54-RELATED (*TSEN54*) **negative**
 PONTOCEREBELLAR HYPOPLASIA, TYPE 1A (*VRK1*) **negative**
 PONTOCEREBELLAR HYPOPLASIA, TYPE 2D (*SEPSEC5*) **negative**
 PONTOCEREBELLAR HYPOPLASIA, VPS53-RELATED (*VPS53*) **negative**
 PRIMARY CILIARY DYSKINESIA, CCDC103-RELATED (*CCDC103*) **negative**
 PRIMARY CILIARY DYSKINESIA, CCDC39-RELATED (*CCDC39*) **negative**
 PRIMARY CILIARY DYSKINESIA, DNAH11-RELATED (*DNAH11*) **negative**
 PRIMARY CILIARY DYSKINESIA, DNAH5-RELATED (*DNAH5*) **negative**
 PRIMARY CILIARY DYSKINESIA, DNAI1-RELATED (*DNAI1*) **negative**
 PRIMARY CILIARY DYSKINESIA, DNAI2-RELATED (*DNAI2*) **negative**
 PRIMARY CONGENITAL GLAUCOMA/PETERS ANOMALY (*CYP1B1*) **negative**
 PRIMARY HYPEROXALURIA, TYPE 1 (*AGXT*) **negative**
 PRIMARY HYPEROXALURIA, TYPE 2 (*GRHPR*) **negative**
 PRIMARY HYPEROXALURIA, TYPE 3 (*HOGA1*) **negative**
 PRIMARY MICROCEPHALY 1, AUTOSOMAL RECESSIVE (*MCPH1*) **negative**
 PROGRESSIVE EARLY-ONSET ENCEPHALOPATHY WITH BRAIN ATROPHY AND THIN CORPUS CALLOSUM (*TBCD*) **negative**
 PROGRESSIVE FAMILIAL INTRAHEPATIC CHOLESTASIS, ABCB4-RELATED (*ABCB4*) **negative**
 PROGRESSIVE FAMILIAL INTRAHEPATIC CHOLESTASIS, TYPE 1 (*PFIC1*) (*ATP8B1*) **negative**
 PROGRESSIVE FAMILIAL INTRAHEPATIC CHOLESTASIS, TYPE 2 (*ABCB11*) **negative**
 PROGRESSIVE FAMILIAL INTRAHEPATIC CHOLESTASIS, TYPE 4 (*PFIC4*) (*TJP2*) **negative**
 PROGRESSIVE PSEUDORHEUMATOID DYSPLASIA (*CCN6*) **negative**
 PROLIDASE DEFICIENCY (*PEPD*) **negative**
 PROPIONIC ACIDEMIA, PCCA-RELATED (*PCCA*) **negative**
 PROPIONIC ACIDEMIA, PCCB-RELATED (*PCCB*) **negative**
 PSEUDOXANTHOMA ELASTICUM (*ABCC6*) **negative**
 PTERIN-4 ALPHA-CARBINOLAMINE DEHYDRATASE (*PCD*) DEFICIENCY (*PCBD1*) **negative**
 PYCNODYSTOSIS (*CTS5*) **negative**
 PYRIDOXAL 5'-PHOSPHATE-DEPENDENT EPILEPSY (*PNPO*) **negative**
 PYRIDOXINE-DEPENDENT EPILEPSY (*ALDH7A1*) **negative**
 PYRUVATE CARBOXYLASE DEFICIENCY (*PC*) **negative**
 PYRUVATE DEHYDROGENASE DEFICIENCY, PDHB-RELATED (*PDHB*) **negative**

R

REFSUM DISEASE, PHYH-RELATED (*PHYH*) **negative**
 RENAL TUBULAR ACIDOSIS AND DEAFNESS, ATP6V1B1-RELATED (*ATP6V1B1*) **negative**
 RENAL TUBULAR ACIDOSIS, PROXIMAL, WITH OCULAR ABNORMALITIES AND MENTAL RETARDATION (*SLC4A4*) **negative**
 RETINITIS PIGMENTOSA 25 (*EYS*) **negative**
 RETINITIS PIGMENTOSA 26 (*CERKL*) **negative**
 RETINITIS PIGMENTOSA 28 (*FAM161A*) **negative**
 RETINITIS PIGMENTOSA 36 (*PRCD*) **negative**
 RETINITIS PIGMENTOSA 59 (*DHDDS*) **negative**
 RETINITIS PIGMENTOSA 62 (*MAK*) **negative**
 RHIZOMELIC CHONDRODYSPLASIA PUNCTATA, TYPE 1 (*PEX7*) **negative**
 RHIZOMELIC CHONDRODYSPLASIA PUNCTATA, TYPE 2 (*GNPAT*) **negative**
 RHIZOMELIC CHONDRODYSPLASIA PUNCTATA, TYPE 3 (*AGPS*) **negative**
 RLBP1-RELATED RETINOPATHY (*RLBP1*) **negative**
 ROBERTS SYNDROME (*ESCO2*) **negative**
 RYR1-RELATED CONDITIONS (*RYR1*) **negative**

S

SALLA DISEASE (*SLC17A5*) **negative**
 SANDHOFF DISEASE (*HEXB*) **negative**
 SCHIMKE IMMUNOSSESQUOUS DYSPLASIA (*SMARCAL1*) **negative**
 SCHINDLER DISEASE (*NAGA*) **negative**
 SEGAWA SYNDROME, TH-RELATED (*TH*) **negative**
 SENIOR-LOKEN SYNDROME 4/NEPHRONOPHTHISIS 4 (*NPHP4*) **negative**
 SEPIAPTERIN REDUCTASE DEFICIENCY (*SPR*) **negative**
 SEVERE COMBINED IMMUNODEFICIENCY (SCID), CD3D-RELATED (*CD3D*) **negative**
 SEVERE COMBINED IMMUNODEFICIENCY (SCID), CD3E-RELATED (*CD3E*) **negative**
 SEVERE COMBINED IMMUNODEFICIENCY (SCID), FOXP1-RELATED (*FOXP1*) **negative**
 SEVERE COMBINED IMMUNODEFICIENCY (SCID), IKBKB-RELATED (*IKBKB*) **negative**
 SEVERE COMBINED IMMUNODEFICIENCY (SCID), IL7R-RELATED (*IL7R*) **negative**
 SEVERE COMBINED IMMUNODEFICIENCY (SCID), JAK3-RELATED (*JAK3*) **negative**
 SEVERE COMBINED IMMUNODEFICIENCY (SCID), PTPRC-RELATED (*PTPRC*) **negative**
 SEVERE COMBINED IMMUNODEFICIENCY (SCID), RAG1-RELATED (*RAG1*) **negative**
 SEVERE COMBINED IMMUNODEFICIENCY, ADA-Related (*ADA*) **negative**
 SEVERE COMBINED IMMUNODEFICIENCY, TYPE ATHABASKAN (*DCLRE1C*) **negative**
 SHORT-RIB THORACIC DYSPLASIA 3 WITH OR WITHOUT POLYDACTYLY (*DYNC2H1*) **negative**
 SHWACHMAN-DIAMOND SYNDROME, SBDS-RELATED (*SBDS*) **negative**
 SIALIDOSIS (*NEU1*) **negative**
 SJÖGREN-LARSSON SYNDROME (*ALDH3A2*) **negative**
 SMITH-LEMLI-OPITZ SYNDROME (*DHCR7*) **negative**
 SPASTIC PARAPLEGIA, TYPE 15 (*ZFYVE26*) **negative**

SPASTIC TETRAPLEGIA, THIN CORPUS CALLOSUM, AND PROGRESSIVE MICROCEPHALY (SPATCCM) (*SLC1A4*) **negative**
 SPG11-RELATED CONDITIONS (*SPG11*) **negative**
 SPINAL MUSCULAR ATROPHY (*SMN1*) **negative** SMN1: Two copies; g.27134T>G: absent; the absence of the g.27134T>G variant decreases the chance to be a silent (2+0) carrier.
 SPINAL MUSCULAR ATROPHY WITH RESPIRATORY DISTRESS TYPE 1 (*IGHMBP2*) **negative**
 SPINOCEREBELLAR ATAXIA, AUTOSOMAL RECESSIVE 10 (*ANO10*) **negative**
 SPINOCEREBELLAR ATAXIA, AUTOSOMAL RECESSIVE 12 (*WWOX*) **negative**
 SPONDYLOCOSTAL DYSOSTOSIS 1 (*DLL3*) **negative**
 SPONDYLOTHORACIC DYSOSTOSIS, MESP2-Related (*MESP2*) **negative**
 STEEL SYNDROME (*COL27A1*) **negative**
 STEROID-RESISTANT NEPHROTIC SYNDROME (*NPHS2*) **see first page**
 STUVE-WIEDEMANN SYNDROME (*LIFR*) **negative**
 SURF1-RELATED CONDITIONS (*SURF1*) **negative**
 SURFACTANT DYSFUNCTION, ABCA3-RELATED (*ABCA3*) **negative**

T

TAY-SACHS DISEASE (*HEXA*) **negative**
 TBCE-RELATED CONDITIONS (*TBCE*) **negative**
 THIAMINE-RESPONSIVE MEGALOBlastic ANEMIA SYNDROME (*SLC19A2*) **negative**
 THYROID DYSHORMONOGENESIS 1 (*SLC5A5*) **negative**
 THYROID DYSHORMONOGENESIS 2A (*TPO*) **negative**
 THYROID DYSHORMONOGENESIS 3 (*TG*) **negative**
 THYROID DYSHORMONOGENESIS 6 (*DUOX2*) **negative**
 TRANSCOBALAMIN II DEFICIENCY (*TCN2*) **negative**
 TRICHOHEPATOENTERIC SYNDROME, SKIC2-RELATED (*SKIC2*) **negative**
 TRICHOHEPATOENTERIC SYNDROME, TTC37-RELATED (*TTC37*) **negative**
 TRICHOHYDROSTROPHY 1/XERODERMA PIGMENTOSUM, GROUP D (*ERCC2*) **negative**
 TRIMETHYLAMINURIA (*FMO3*) **negative**
 TRIPLE A SYNDROME (*AAAS*) **negative**
 TSHR-RELATED CONDITIONS (*TSHR*) **negative**
 TYROSINEMIA TYPE III (*HPD*) **negative**
 TYROSINEMIA, TYPE 1 (*FAH*) **negative**
 TYROSINEMIA, TYPE 2 (*TAT*) **negative**

U

USHER SYNDROME, TYPE 1B (*MYO7A*) **negative**
 USHER SYNDROME, TYPE 1C (*USH1C*) **negative**
 USHER SYNDROME, TYPE 1D (*CDH23*) **negative**
 USHER SYNDROME, TYPE 1F (*PCDH15*) **negative**
 USHER SYNDROME, TYPE 1J/DEAFNESS, AUTOSOMAL RECESSIVE, 48 (*CIB2*) **negative**
 USHER SYNDROME, TYPE 2A (*USH2A*) **negative**
 USHER SYNDROME, TYPE 2C (*ADGRV1*) **negative**
 USHER SYNDROME, TYPE 3 (*CLRN1*) **negative**

V

VERY LONG-CHAIN ACYL-CoA DEHYDROGENASE DEFICIENCY (*ACADVL*) **negative**
 VICI SYNDROME (*EPG5*) **negative**
 VITAMIN D-DEPENDENT RICKETS, TYPE 1A (*CYP27B1*) **negative**
 VITAMIN D-RESISTANT RICKETS TYPE 2A (*VDR*) **negative**
 VLDLR-ASSOCIATED CEREBELLAR HYPOPLASIA (*VLDLR*) **negative**

W

WALKER-WARBURG SYNDROME, CRPPA-RELATED (*CRPPA*) **negative**
 WALKER-WARBURG SYNDROME, FKTN-RELATED (*FKTN*) **negative**
 WALKER-WARBURG SYNDROME, LARGE1-RELATED (*LARGE1*) **negative**
 WALKER-WARBURG SYNDROME, POMT1-RELATED (*POMT1*) **negative**
 WALKER-WARBURG SYNDROME, POMT2-RELATED (*POMT2*) **negative**
 WARSAW BREAKAGE SYNDROME (*DDX11*) **negative**
 WERNER SYNDROME (*WRN*) **negative**
 WILSON DISEASE (*ATP7B*) **negative**
 WOLCOTT-RALLISON SYNDROME (*EIF2AK3*) **negative**
 WOLMAN DISEASE (*LIPA*) **negative**
 WOODHOUSE-SAKATI SYNDROME (*DCAF17*) **negative**

X

XERODERMA PIGMENTOSUM VARIANT TYPE (*POLH*) **negative**
 XERODERMA PIGMENTOSUM, GROUP A (*XPA*) **negative**
 XERODERMA PIGMENTOSUM, GROUP C (*XPC*) **negative**

Z

ZELLWEGER SPECTRUM DISORDER, PEX13-RELATED (*PEX13*) **negative**
 ZELLWEGER SPECTRUM DISORDER, PEX16-RELATED (*PEX16*) **negative**
 ZELLWEGER SPECTRUM DISORDER, PEX5-RELATED (*PEX5*) **negative**
 ZELLWEGER SPECTRUM DISORDERS, PEX10-RELATED (*PEX10*) **negative**
 ZELLWEGER SPECTRUM DISORDERS, PEX12-RELATED (*PEX12*) **negative**
 ZELLWEGER SPECTRUM DISORDERS, PEX1-RELATED (*PEX1*) **negative**
 ZELLWEGER SPECTRUM DISORDERS, PEX26-RELATED (*PEX26*) **negative**
 ZELLWEGER SPECTRUM DISORDERS, PEX2-RELATED (*PEX2*) **negative**

Patient Information

Patient Name:

Test Information

Ordering Physician:



Date Of Birth:



Clinic Information:

Case File ID:



Report Date:

Z

ZELLWEGER SPECTRUM DISORDERS, PEX6-RELATED (PEX6) **negative**

Patient Information

Patient Name: [REDACTED]

Test Information

Ordering Physician: [REDACTED]



Clinic Information:

Date Of Birth: [REDACTED]

Case File ID: [REDACTED]

Report Date:

Testing Methodology, Limitations, and Comments:**Next-generation sequencing (NGS)**

Sequencing library prepared from genomic DNA isolated from a patient sample is enriched for targets of interest using standard hybridization capture protocols and PCR amplification (for targets specified below). NGS is then performed to achieve the standards of quality control metrics, including a minimum coverage of 99% of targeted regions at 20X sequencing depth. Sequencing data is aligned to human reference sequence, followed by deduplication, metric collection and variant calling (coding region +/- 20bp). Variants are then classified according to ACMGG/AMP standards of interpretation using publicly available databases including but not limited to ENSEMBL, HGMD Pro, ClinGen, ClinVar, 1000G, ESP and gnomAD. Variants predicted to be pathogenic or likely pathogenic for the specified diseases are reported. It should be noted that the data interpretation is based on our current understanding of the genes and variants at the time of reporting. Putative positive sequencing variants that do not meet internal quality standards or are within highly homologous regions are confirmed by Sanger sequencing or gene-specific long-range PCR as needed prior to reporting.

Copy Number Variant (CNV) analysis is limited to deletions involving two or more exons for all genes on the panel, in addition to specific known recurrent single-exon deletions. CNVs of small size may have reduced detection rate. This method does not detect gene inversions, single-exonic and sub-exonic deletions (unless otherwise specified), and duplications of all sizes (unless otherwise specified). Additionally, this method does not define the exact breakpoints of detected CNV events. Confirmation testing for copy number variation is performed by specific PCR, Multiplex Ligation-dependent Probe Amplification (MLPA), next generation sequencing, or other methodology.

This test may not detect certain variants due to local sequence characteristics, high/low genomic complexity, homologous sequence, or allele dropout (PCR-based assays). Variants within noncoding regions (promoter, 5'UTR, 3'UTR, deep intronic regions, unless otherwise specified), small deletions or insertions larger than 25bp, low-level mosaic variants, structural variants such as inversions, and/or balanced translocations may not be detected with this technology.

SPECIAL NOTES

For ABCC6, variants in exons 1-9 are not detected due to the presence of regions of high homology.

For CFTR, when the CFTR R117H variant is detected, reflex analysis of the polythymidine variations (5T, 7T and 9T) at the intron 9 branch/acceptor site of the CFTR gene will be performed.

For CYP21A2, targets were enriched using long-range PCR amplification, followed by next generation sequencing. Duplication analysis will only be performed and reported when c.955C>T (p.Q319*) is detected. Sequencing and CNV analysis may have reduced sensitivity, if variants result from complex rearrangements, in trans with a gene deletion, or CYP21A2 gene duplication on one chromosome and deletion on the other chromosome. This analysis cannot detect sequencing variants located on the CYP21A2 duplicated copy.

For DDX11, only NM_030653.3:c.1763 - 1G > C variant will be analyzed and reported.

For HBA1/HBA2, CNV analysis is offered to detect common deletions of -alpha3.7, -alpha4.2, --MED, --SEA, --FIL, --THAI, --alpha20.5, and/or HS-40.

For OTOA, variants in exons 20 - 28 are not analyzed due to high sequence homology.

For RPGRIP1L, variants in exon 23 are not detected due to assay limitation.

For SAMD9, only p.K1495E variant will be analyzed and reported.

Friedreich Ataxia (FXN)

The GAA repeat region of the FXN gene is assessed by trinucleotide PCR assay and capillary electrophoresis. Variances of +/-1 repeat for normal alleles and up to +/-3 repeats for premutation alleles may occur. For fully penetrant expanded alleles, the precise repeat size cannot be determined, therefore the approximate allele size is reported. Sequencing and copy number variants are analyzed by next-generation sequencing analysis.

Friedreich Ataxia Repeat Categories

Categories	GAA Repeat Sizes
Normal	<34
Premutation	34 - 65
Full	>65

Patient Information

Patient Name: [REDACTED]

Test Information

Ordering Physician: [REDACTED]



Date Of Birth: [REDACTED]

Clinic Information:

Case File ID: [REDACTED]

Report Date:

Spinal Muscular Atrophy (SMN1)

The total combined copy number of SMN1 and SMN2 exon 7 is quantified based on NGS read depth. The ratio of SMN1 to SMN2 is calculated based on the read depth of a single nucleotide that distinguishes these two genes in exon 7. In addition to copy number analysis, testing for the presence or absence of a single nucleotide polymorphism (g.27134T>G in intron 7 of SMN1) associated with the presence of a SMN1 duplication allele is performed using NGS.

Ethnicity	Two SMN1 copies carrier risk before g.27134T>G testing	Carrier risk after g.27134T>G testing	
		g.27134T>G ABSENT	g.27134T>G PRESENT
Caucasian	1 in 632	1 in 769	1 in 29
Ashkenazi Jewish	1 in 350	1 in 580	LIKELY CARRIER
Asian	1 in 628	1 in 702	LIKELY CARRIER
African-American	1 in 121	1 in 396	1 in 34
Hispanic	1 in 1061	1 in 1762	1 in 140

Variant Classification

Only pathogenic or likely pathogenic variants are reported. Other variants including benign variants, likely benign variants, variants of uncertain significance, or inconclusive variants identified during this analysis may be reported in certain circumstances. Our laboratory's variant classification criteria are based on the ACMG and internal guidelines and our current understanding of the specific genes. This interpretation may change over time as more information about a gene and/or variant becomes available. Natera and its lab partner(s) may reclassify variants at certain intervals but may not release updated reports without a specific request made to Natera by the ordering provider. Natera may disclose incidental findings if deemed clinically pertinent to the test performed.

Negative Results

A negative carrier screening result reduces the risk for a patient to be a carrier of a specific disease but does not completely rule out carrier status. Please visit <https://www.natera.com/panel-option/h-all/> for a table of carrier rates, detection rates, residual risks and promised variants/exons per gene. Carrier rates before and after testing vary by ethnicity and assume a negative family history for each disease screened and the absence of clinical symptoms in the patient. Any patient with a family history for a specific genetic disease will have a higher carrier risk prior to testing and, if the disease-causing mutation in their family is not included on the test, their carrier risk would remain unchanged. Genetic counseling is recommended for patients with a family history of genetic disease so that risk figures based on actual family history can be determined and discussed along with potential implications for reproduction. Horizon carrier screening has been developed to identify the reproductive risks for monogenic inherited conditions. Even when one or both members of a couple screen negative for pathogenic variants in a specific gene, the disease risk for their offspring is not zero. There is still a low risk for the condition in their offspring due to a number of different mechanisms that are not detected by Horizon including, but not limited to, pathogenic variant(s) in the tested gene or in a different gene not included on Horizon, pathogenic variant(s) in an upstream regulator, uniparental disomy, de novo mutation(s), or digenic or polygenic inheritance.

Additional Comments

These analyses generally provide highly accurate information regarding the patient's carrier status. Despite this high level of accuracy, it should be kept in mind that there are many potential sources of diagnostic error, including misidentification of samples, polymorphisms, or other rare genetic variants that interfere with analysis. Families should understand that rare diagnostic errors may occur for these reasons.

7414, DONOR ▲

DOB: [REDACTED] Age: [REDACTED] Specimen: [REDACTED] Collected: 03/18/2024 11:00
 Sex: M Fasting: [REDACTED] Requisition: 0000772 [REDACTED] Received: 03/19/2024 13:18
 Phone: [REDACTED] Lab Reference: [REDACTED] Reported: 03/25/2024 19:40
 Patient ID: 7414- [REDACTED] Report Status: FINAL / SEE REPORT



=PHILLIPINES, THAILAND


▲ Hemoglobinopathy Evaluation

FINAL

Lab: AMD

Hemoglobinopathy Evaluation	No Historical Data	FINAL
Red Blood Cell Count Reference Range: 4.20-5.80 Mill/uL	5.27 4.20 5.80 No Historical Data	FINAL
HEMOGLOBIN Reference Range: 13.2-17.1 g/dL	16.8 13.2 17.1 No Historical Data	FINAL
Hematocrit	No Historical Data	FINAL
▲ Hematocrit Reference Range: 38.5-50.0 %	51.1 38.5 50.0 No Historical Data	FINAL
MCV Reference Range: 80.0-100.0 fL	97.0 80.0 100.0 No Historical Data	FINAL
MCH Reference Range: 27.0-33.0 pg	31.9 27.0 33.0 No Historical Data	FINAL
RDW Reference Range: 11.0-15.0 %	12.7 11.0 15.0 No Historical Data	FINAL
Hemoglobinopathy Evaluation	No Historical Data	FINAL
Hemoglobin A Reference Range: >96.0 %	97.4 >96.0 No Historical Data	FINAL
Hemoglobin F Reference Range: <2.0 %	0.0 <2.0 No Historical Data	FINAL

Hemoglobin A2 (Quant)
Reference Range: 2.2-3.2 %



FINAL

No Historical Data

Interpretation

No Historical Data

FINAL

NORMAL PATTERN

There is a normal pattern of hemoglobins and normal levels of Hb A2 and Hb F are present. No variant hemoglobins are observed. This is consistent with A/A phenotype. If iron deficiency coexists with a mild/silent beta thalassemia trait Hb A2 may be in the normal range. Rare variant hemoglobins have no separation from hemoglobin A by capillary zone electrophoresis (CZE) or high-performance liquid chromatography (HPLC). If clinically indicated, Thalassemia and Hemoglobinopathy Comprehensive (TC 17365) should be considered.

▲ CBC (includes Differential and Platelets)

FINAL

Lab: AMD


White Blood Cell Count
Reference Range: 3.8-10.8 Thous/uL



FINAL

No Historical Data

Red Blood Cell Count
Reference Range: 4.20-5.80 Mill/uL



FINAL

No Historical Data

HEMOGLOBIN
Reference Range: 13.2-17.1 g/dL



FINAL

No Historical Data


▲ Hematocrit
Reference Range: 38.5-50.0 %



FINAL

No Historical Data


MCV
Reference Range: 80.0-100.0 fL



FINAL

No Historical Data


MCH
Reference Range: 27.0-33.0 pg



FINAL

No Historical Data


MCHC
Reference Range: 32.0-36.0 g/dL



FINAL

No Historical Data

RDW
Reference Range: 11.0-15.0 %



FINAL

No Historical Data

Chromosome Analysis, Blood

FINAL

Lab: AMD

Chromosome Analysis, Blood

No Historical Data

FINAL

Order ID:



Specimen Type: Blood

Clinical Indication: Gamete donor

RESULT:
NORMAL MALE KARYOTYPE

INTERPRETATION:
Chromosome analysis revealed normal G-band patterns within the limits of standard cytogenetic analysis.

Please expect the results of any other concurrent study in a separate report.

NOMENCLATURE:
46,XY

ASSAY INFORMATION:
Method: G-Band (Digital Analysis:
MetaSystems/Ikaros)
Cells Counted: 20
Band Level: 550
Cells Analyzed: 5
Cells Karyotyped: 5

This test does not address genetic disorders that cannot be detected by standard cytogenetic methods or rare events such as low level mosaicism or subtle rearrangements.

Steven A. Schonberg, Ph.D., FACMG, Technical Director, Cytogenetics and Genomics, 703-802-7156

Electronic Signature: 3/25/2024 6:57 PM

For additional information, please refer to <http://education.questdiagnostics.com/faq/chromsblood> (This link is being provided for informational/educational purposes only).

Performing Sites

AMD Quest Diagnostics Nichols Institute, 14225 Newbrook Drive, Chantilly, VA 20151 Laboratory Director: Patrick W Mason, MD PhD

Key

Priority Out of Range Out of Range Pending Result Preliminary Result Final Result Reissued Result

Note: Data displayed only for results that meet strict identification matching. Historical result view may vary based on corrected or updated patient demographics. The reference range displayed may vary due to potential changes in laboratory testing methods. Please refer to the published reference range on each lab report.

Quest, Quest Diagnostics, the associated logo, Nichols Institute, Interactive Insights and all associated Quest Diagnostics marks are the registered trademarks of Quest Diagnostics. All third party marks - ® and ™ - are the property of their respective owners. Privacy policy can be found at: <http://questdiagnostics.com/home/privacy-policy/online-privacy.html>. © 2022 Quest Diagnostics Incorporated. All rights reserved.