



## Donor 7249

### Genetic Testing Summary

Fairfax Cryobank recommends reviewing this genetic testing summary with your healthcare provider to determine suitability.

Last Updated: 08/09/24

Donor Reported Ancestry: Irish, Polish, German

Jewish Ancestry: No

| Genetic Test*   | Result  | Comments/Donor's Residual Risk**   |
|---|---|--|
| Chromosome analysis (karyotype)   | Normal male karyotype   | No evidence of clinically significant chromosome abnormalities   |
| Hemoglobin evaluation   | Normal hemoglobin fractionation and MCV/MCH results   | Reduced risk to be a carrier for sickle cell anemia, beta thalassemia, alpha thalassemia trait (aa/-- and a-/a-) and other hemoglobinopathies  |
| Expanded Genetic Disease Carrier Screening Panel attached- 514 diseases by gene sequencing. | <p>Carrier: ABCA4-related conditions (ABCA4)</p> <p>Carrier: Nonsyndromic deafness (TMPRSS3-related)</p> <p>Carrier: Sandhoff disease (HEXB)</p> <p>Negative for other genes sequenced.</p> | <p>Partner testing is recommended before using this donor.</p> <p>Residual risks for negative results can be seen here:</p> <p><a href="https://fairfaxcryobank.com/invitae-residual-risk-table">https://fairfaxcryobank.com/invitae-residual-risk-table</a></p> |
| <b>Special Testing</b>  |   |  |
| Gene: SLC6A19   | Negative by gene sequencing   |  |

\*No single test can screen for all genetic disorders. A negative screening result significantly reduces, but cannot eliminate, the risk for these conditions in a pregnancy.

\*\*Donor residual risk is the chance the donor is still a carrier after testing negative.

|  |  |                                  |
|--|--|----------------------------------|
| <b>Patient name:</b> Donor 7249          | <b>Sample type:</b> Blood                  | <b>Report date:</b> 30-OCT-2023  |
| <b>DOB:</b> [REDACTED]                   | <b>Sample collection date:</b> 20-OCT-2023 | <b>Invitae #:</b> [REDACTED]     |
| <b>Sex assigned at birth:</b> Male       | <b>Sample accession date:</b> 23-OCT-2023  | <b>Clinical team:</b> [REDACTED] |
| <b>Gender:</b>                           |  | [REDACTED]                       |
| <b>Patient ID (MRN):</b> 7249-[REDACTED] |  |                                  |

**Reason for testing**

Gamete donor

**Test performed**

Invitae Carrier Screen


**RESULT: POSITIVE**

This carrier test evaluated 514 gene(s) for genetic changes (variants) that are associated with an increased risk of having a child with a genetic condition. Knowledge of carrier status for one of these conditions may provide information that can be used to assist with family planning and/or preparation. Carrier screening is not intended for diagnostic purposes. To identify a potential genetic basis for a condition in the individual being tested, diagnostic testing for the gene(s) of interest is recommended.

This test shows the presence of clinically significant genetic change(s) in this individual in the gene(s) indicated below. No other clinically significant changes were identified in the remaining genes evaluated with this test.

| RESULTS   | GENE    | VARIANT(S)                | INHERITANCE         | PARTNER TESTING RECOMMENDED |
|---|---------|---------------------------|---------------------|-----------------------------|
| <b>Carrier:</b> ABCA4-related conditions                | ABCA4   | c.2588G>C (p.Gly863Ala) § | Autosomal recessive | Yes                         |
| <b>Carrier:</b> Nonsyndromic deafness (TMPRSS3-related) | TMPRSS3 | c.1028G>A (p.Trp343*)     | Autosomal recessive | Yes                         |
| <b>Carrier:</b> Sandhoff disease                        | HEXB    | Deletion (Exons 1-5)      | Autosomal recessive | Yes                         |

§ This variant is known to have low penetrance. See Clinical summary and/or Variant details on following pages for more information.

## Next steps

- See the table above for recommendations regarding testing of this individual's reproductive partner.
- Even for genes that have a negative test result, there is always a small risk that an individual could still be a carrier. This is called “residual risk.” See the Carrier detection rates and residual risks document.
- Discussion with a physician and/or genetic counselor is recommended to further review the implications of this test result and to understand these results in the context of any family history of a genetic condition.
- All patients, regardless of result, may wish to consider additional screening for hemoglobinopathies by complete blood count (CBC) and hemoglobin electrophoresis, if this has not already been completed.
- Individuals can register their tests at <https://www.invitae.com/patients/> to access online results, educational resources, and next steps.

## Clinical summary

### **RESULT: CARRIER**

#### **ABCA4-related conditions**

A single Pathogenic (low penetrance) variant, c.2588G>C (p.Gly863Ala), was identified in ABCA4.

#### **What are ABCA4-related conditions?**

ABCA4-related conditions are a spectrum of inherited retinal disorders that cause impaired vision.

Cone-rod dystrophy (CRD) typically presents during childhood or adolescence and symptoms become more severe over time. Symptoms include reduced visual acuity (farsightedness or nearsightedness), loss of color perception, increased sensitivity to light (photophobia), and difficulty seeing in low light settings (night blindness). Some affected individuals develop involuntary eye movements (nystagmus), and many are legally blind by mid-adulthood.

Stargardt disease typically presents during childhood to early adulthood, although the severity and progression are highly variable. Affected individuals experience symptoms including a dark spot appearing in the center of their vision, having difficulty reading, driving or recognizing faces, difficulty transitioning from an area of light to dark, and photophobia. Individuals can also develop problems with night or color vision over time. Upon retinal exam, there is a characteristic build up of an orange-yellow fatty substance called lipofuscin at the macula at the back of the eye, which is the part of the eye that is responsible for central vision.

Retinitis pigmentosa (RP) typically presents with night blindness, which usually occurs during childhood or adolescence. Vision loss continues over years or decades and typically progresses to a loss of side (peripheral) vision, causing tunnel vision. Ultimately, central vision loss occurs. Many individuals with RP are legally blind by adulthood, though the severity of symptoms and age of onset varies by individual.

Not everyone with a genetic change in ABCA4 will present the same; symptoms and severity can vary, even between family members with the same genetic change. Follow-up depends on each affected individual's specific situation, and discussion with a healthcare provider should be considered.

### Next steps


Carrier testing for the reproductive partner is recommended.

#### **If your partner tests positive:**

In autosomal recessive inheritance, an individual must have disease-causing genetic changes in each copy of the ABCA4 gene to be affected. Carriers, who have a disease-causing genetic change in only one copy of the gene, typically do not have symptoms. When both reproductive partners are carriers of an autosomal recessive condition, there is a 25% chance for each child to have the condition.

#### **If your partner tests negative:**

A negative carrier test result reduces, but does not eliminate, the chance that a person may be a carrier. The risk that a person could still be a carrier, even after a negative test result, is called a residual risk. See the table below for your partner's hypothetical residual risk after testing negative for ABCA4-related conditions. These values are provided only as a guide, are based on the detection rate for the condition as tested at Invitae, and assume a negative family history, the absence of symptoms, and vary based on the ethnic background of an individual. For genes associated with both dominant and recessive inheritance, the numbers provided apply to the recessive condition(s) associated with the gene.





Patient name: Donor 7249

DOB: [REDACTED]

Invitae #: [REDACTED]

| DISORDER (INHERITANCE)                       | GENE  | ETHNICITY  | CARRIER FREQUENCY BEFORE SCREENING | CARRIER RESIDUAL RISK AFTER NEGATIVE RESULT |
|--|-------|------------|------------------------------------|---|
| ABCA4-related conditions (AR)<br>NM_000350.2 | ABCA4 | Pan-ethnic | 1 in 45                            | 1 in 441                                    |


**RESULT: CARRIER**

## Nonsyndromic deafness (TMPRSS3-related)

A single Pathogenic variant, c.1028G>A (p.Trp343\*), was identified in TMPRSS3.

### What is nonsyndromic deafness (TMPRSS3-related)?

Nonsyndromic deafness is a condition that affects an individual's ability to hear. It can be caused by changes in several different genes. Nonsyndromic deafness does not affect any other part of the body. Affected individuals are born with mild to profound deafness that typically does not worsen over time. Severity of deafness may vary, even among members of the same family. Intellect and life span are not impacted. Follow-up depends on each affected individual's specific situation, and discussion with a healthcare provider should be considered.

### Next steps


Carrier testing for the reproductive partner is recommended.


**If your partner tests positive:**

In autosomal recessive inheritance, an individual must have disease-causing genetic changes in each copy of the TMPRSS3 gene to be affected. Carriers, who have a disease-causing genetic change in only one copy of the gene, typically do not have symptoms. When both reproductive partners are carriers of an autosomal recessive condition, there is a 25% chance for each child to have the condition.


**If your partner tests negative:**

A negative carrier test result reduces, but does not eliminate, the chance that a person may be a carrier. The risk that a person could still be a carrier, even after a negative test result, is called a residual risk. See the table below for your partner's hypothetical residual risk after testing negative for nonsyndromic deafness (TMPRSS3-related). These values are provided only as a guide, are based on the detection rate for the condition as tested at Invitae, and assume a negative family history, the absence of symptoms, and vary based on the ethnic background of an individual. For genes associated with both dominant and recessive inheritance, the numbers provided apply to the recessive condition(s) associated with the gene.



| DISORDER (INHERITANCE)                                      | GENE    | ETHNICITY  | CARRIER FREQUENCY BEFORE SCREENING | CARRIER RESIDUAL RISK AFTER NEGATIVE RESULT |
|---|---------|------------|------------------------------------|---|
| Nonsyndromic deafness (TMPRSS3-related) (AR)<br>NM_024022.2 | TMPRSS3 | Pan-ethnic | ≤1 in 500                          | Reduced                                     |


**RESULT: CARRIER**

## Sandhoff disease

A single Pathogenic variant, Deletion (Exons 1-5), was identified in HEXB.

### What is Sandhoff disease?

Sandhoff disease is a condition that affects lysosomes, which are structures in the cell that break down and recycle other molecules. Due to absent or reduced activity of the enzymes beta-hexosaminidase A and B (HEXA and HEXB), individuals with Sandhoff disease have difficulty breaking down a fatty substance called GM2 ganglioside and other substances. These substances accumulate in the cells, and are particularly toxic to the nerve cells in the central nervous system, leading to the destruction of neurons in the brain and spinal cord. The severity and age of onset of Sandhoff disease can vary, but the vast majority present in infancy with progressive weakness, loss of motor skills, and an increased startle reflex. Symptoms progress to include intellectual disability, hearing and vision loss, and seizures, with abnormal muscle tensing (spasticity). Affected individuals typically also have a characteristic cherry red spot at the back of the eye. Death usually occurs by age 3 or 4. Milder forms of the condition may be characterized by later onset, slower symptom progression, and more variable neurologic findings, including difficulty coordinating movements (ataxia) and psychiatric illness. Follow-up depends on each affected individual's specific situation, and discussion with a healthcare provider should be considered.

### Next steps


Carrier testing for the reproductive partner is recommended.


**If your partner tests positive:**

In autosomal recessive inheritance, an individual must have disease-causing genetic changes in each copy of the HEXB gene to be affected. Carriers, who have a disease-causing genetic change in only one copy of the gene, typically do not have symptoms. When both reproductive partners are carriers of an autosomal recessive condition, there is a 25% chance for each child to have the condition.


**If your partner tests negative:**

A negative carrier test result reduces, but does not eliminate, the chance that a person may be a carrier. The risk that a person could still be a carrier, even after a negative test result, is called a residual risk. See the table below for your partner's hypothetical residual risk after testing negative for Sandhoff disease. These values are provided only as a guide, are based on the detection rate for the condition as tested at Invitae, and assume a negative family history, the absence of symptoms, and vary based on the ethnic background of an individual. For genes associated with both dominant and recessive inheritance, the numbers provided apply to the recessive condition(s) associated with the gene.



| DISORDER (INHERITANCE)               | GENE | ETHNICITY  | CARRIER FREQUENCY BEFORE SCREENING | CARRIER RESIDUAL RISK AFTER NEGATIVE RESULT |
|--------------------------------------|------|------------|------------------------------------|---|
| Sandhoff disease (AR)<br>NM_000521.3 | HEXB | Pan-ethnic | 1 in 180                           | 1 in 17900                                  |

## Results to note

---

### SMN1

- Negative result. SMN1: 2 copies; c.\*3+80T>G not detected.

### Pseudodeficiency allele(s)

- Benign change, c.742G>A (p.Asp248Asn), known to be a pseudodeficiency allele, identified in the GALC gene. Pseudodeficiency alleles are not known to be associated with disease, including Krabbe disease.
- The presence of a pseudodeficiency allele does not impact this individual's risk to be a carrier. Individuals with pseudodeficiency alleles may exhibit false positive results on related biochemical tests, including newborn screening. However, pseudodeficiency alleles are not known to cause disease, even when there are two copies of the variant (homozygous) or when in combination with another disease-causing variant (compound heterozygous). Carrier testing for the reproductive partner is not indicated based on this result.

## Variant details

---

### ABCA4, Exon 17, c.2588G>C (p.Gly863Ala), heterozygous, Pathogenic (low penetrance)

- This sequence change replaces glycine, which is neutral and non-polar, with alanine, which is neutral and non-polar, at codon 863 of the ABCA4 protein (p.Gly863Ala).
- This variant is present in population databases (rs76157638, gnomAD 0.8%), including at least one homozygous and/or hemizygous individual.
- This variant has been reported in the compound-heterozygous state in several individuals and families affected with Stargardt disease and retinitis pigmentosa (PMID: 10612508, 10634594, 10090887, 12192456, 9054934, 23695285, 26247787, 25097241, 28041643). However, studies suggest that this is a mild variant that may only cause disease when in combination with a severe, pathogenic ABCA4 variant (PMID: 10090887).
- ClinVar contains an entry for this variant (Variation ID: 7879).
- An algorithm developed to predict the effect of missense changes on protein structure and function (PolyPhen-2) suggests that this variant is likely to be disruptive.
- Experimental studies have shown that this variant results in the production of two transcripts: one that lacks glycine 863 and the other with the Gly863Ala missense change (PMID: 10090887). Additional functional studies have shown that this missense change affects nucleotide hydrolysis and reduces the interaction of ABCA4 with 11-cis-retinal (PMID: 11919200, 23144455, 11017087).
- Algorithms developed to predict the effect of sequence changes on RNA splicing suggest that this variant may create or strengthen a splice site.
- In summary, this variant is reported to cause autosomal recessive Stargardt disease and retinitis pigmentosa. However, as this variant is associated with a lower penetrance than other pathogenic alleles in the ABCA4 gene, and as it may not result in disease in the homozygous state, it has been classified as Pathogenic (low penetrance).

### HEXB, Deletion (Exons 1-5), heterozygous, PATHOGENIC

- This variant is a gross deletion of the genomic region encompassing exon(s) 1-5 of the HEXB gene, which includes the initiator codon. This deletion extends beyond the assayed region for this gene and therefore may encompass additional genes. It is expected to result in an absent or disrupted protein product. Loss-of-function variants in HEXB are known to be pathogenic (PMID: 7550345, 18758829).
- A similar copy number variant has been observed in individuals with Sandhoff disease (PMID: 2921040, 23010210).
- For these reasons, this variant has been classified as Pathogenic.

### TMPRSS3, Exon 10, c.1028G>A (p.Trp343\*), heterozygous, PATHOGENIC

- This sequence change creates a premature translational stop signal (p.Trp343\*) in the TMPRSS3 gene. It is expected to result in an absent or disrupted protein product. Loss-of-function variants in TMPRSS3 are known to be pathogenic (PMID: 16021470, 26969326).
- This variant is not present in population databases (gnomAD no frequency).

- This variant has not been reported in the literature in individuals affected with TMPRSS3-related conditions.
- Algorithms developed to predict the effect of sequence changes on RNA splicing suggest that this variant may disrupt the consensus splice site.
- For these reasons, this variant has been classified as Pathogenic.

## Residual risk

---

No carrier test can detect 100% of carriers. There still remains a small risk of being a carrier after a negative test (residual risk). Residual risk values assume a negative family history and are inferred from published carrier frequencies and estimated detection rates based on testing technologies used at Invitae. You can view Invitae's complete Carrier detection rates and residual risks document (containing all carrier genes) online at <https://www.invitae.com/carrier-residual-risks/>. Additionally, the order-specific information for this report is available to download in the portal (under this order's documents) or can be requested by contacting Invitae Client Services. The complete Carrier detection rates and residual risks document will not be applicable for any genes with specimen-specific limitations in sequencing and/or deletion/duplication coverage. Please see the final bullet point in the Limitations section of this report to view if this specimen had any gene-specific coverage gaps.



## Genes analyzed

This table represents a complete list of genes analyzed for this individual, including the relevant gene transcript(s). If more than one transcript is listed for a single gene, variants were reported using the first transcript listed unless otherwise indicated in the report. An asterisk (\*) indicates that this gene has a limitation. Please see the Limitations section for details. Results are negative, unless otherwise indicated in the report.

| GENE     | TRANSCRIPT  | GENE     | TRANSCRIPT              | GENE     | TRANSCRIPT     |
|----------|-------------|----------|-------------------------|----------|----------------|
| AAAS     | NM_015665.5 | AP1S1    | NM_001283.3             | CBS      | NM_000071.2    |
| ABCA12   | NM_173076.2 | AQP2     | NM_000486.5             | CC2D1A   | NM_017721.5    |
| ABCA3    | NM_001089.2 | ARG1     | NM_000045.3             | CC2D2A   | NM_001080522.2 |
| ABCA4    | NM_000350.2 | ARL6     | NM_177976.2             | CCDC103  | NM_213607.2    |
| ABCB11   | NM_003742.2 | ARSA     | NM_000487.5             | CCDC39   | NM_181426.1    |
| ABCB4    | NM_000443.3 | ARSB     | NM_000046.3             | CCDC88C  | NM_001080414.3 |
| ABCC2*   | NM_000392.4 | ASL      | NM_000048.3             | CD3D     | NM_000732.4    |
| ABCC8    | NM_000352.4 | ASNS     | NM_133436.3             | CD3E     | NM_000733.3    |
| ACAD9    | NM_014049.4 | ASPA     | NM_000049.2             | CD40     | NM_001250.5    |
| ACADM    | NM_000016.5 | ASS1     | NM_000050.4             | CD59     | NM_203330.2    |
| ACADVL   | NM_000018.3 | ATM*     | NM_000051.3             | CDH23    | NM_022124.5    |
| ACAT1    | NM_000019.3 | ATP6V1B1 | NM_001692.3             | CEP152   | NM_014985.3    |
| ACOX1    | NM_004035.6 | ATP7B    | NM_000053.3             | CEP290   | NM_025114.3    |
| ACSF3    | NM_174917.4 | ATP8B1*  | NM_005603.4             | CERKL    | NM_001030311.2 |
| ADA      | NM_000022.2 | BBS1     | NM_024649.4             | CFTR*    | NM_000492.3    |
| ADAMTS2  | NM_014244.4 | BBS10    | NM_024685.3             | CHAT     | NM_020549.4    |
| ADAMTSL4 | NM_019032.5 | BBS12    | NM_152618.2             | CHRNE    | NM_000080.3    |
| ADGRG1   | NM_005682.6 | BBS2     | NM_031885.3             | CHRNA3   | NM_005199.4    |
| ADGRV1   | NM_032119.3 | BBS4     | NM_033028.4             | CIITA    | NM_000246.3    |
| AGA      | NM_000027.3 | BBS5     | NM_152384.2             | CLCN1    | NM_000083.2    |
| AGL      | NM_000642.2 | BBS7     | NM_176824.2             | CLN3     | NM_001042432.1 |
| AGPS     | NM_003659.3 | BBS9*    | NM_198428.2             | CLN5     | NM_006493.2    |
| AGXT     | NM_000030.2 | BCKDHA   | NM_000709.3             | CLN6     | NM_017882.2    |
| AHI1     | NM_017651.4 | BCKDHB   | NM_183050.2             | CLN8     | NM_018941.3    |
| AIPL1*   | NM_014336.4 | BCS1L    | NM_004328.4             | CLRN1    | NM_174878.2    |
| AIRE     | NM_000383.3 | BLM      | NM_000057.3             | CNGB3    | NM_019098.4    |
| ALDH3A2  | NM_000382.2 | BLOC1S3  | NM_212550.4             | COL11A2* | NM_080680.2    |
| ALDH7A1  | NM_001182.4 | BLOC1S6  | NM_012388.3             | COL17A1  | NM_000494.3    |
| ALDOB    | NM_000035.3 | BMP1     | NM_006129.4;NM_001199.3 | COL27A1  | NM_032888.3    |
| ALG1     | NM_019109.4 | BRIP1    | NM_032043.2             | COL4A3   | NM_000091.4    |
| ALG6     | NM_013339.3 | BSND     | NM_057176.2             | COL4A4   | NM_000092.4    |
| ALMS1    | NM_015120.4 | BTD      | NM_000060.3             | COL7A1   | NM_000094.3    |
| ALPL     | NM_000478.5 | CAD      | NM_004341.4             | COX15    | NM_004376.6    |
| AMN*     | NM_030943.3 | CANT1    | NM_138793.3             | CPS1     | NM_001875.4    |
| AMT      | NM_000481.3 | CAPN3    | NM_000070.2             | CPT1A    | NM_001876.3    |
| ANO10*   | NM_018075.3 | CASQ2    | NM_001232.3             | CPT2     | NM_000098.2    |



Patient name: Donor 7249

DOB: [REDACTED]

Invitae #: [REDACTED]

| GENE     | TRANSCRIPT     |
|----------|----------------|
| CRB1     | NM_201253.2    |
| CRTAP    | NM_006371.4    |
| CTNS     | NM_004937.2    |
| CTSA     | NM_000308.3    |
| CTSC     | NM_001814.5    |
| CTSD     | NM_001909.4    |
| CTSK     | NM_000396.3    |
| CYBA     | NM_000101.3    |
| CYP11A1  | NM_000781.2    |
| CYP11B1  | NM_000497.3    |
| CYP11B2  | NM_000498.3    |
| CYP17A1  | NM_000102.3    |
| CYP19A1  | NM_031226.2    |
| CYP1B1   | NM_000104.3    |
| CYP21A2* | NM_000500.7    |
| CYP27A1  | NM_000784.3    |
| CYP27B1  | NM_000785.3    |
| CYP7B1   | NM_004820.3    |
| DBT      | NM_001918.3    |
| DCAF17   | NM_025000.3    |
| DCLRE1C  | NM_001033855.2 |
| DDX11*   | NM_030653.3    |
| DFNB59   | NM_001042702.3 |
| DGAT1    | NM_012079.5    |
| DGUOK    | NM_080916.2    |
| DHCR7    | NM_001360.2    |
| DHDDS    | NM_024887.3    |
| DLD      | NM_000108.4    |
| DLL3     | NM_016941.3    |
| DNAH11   | NM_001277115.1 |
| DNAH5    | NM_001369.2    |
| DNAI1    | NM_012144.3    |
| DNAI2    | NM_023036.4    |
| DNMT3B   | NM_006892.3    |
| DOK7     | NM_173660.4    |
| DUOX2*   | NM_014080.4    |
| DYNC2H1  | NM_001080463.1 |
| DYSF     | NM_003494.3    |
| EIF2AK3  | NM_004836.6    |

| GENE    | TRANSCRIPT     |
|---------|----------------|
| EIF2B1  | NM_001414.3    |
| EIF2B2  | NM_014239.3    |
| EIF2B3  | NM_020365.4    |
| EIF2B4  | NM_015636.3    |
| EIF2B5  | NM_003907.2    |
| ELP1    | NM_003640.3    |
| EPG5    | NM_020964.2    |
| ERCC2   | NM_000400.3    |
| ERCC6   | NM_000124.3    |
| ERCC8   | NM_000082.3    |
| ESCO2   | NM_001017420.2 |
| ETFA    | NM_000126.3    |
| ETFB    | NM_001985.2    |
| ETFDH   | NM_004453.3    |
| ETHE1   | NM_014297.3    |
| EVC     | NM_153717.2    |
| EVC2    | NM_147127.4    |
| EXOSC3  | NM_016042.3    |
| EYS*    | NM_001142800.1 |
| FAH*    | NM_000137.2    |
| FAM161A | NM_001201543.1 |
| FANCA   | NM_000135.2    |
| FANCC   | NM_000136.2    |
| FANCD2* | NM_033084.3    |
| FANCE   | NM_021922.2    |
| FANCG   | NM_004629.1    |
| FANCI   | NM_001113378.1 |
| FANCL*  | NM_018062.3    |
| FBP1    | NM_000507.3    |
| FBXO7   | NM_012179.3    |
| FH*     | NM_000143.3    |
| FKBP10  | NM_021939.3    |
| FKRP    | NM_024301.4    |
| FKTN    | NM_001079802.1 |
| FMO3    | NM_006894.6    |
| FOXN1   | NM_003593.2    |
| FOXRED1 | NM_017547.3    |
| FRAS1   | NM_025074.6    |
| FREM2   | NM_207361.5    |

| GENE   | TRANSCRIPT     |
|--------|----------------|
| FUCA1  | NM_000147.4    |
| G6PC   | NM_000151.3    |
| G6PC3  | NM_138387.3    |
| GAA    | NM_000152.3    |
| GALC*  | NM_000153.3    |
| GALE*  | NM_000403.3    |
| GALK1  | NM_000154.1    |
| GALNS  | NM_000512.4    |
| GALNT3 | NM_004482.3    |
| GALT   | NM_000155.3    |
| GAMT   | NM_000156.5    |
| GATM   | NM_001482.2    |
| GBA*   | NM_001005741.2 |
| GBE1   | NM_000158.3    |
| GCDH   | NM_000159.3    |
| GCH1   | NM_000161.2    |
| GDF5   | NM_000557.4    |
| GFM1   | NM_024996.5    |
| GHR*   | NM_000163.4    |
| GJB2   | NM_004004.5    |
| GLB1   | NM_000404.2    |
| GLDC   | NM_000170.2    |
| GLE1   | NM_001003722.1 |
| GNB3*  | NM_001128227.2 |
| GNPAT  | NM_014236.3    |
| GNPTAB | NM_024312.4    |
| GNPTG  | NM_032520.4    |
| GNS    | NM_002076.3    |
| GORAB  | NM_152281.2    |
| GRHPR  | NM_012203.1    |
| GRIP1  | NM_021150.3    |
| GSS    | NM_000178.2    |
| GUCY2D | NM_000180.3    |
| GUSB   | NM_000181.3    |
| HADH   | NM_005327.4    |
| HADHA  | NM_000182.4    |
| HADHB  | NM_000183.2    |
| HAMP   | NM_021175.2    |
| HAX1   | NM_006118.3    |



Patient name: Donor 7249    DOB: ██████████

Invitae #: ██████████

| GENE    | TRANSCRIPT     |
|---------|----------------|
| HBA1*   | NM_000558.4    |
| HBA2    | NM_000517.4    |
| HBB     | NM_000518.4    |
| HEXA    | NM_000520.4    |
| HEXB    | NM_000521.3    |
| HGSNAT  | NM_152419.2    |
| HJV     | NM_213653.3    |
| HLCS    | NM_000411.6    |
| HMGCL   | NM_000191.2    |
| HMOX1   | NM_002133.2    |
| HOGA1   | NM_138413.3    |
| HPD     | NM_002150.2    |
| HPS1    | NM_000195.4    |
| HPS3    | NM_032383.4    |
| HPS4    | NM_022081.5    |
| HPS5    | NM_181507.1    |
| HPS6    | NM_024747.5    |
| HSD17B3 | NM_000197.1    |
| HSD17B4 | NM_000414.3    |
| HSD3B2  | NM_000198.3    |
| HYAL1   | NM_153281.1    |
| HYLS1   | NM_145014.2    |
| IDUA    | NM_000203.4    |
| IGHMBP2 | NM_002180.2    |
| IKBKB   | NM_001556.2    |
| IL7R    | NM_002185.3    |
| INVS    | NM_014425.3    |
| ITGA6   | NM_000210.3    |
| ITGB3   | NM_000212.2    |
| ITGB4   | NM_001005731.2 |
| IVD     | NM_002225.3    |
| JAK3    | NM_000215.3    |
| KCNJ1   | NM_000220.4    |
| KCNJ11  | NM_000525.3    |
| LAMA2   | NM_000426.3    |
| LAMA3   | NM_000227.4    |
| LAMB3   | NM_000228.2    |
| LAMC2   | NM_005562.2    |
| LARGE1  | NM_004737.4    |

| GENE    | TRANSCRIPT     |
|---------|----------------|
| LCA5    | NM_181714.3    |
| LDLR    | NM_000527.4    |
| LDLRAP1 | NM_015627.2    |
| LHX3    | NM_014564.4    |
| LIFR*   | NM_002310.5    |
| LIG4    | NM_002312.3    |
| LIPA    | NM_000235.3    |
| LMBRD1  | NM_018368.3    |
| LOXHD1  | NM_144612.6    |
| LPL     | NM_000237.2    |
| LRAT    | NM_004744.4    |
| LRP2    | NM_004525.2    |
| LRPPRC  | NM_133259.3    |
| LYST    | NM_000081.3    |
| MAK     | NM_001242957.2 |
| MAN2B1  | NM_000528.3    |
| MANBA   | NM_005908.3    |
| MCEE    | NM_032601.3    |
| MCOLN1  | NM_020533.2    |
| MCPH1   | NM_024596.4    |
| MECR    | NM_016011.3    |
| MED17   | NM_004268.4    |
| MESP2   | NM_001039958.1 |
| MFSD8   | NM_152778.2    |
| MKKS    | NM_018848.3    |
| MKS1    | NM_017777.3    |
| MLC1*   | NM_015166.3    |
| MLYCD   | NM_012213.2    |
| MMAA    | NM_172250.2    |
| MMAB    | NM_052845.3    |
| MMACHC  | NM_015506.2    |
| MMADHC  | NM_015702.2    |
| MOCS1   | NM_001358530.2 |
| MOCS2A  | NM_176806.3    |
| MOCS2B  | NM_004531.4    |
| MPI     | NM_002435.2    |
| MPL     | NM_005373.2    |
| MPV17   | NM_002437.4    |
| MRE11   | NM_005591.3    |

| GENE    | TRANSCRIPT              |
|---------|-------------------------|
| MTHFR*  | NM_005957.4             |
| MTR     | NM_000254.2             |
| MTRR    | NM_002454.2             |
| MTTP    | NM_000253.3             |
| MUSK    | NM_005592.3             |
| MUT     | NM_000255.3             |
| MVK     | NM_000431.3             |
| MYO15A  | NM_016239.3             |
| MYO7A   | NM_000260.3             |
| NAGA    | NM_000262.2             |
| NAGLU   | NM_000263.3             |
| NAGS    | NM_153006.2             |
| NBN     | NM_002485.4             |
| NCF2    | NM_000433.3             |
| NDRG1   | NM_006096.3             |
| NDUFAF2 | NM_174889.4             |
| NDUFAF5 | NM_024120.4             |
| NDUFS4  | NM_002495.3             |
| NDUFS6  | NM_004553.4             |
| NDUFS7  | NM_024407.4             |
| NDUFV1  | NM_007103.3             |
| NEB*    | NM_001271208.1          |
| NEU1    | NM_000434.3             |
| NGLY1   | NM_018297.3             |
| NPC1    | NM_000271.4             |
| NPC2    | NM_006432.3             |
| NPHP1   | NM_000272.3             |
| NPHS1   | NM_004646.3             |
| NPHS2   | NM_014625.3             |
| NR2E3   | NM_014249.3             |
| NSMCE3  | NM_138704.3             |
| NTRK1   | NM_001012331.1          |
| OAT*    | NM_000274.3             |
| OCA2    | NM_000275.2             |
| OPA3    | NM_025136.3             |
| OSTM1   | NM_014028.3             |
| OTOA*   | NM_144672.3             |
| OTOF    | NM_194248.2;NM_194323.2 |
| P3H1    | NM_022356.3             |



Patient name: Donor 7249    DOB: ██████████

Invitae #: ██████████

| GENE    | TRANSCRIPT                     |
|---------|--------------------------------|
| PAH     | NM_000277.1                    |
| PANK2   | NM_153638.2                    |
| PC      | NM_000920.3                    |
| PCBD1   | NM_000281.3                    |
| PCCA    | NM_000282.3                    |
| PCCB    | NM_000532.4                    |
| PCDH15  | NM_033056.3                    |
| PCNT    | NM_006031.5                    |
| PDHB    | NM_000925.3                    |
| PEPD    | NM_000285.3                    |
| PET100  | NM_001171155.1                 |
| PEX1*   | NM_000466.2                    |
| PEX10   | NM_153818.1                    |
| PEX12   | NM_000286.2                    |
| PEX13   | NM_002618.3                    |
| PEX16   | NM_004813.2                    |
| PEX2    | NM_000318.2                    |
| PEX26   | NM_017929.5                    |
| PEX5    | NM_001131025.1                 |
| PEX6    | NM_000287.3                    |
| PEX7    | NM_000288.3                    |
| PFKM    | NM_000289.5                    |
| PGM3    | NM_001199917.1                 |
| PHGDH   | NM_006623.3                    |
| PHKB    | NM_000293.2;NM_00103183<br>5.2 |
| PHKG2   | NM_000294.2                    |
| PHYH    | NM_006214.3                    |
| PIGN    | NM_176787.4                    |
| PKHD1*  | NM_138694.3                    |
| PLA2G6  | NM_003560.2                    |
| PLEKHG5 | NM_020631.4                    |
| PLOD1   | NM_000302.3                    |
| PMM2    | NM_000303.2                    |
| PNPO    | NM_018129.3                    |
| POLG    | NM_002693.2                    |
| POLH    | NM_006502.2                    |
| POMGNT1 | NM_017739.3                    |
| POMT1   | NM_007171.3                    |
| POMT2   | NM_013382.5                    |

| GENE     | TRANSCRIPT     |
|----------|----------------|
| POR      | NM_000941.2    |
| POU1F1   | NM_000306.3    |
| PPT1     | NM_000310.3    |
| PRCD     | NM_001077620.2 |
| PRDM5    | NM_018699.3    |
| PRF1     | NM_001083116.1 |
| PROP1    | NM_006261.4    |
| PSAP     | NM_002778.3    |
| PTPRC*   | NM_002838.4    |
| PTS      | NM_000317.2    |
| PUS1     | NM_025215.5    |
| PYGM     | NM_005609.3    |
| QDPR     | NM_000320.2    |
| RAB23    | NM_183227.2    |
| RAG1     | NM_000448.2    |
| RAG2     | NM_000536.3    |
| RAPSN    | NM_005055.4    |
| RARS2    | NM_020320.3    |
| RDH12    | NM_152443.2    |
| RLBP1    | NM_000326.4    |
| RMRP     | NR_003051.3    |
| RNASEH2A | NM_006397.2    |
| RNASEH2B | NM_024570.3    |
| RNASEH2C | NM_032193.3    |
| RPE65    | NM_000329.2    |
| RPGRIP1L | NM_015272.2    |
| RTEL1    | NM_001283009.1 |
| RXYLT1   | NM_014254.2    |
| RYR1     | NM_000540.2    |
| SACS     | NM_014363.5    |
| SAMD9    | NM_017654.3    |
| SAMHD1   | NM_015474.3    |
| SCO2     | NM_005138.2    |
| SEC23B   | NM_006363.4    |
| SEPSECS  | NM_016955.3    |
| SGCA     | NM_000023.2    |
| SGCB     | NM_000232.4    |
| SGCD     | NM_000337.5    |
| SGCG     | NM_000231.2    |

| GENE     | TRANSCRIPT     |
|----------|----------------|
| SGSH     | NM_000199.3    |
| SKIV2L   | NM_006929.4    |
| SLC12A1  | NM_000338.2    |
| SLC12A3  | NM_000339.2    |
| SLC12A6  | NM_133647.1    |
| SLC17A5  | NM_012434.4    |
| SLC19A2  | NM_006996.2    |
| SLC19A3  | NM_025243.3    |
| SLC1A4   | NM_003038.4    |
| SLC22A5  | NM_003060.3    |
| SLC25A13 | NM_014251.2    |
| SLC25A15 | NM_014252.3    |
| SLC25A20 | NM_000387.5    |
| SLC26A2  | NM_000112.3    |
| SLC26A3  | NM_000111.2    |
| SLC26A4  | NM_000441.1    |
| SLC27A4  | NM_005094.3    |
| SLC35A3  | NM_012243.2    |
| SLC37A4  | NM_001164277.1 |
| SLC38A8  | NM_001080442.2 |
| SLC39A4  | NM_130849.3    |
| SLC45A2  | NM_016180.4    |
| SLC4A11  | NM_032034.3    |
| SLC5A5   | NM_000453.2    |
| SLC7A7   | NM_001126106.2 |
| SMARCA11 | NM_014140.3    |
| SMN1*    | NM_000344.3    |
| SMPD1    | NM_000543.4    |
| SNAP29   | NM_004782.3    |
| SPG11    | NM_025137.3    |
| SPR      | NM_003124.4    |
| SRD5A2   | NM_000348.3    |
| ST3GAL5  | NM_003896.3    |
| STAR     | NM_000349.2    |
| STX11    | NM_003764.3    |
| STXBP2   | NM_006949.3    |
| SUMF1    | NM_182760.3    |
| SUOX     | NM_000456.2    |
| SURF1    | NM_003172.3    |



Patient name: Donor 7249

DOB: [REDACTED]

Invitae #: [REDACTED]

| GENE    | TRANSCRIPT     |
|---------|----------------|
| SYNE4   | NM_001039876.2 |
| TANGO2  | NM_152906.6    |
| TAT     | NM_000353.2    |
| TBCD    | NM_005993.4    |
| TBCE*   | NM_003193.4    |
| TCIRG1  | NM_006019.3    |
| TCN2    | NM_000355.3    |
| TECPR2  | NM_014844.3    |
| TERT    | NM_198253.2    |
| TF      | NM_001063.3    |
| TFR2    | NM_003227.3    |
| TG*     | NM_003235.4    |
| TGM1    | NM_000359.2    |
| TH      | NM_199292.2    |
| TK2     | NM_004614.4    |
| TMC1    | NM_138691.2    |
| TMEM216 | NM_001173990.2 |
| TMEM67  | NM_153704.5    |
| TMPRSS3 | NM_024022.2    |
| TPO     | NM_000547.5    |
| TPP1    | NM_000391.3    |
| TREX1   | NM_033629.4    |
| TRIM32  | NM_012210.3    |
| TRIM37  | NM_015294.4    |
| TRMU    | NM_018006.4    |
| TSEN54  | NM_207346.2    |
| TSFM*   | NM_001172696.1 |
| TSHB    | NM_000549.4    |
| TSHR    | NM_000369.2    |
| TTC37   | NM_014639.3    |
| TTPA    | NM_000370.3    |
| TULP1   | NM_003322.4    |
| TYMP    | NM_001953.4    |
| TYR*    | NM_000372.4    |
| TYRP1   | NM_000550.2    |
| UBR1    | NM_174916.2    |
| UNC13D  | NM_199242.2    |
| USH1C*  | NM_005709.3    |
| USH2A   | NM_206933.2    |

| GENE    | TRANSCRIPT     |
|---------|----------------|
| VDR     | NM_001017535.1 |
| VLDLR   | NM_003383.4    |
| VPS11   | NM_021729.5    |
| VPS13A* | NM_033305.2    |
| VPS13B  | NM_017890.4    |
| VPS45   | NM_007259.4    |
| VPS53*  | NM_001128159.2 |
| VRK1    | NM_003384.2    |
| VSX2    | NM_182894.2    |
| WISP3   | NM_003880.3    |
| WNT10A  | NM_025216.2    |
| WRN*    | NM_000553.4    |
| XPA     | NM_000380.3    |
| XPC     | NM_004628.4    |
| ZBTB24  | NM_014797.2    |
| ZFYVE26 | NM_015346.3    |
| ZNF469  | NM_001127464.2 |

## Methods

- Genomic DNA obtained from the submitted sample is enriched for targeted regions using a hybridization-based protocol, and sequenced using Illumina technology. Unless otherwise indicated, all targeted regions are sequenced with  $\geq 50\times$  depth or are supplemented with additional analysis. Reads are aligned to a reference sequence (GRCh37), and sequence changes are identified and interpreted in the context of a single clinically relevant transcript, indicated in the Genes Analyzed table. Enrichment and analysis focus on the coding sequence of the indicated transcripts, 20bp of flanking intronic sequence, and other specific genomic regions demonstrated to be causative of disease at the time of assay design. Promoters, untranslated regions, and other non-coding regions are not otherwise interrogated. Exonic deletions and duplications are called using an in-house algorithm that determines copy number at each target by comparing the read depth for each target in the proband sequence with both mean read-depth and read-depth distribution, obtained from a set of clinical samples. Markers across the X and Y chromosomes are analyzed for quality control purposes and may detect deviations from the expected sex chromosome complement. Such deviations may be included in the report in accordance with internal guidelines. Variants are reported according to the Human Genome Variation Society (HGVS) guidelines. Confirmation of the presence and location of reportable variants is performed as needed based on stringent criteria using one of several validated orthogonal approaches (PubMed ID 30610921). Sequencing is performed by Invitae Corporation (1400 16th Street, San Francisco, CA 94103, #05D2040778). Confirmatory sequencing is performed by Invitae Corporation (1400 16th Street, San Francisco, CA 94103, #05D2040778).

The following additional analyses are performed if relevant to the requisition. For GBA the reference genome has been modified to mask the sites of polymorphic paralog sequence variants (PSVs) in both the gene and pseudogene. For CYP21A2 and GBA, if one or more reportable variants, gene conversion, or fusion event is identified via our NGS pipeline (see Limitations), these variants are confirmed by PacBio sequencing of an amplicon generated by long-range PCR and subsequent short-range PCR. In some cases, it may not be possible to disambiguate between the gene and pseudogene. For GJB2, the reportable range includes large upstream deletions overlapping GJB6. For HBA1/2, the reference genome has been modified to force some sequencing reads derived from HBA1 to align to HBA2, and variant calling algorithms are modified to support an expectation of 4 alleles in these regions. HBA1/2 copy number calling is performed by a custom hypothesis testing algorithm which generates diplotype calls. If sequence data for a sample does not support a unique high confidence match from among hypotheses tested, that sample is flagged for manual review. Copy number variation is only reported for coding sequence of HBA1 and HBA2 and the HS-40 region. This assay does not distinguish among the  $\alpha 3.7$  subtypes, and all  $\alpha 3.7$  variants are called as HBA1 deletions. This assay may not detect overlapping copy gain and copy loss events when the breakpoints of those events are similar. For FMR1, cytosine-guanine-guanine (CGG) triplet repeats in the 5' untranslated region (5' UTR) of the FMR1 gene are detected by triplet repeat-primed PCR (RP-PCR) with fluorescently labeled primers followed by capillary electrophoresis. Reference ranges: Normal:  $<45$  CGG repeats, intermediate: 45-54 CGG repeats, premutation: 55-200 CGG repeats, full mutation:  $>200$  CGG repeats. For alleles with 55-90 triplet repeats, the region surrounding the FMR1 repeat is amplified by PCR. The PCR amplicons are then processed through PacBio SMRTBell library prep and sequenced using PacBio long read technology. The number of AGG interruptions within the 55-90 triplet repeat is read directly from the resulting DNA sequences.

- This report only includes variants that have a clinically significant association with the conditions tested as of the report date. Variants of uncertain significance, benign variants, and likely benign variants are not included in this report. However, if additional evidence becomes available to indicate that the clinical significance of a variant has changed, Invitae may update this report and provide notification.
- A PMID is a unique identifier referring to a published, scientific paper. Search by PMID at <http://www.ncbi.nlm.nih.gov/pubmed>.
- An rsID is a unique identifier referring to a single genomic position, and is used to associate population frequency information with sequence changes at that position. Reported population frequencies are derived from a number of public sites that aggregate data from large-scale population sequencing projects, including ExAC (<http://exac.broadinstitute.org>), gnomAD (<http://gnomad.broadinstitute.org>), and dbSNP (<http://ncbi.nlm.nih.gov/SNP>).

## Disclaimer

DNA studies do not constitute a definitive test for the selected condition(s) in all individuals. It should be realized that there are possible sources of error. Errors can result from trace contamination, rare technical errors, rare genetic variants that interfere with analysis, recent scientific developments, and alternative classification systems. This test should be one of many aspects used by the healthcare provider to help with a diagnosis and treatment plan, but it is not a diagnosis itself. This test was developed and its performance characteristics determined by Invitae. It has not been cleared or approved by

the FDA. The laboratory is regulated under the Clinical Laboratory Improvement Act (CLIA) as qualified to perform high-complexity clinical tests (CLIA ID: 05D2040778). This test is used for clinical purposes. It should not be regarded as investigational or for research.

## Limitations

- Based on validation study results, this assay achieves >99% analytical sensitivity and specificity for single nucleotide variants, insertions and deletions <15bp in length, and exon-level deletions and duplications. Invitae's methods also detect insertions and deletions larger than 15bp but smaller than a full exon but sensitivity for these may be marginally reduced. Invitae's deletion/duplication analysis determines copy number at a single exon resolution at virtually all targeted exons. However, in rare situations, single-exon copy number events may not be analyzed due to inherent sequence properties or isolated reduction in data quality. Certain types of variants, such as structural rearrangements (e.g. inversions, gene conversion events, translocations, etc.) or variants embedded in sequence with complex architecture (e.g. short tandem repeats or segmental duplications), may not be detected. Additionally, it may not be possible to fully resolve certain details about variants, such as mosaicism, phasing, or mapping ambiguity. Unless explicitly guaranteed, sequence changes in the promoter, non-coding exons, and other non-coding regions are not covered by this assay. Please consult the test definition on our website for details regarding regions or types of variants that are covered or excluded for this test. This report reflects the analysis of an extracted genomic DNA sample. While this test is intended to reflect the analysis of extracted genomic DNA from a referred patient, in very rare cases the analyzed DNA may not represent that individual's constitutional genome, such as in the case of a circulating hematolymphoid neoplasm, bone marrow transplant, blood transfusion, chimerism, culture artifact or maternal cell contamination. Interpretations are made on the assumption that any clinical information provided, including specimen identity, is accurate.
- ANO10: Sequencing analysis for exons 8 includes only cds +/- 0 bp. ATP8B1: Sequencing analysis for exons 19 includes only cds +/- 10 bp. AIPL1: Sequencing analysis for exons 2 includes only cds +/- 10 bp. GHR: Deletion/duplication and sequencing analysis is not offered for exon 3. TBCE: Sequencing analysis for exons 2 includes only cds +/- 10 bp. CYP21A2: Analysis includes the most common variants (c.92C>T(p.Pro31Leu), c.293-13C>G (intronic), c.332\_339delGAGACTAC (p.Gly111Valfs\*21), c.518T>A (p.Ile173Asn), c.710T>A (p.Ile237Asn), c.713T>A (p.Val238Glu), c.719T>A (p.Met240Lys), c.844G>T (p.Val282Leu), c.923dupT (p.Leu308Phefs\*6), c.955C>T (p.Gln319\*), c.1069C>T(p.Arg357Trp), c.1360C>T (p.Pro454Ser) and the 30Kb deletion) as well as select rare HGMD variants only (list available upon request). Full gene duplications are reported only in the presence of a pathogenic variant(s). When a duplication and a pathogenic variant(s) is identified, phase (cis/trans) cannot be determined. Full gene deletion analysis is not offered. Sensitivity to detect these variants, if they result from complex gene conversion/fusion events, may be reduced. TYR: Deletion/duplication and sequencing analysis is not offered for exon 5. PTPRC: Sequencing analysis is not offered for exons 3, 15. ABCC2: Deletion/duplication analysis is not offered for exons 24-25. OTOA: Deletion/duplication and sequencing analysis is not offered for exons 20-28. DUOX2: Deletion/duplication and sequencing analysis is not offered for exons 6-7. TG: Deletion/duplication analysis is not offered for exon 18. Sequencing analysis for exons 44 includes only cds +/- 0 bp. FANCD2: Deletion/duplication analysis is not offered for exons 14-17, 22 and sequencing analysis is not offered for exons 15-17. Sequencing analysis for exons 6, 14, 18, 20, 23, 25, 34 includes only cds +/- 10 bp. FANCL: Sequencing analysis for exons 4, 10 includes only cds +/- 10 bp. ATM: Sequencing analysis for exons 6, 24, 43 includes only cds +/- 10 bp. CFTR: Sequencing analysis for exons 7 includes only cds +/- 10 bp. EYS: Sequencing analysis for exons 30 includes only cds +/- 0 bp. FAH: Deletion/duplication analysis is not offered for exon 14. FH: Sequencing analysis for exons 9 includes only cds +/- 10 bp. GALC: Deletion/duplication analysis is not offered for exon 6. GBA: c.84dupG (p.Leu29Alafs\*18), c.115+1G>A (Splice donor), c.222\_224delTAC (p.Thr75del), c.475C>T (p.Arg159Trp), c.595\_596delCT (p.Leu199Aspfs\*62), c.680A>G (p.Asn227Ser), c.721G>A (p.Gly241Arg), c.754T>A (p.Phe252Ile), c.1226A>G (p.Asn409Ser), c.1246G>A (p.Gly416Ser), c.1263\_1317del (p.Leu422Profs\*4), c.1297G>T (p.Val433Leu), c.1342G>C (p.Asp448His), c.1343A>T (p.Asp448Val), c.1448T>C (p.Leu483Pro), c.1504C>T (p.Arg502Cys), c.1505G>A (p.Arg502His), c.1603C>T (p.Arg535Cys), c.1604G>A (p.Arg535His) variants only. Rarely, sensitivity to detect these variants may be reduced. When sensitivity is reduced, zygosity may be reported as "unknown". GNE: Sequencing analysis for exons 8 includes only cds +/- 10 bp. HBA1/2: This assay is designed to detect deletions and duplications of HBA1 and/or HBA2, resulting from the -alpha20.5, --MED, --SEA, --FIL/--THAI, -alpha3.7, -alpha4.2, anti3.7 and anti4.2. Sensitivity to detect other copy number variants may be reduced. Detection of overlapping deletion and duplication events will be limited to combinations of events with significantly differing boundaries. In addition, deletion of the enhancer element HS-40 and the sequence variant, Constant Spring (NM\_000517.4:c.427T>C), can be identified by this assay. LIFR: Sequencing analysis for exons 3 includes only cds +/- 5 bp. MLC1: Sequencing analysis for exons 11 includes only cds +/- 10 bp. MTHFR: The NM\_005957.4:c.665C>T (p.Ala222Val) (aka 677C>T) and c.1286A>C (p.Glu429Ala) (aka 1298A>C) variants are not reported in our primary report. NEB: Deletion/duplication analysis is not offered for exons 82-105. NEB variants in this region with no evidence towards pathogenicity are not included in this report, but are available upon request. OAT: Deletion/duplication analysis is not offered for exon 2. PEX1: Sequencing analysis for exons 16 includes only cds +/- 0 bp. PKHD1: Deletion/duplication analysis is not offered for exon 13. SMN1: Systematic exon numbering is used for all genes, including SMN1, and for this reason the exon typically referred to as exon 7 in the literature (PMID: 8838816) is referred to as exon 8 in this report. This assay unambiguously detects SMN1 exon 8 copy number. The



Patient name: Donor 7249

DOB: [REDACTED]

Invitae #: [REDACTED]

presence of the g.27134T>G variant (also known as c.\*3+80T>G) is reported if SMN1 copy number = 2. SMN1 or SMN2: NM\_000344.3:c.\*3+80T>G variant only. TSM: Sequencing analysis is not offered for exon 5. USH1C: Deletion/duplication analysis is not offered for exons 5-6. VPS13A: Deletion/duplication analysis is not offered for exons 2-3, 27-28. VPS53: Sequencing analysis for exons 14 includes only cds +/- 5 bp. AMN: Deletion/duplication analysis is not offered for exon 1. GALE: Sequencing analysis for exons 10 includes only cds +/- 5 bp. DDX11: NM\_030653.3:c.1763-1G>C variant only. BBS9: Deletion/duplication analysis is not offered for exon 4. COL11A2: Deletion/duplication analysis is not offered for exon 36. WRN: Deletion/duplication analysis is not offered for exons 10-11. Sequencing analysis for exons 8, 10-11 includes only cds +/- 10 bp.

### This report has been reviewed and approved by:



Matteo Vatta, Ph.D., FACMG  
Clinical Molecular Geneticist



PATIENT INFORMATION

7249, DONOR

REPORT STATUS **Final**

Nichols Institute, Chantilly

DOB: [REDACTED] Age: [REDACTED]

SEX: M

ORDERING PHYSICIAN

CLIENT INFORMATION

ID: 7249-[REDACTED]

SPECIMEN INFORMATION

SPECIMEN: [REDACTED]

REQUISITION: [REDACTED]

LAB REF NO: 7249-[REDACTED]

COLLECTED: 10/20/2023 00:00  
 RECEIVED: 10/21/2023 12:33  
 REPORTED: 10/30/2023 21:57

| Test Name                   | In Range | Out of Range | Reference Range   | Lab |
|-----------------------------|----------|--------------|-------------------|-----|
| Hemoglobinopathy Evaluation |          |              |                   | AMD |
| Red Blood Cell Count        | 5.10     |              | 4.20-5.80 Mill/uL |     |
| HEMOGLOBIN                  | 15.2     |              | 13.2-17.1 g/dL    |     |
| Hematocrit                  |          |              |                   |     |
| Hematocrit                  | 46.9     |              | 38.5-50.0 %       |     |
| MCV                         | 92.0     |              | 80.0-100.0 fL     |     |
| MCH                         | 29.8     |              | 27.0-33.0 pg      |     |
| RDW                         | 12.6     |              | 11.0-15.0 %       |     |
| Hemoglobin A                | 97.6     |              | >96.0 %           |     |
| Hemoglobin F                | 0.0      |              | <2.0 %            |     |
| Hemoglobin A2 (Quant)       | 2.4      |              | 2.2-3.2 %         |     |
| Interpretation              |          |              |                   |     |

NORMAL PATTERN

There is a normal pattern of hemoglobins and normal levels of Hb A2 and Hb F are present. No variant hemoglobins are observed. This is consistent with A/A phenotype. If iron deficiency coexists with a mild/silent beta thalassemia trait Hb A2 may be in the normal range. Rare variant hemoglobins have no separation from hemoglobin A by capillary zone electrophoresis (CZE) or high-performance liquid chromatography (HPLC). If clinically indicated, Thalassemia and Hemoglobinopathy Comprehensive (TC 17365) should be considered.

|   |      |  |                   |     |
|---|------|--|-------------------|-----|
| CBC (includes Differential and Platelets) |      |  |                   | AMD |
| CBC (includes Differential and Platelets) |      |  |                   |     |
| White Blood Cell Count                    | 4.9  |  | 3.8-10.8 Thous/uL |     |
| Red Blood Cell Count                      | 5.10 |  | 4.20-5.80 Mill/uL |     |
| HEMOGLOBIN                                | 15.2 |  | 13.2-17.1 g/dL    |     |
| Hematocrit                                | 46.9 |  | 38.5-50.0 %       |     |
| MCV                                       | 92.0 |  | 80.0-100.0 fL     |     |
| MCH                                       | 29.8 |  | 27.0-33.0 pg      |     |
| MCHC                                      | 32.4 |  | 32.0-36.0 g/dL    |     |
| RDW                                       | 12.6 |  | 11.0-15.0 %       |     |
| PLATELET COUNT                            | 267  |  | 140-400 Thous/uL  |     |
| MPV                                       | 10.6 |  | 7.5-12.5 fl       |     |

PATIENT INFORMATION  
7249, DONOR

REPORT STATUS **Final**

Nichols Institute, Chantilly

ORDERING PHYSICIAN

DOB: [REDACTED] Age: [REDACTED]  
SEX: M  
ID: 7249-[REDACTED]

COLLECTED: 10/20/2023 00:00  
REPORTED: 10/30/2023 21:57

| Test Name   | In Range   | Out of Range | Reference Range    | Lab        |
|---|------------|--------------|--------------------|------------|
| CBC (includes Differential and Platelets) (Continued) |            |              |                    |            |
| Absolute Neutrophils                                  | 1882       |              | 1500-7800 cells/uL |            |
| Absolute Lymphocytes                                  | 2465       |              | 850-3900 cells/uL  |            |
| Absolute Monocytes                                    | 323        |              | 200-950 cells/uL   |            |
| Absolute Eosinophils                                  | 181        |              | 15-500 cells/uL    |            |
| Absolute Basophils                                    | 49         |              | 0-200 cells/uL     |            |
| Neutrophils   | 38.4       |              | %                  |            |
| Lymphocytes   | 50.3       |              | %                  |            |
| Monocytes   | 6.60       |              | %                  |            |
| Eosinophils   | 3.7        |              | %                  |            |
| Basophils   | 1.00       |              | %                  |            |
| Nucleated RBC   | 0.00       |              | 0 /100 WBC         |            |
| [REDACTED]  |            | [REDACTED]   | [REDACTED]         | [REDACTED] |
| [REDACTED]  | [REDACTED] |              | [REDACTED]         | [REDACTED] |
| [REDACTED]  | [REDACTED] |              | [REDACTED]         | [REDACTED] |

Chromosome Analysis, Blood AMD  
Chromosome Analysis, Blood  
Chromosome Analysis, Blood

Order ID: [REDACTED]

Specimen Type: Blood

Clinical Indication: Gamete donor, rule out chromosome abnormality

RESULT:  
NORMAL MALE KARYOTYPE

INTERPRETATION:  
Chromosome analysis revealed normal G-band patterns within the limits of standard cytogenetic analysis.

Please expect the results of any other concurrent study in a separate report.

NOMENCLATURE:  
46,XY

ASSAY INFORMATION:  
Method: G-Band (Digital Analysis:  
MetaSystems/Ikaros)  
Cells Counted: 20  
Band Level: 550  
Cells Analyzed: 5  
Cells Karyotyped: 5

This test does not address genetic disorders that cannot be detected by standard cytogenetic methods or rare events such as low level mosaicism or subtle rearrangements.

PATIENT INFORMATION

7249, DONOR

REPORT STATUS **Final**

Nichols Institute, Chantilly

ORDERING PHYSICIAN

DOB: [REDACTED]

Age: [REDACTED]

SEX: M

ID: 7249-[REDACTED]

COLLECTED: 10/20/2023 00:00

REPORTED: 10/30/2023 21:57

| Test Name | In Range | Out of Range | Reference Range | Lab |
|-----------|----------|--------------|-----------------|-----|
|-----------|----------|--------------|-----------------|-----|

Chromosome Analysis, Blood (Continued)

Chromosome Analysis, Blood (Continued)


Debra Boles, Ph.D., FACMG, Technical Director, Cytogenetics and Genomics, 703-802-7156

Electronic Signature: 10/30/2023 9:12 PM

For additional information, please refer to <http://education.questdiagnostics.com/faq/chromsblood> (This link is being provided for informational/educational purposes only).

-----  
**Performing Laboratory Information:**

AMD Quest Diagnostics Nichols Institute 14225 Newbrook Drive Chantilly VA 20151 Laboratory Director: Patrick W Mason, MD PhD



Patient Information:

7249, Donor

DOB: [REDACTED]

Sex: M

MR#: 7249

Patient#: [REDACTED]

Partner Information:

Not Tested

Physician:

Seitz, Suzanne

ATTN: Seitz, Suzanne

Fairfax Cryobank

3015 Williams Drive

Fairfax, VA 22031

Laboratory:

Fulgent Therapeutics LLC

CAP#: 8042697

CLIA#: 05D2043189

Laboratory Director:

Lawrence M. Weiss, MD

Report Date: Jul 28, 2024

Accession:

[REDACTED]

Test#: [REDACTED]

Specimen Type: DNA

Collected: Not Provided

Accession:

N/A

## FINAL RESULTS



No carrier mutations identified

## TEST PERFORMED

### Single Gene Carrier Screening: SLC6A19

(1 Gene Panel: *SLC6A19*; gene sequencing with deletion and duplication analysis)

## INTERPRETATION:

### Notes and Recommendations:

- No carrier mutations were identified in the submitted specimen. A negative result does not rule out the possibility of a genetic predisposition nor does it rule out any pathogenic mutations in areas not assessed by this test or in regions that were covered at a level too low to reliably assess. Also, it does not rule out mutations that are of the sort not queried by this test; see Methods and Limitations for more information. A negative result reduces, but does not eliminate, the chance to be a carrier for any condition included in this screen. Please see the supplemental table for details.
- This carrier screening test does not screen for all possible genetic conditions, nor for all possible mutations in every gene tested. This report does not include variants of uncertain significance; only variants classified as pathogenic or likely pathogenic at the time of testing, and considered relevant for reproductive carrier screening, are reported. Please see the gene specific notes for details. Please note that the classification of variants can change over time.
- Patients may wish to discuss any carrier results with blood relatives, as there is an increased chance that they are also carriers. These results should be interpreted in the context of this individual's clinical findings, biochemical profile, and family history.
- Gene specific notes and limitations may be present. See below.
- Genetic counseling is recommended. Available genetic counselors and additional resources can be found at the National Society of Genetic Counselors (NSGC; <https://www.nsgc.org>)



## GENES TESTED:

---

### Custom Beacon Carrier Screening Panel - Gene

This analysis was run using the Custom Beacon Carrier Screening Panel gene list. 1 genes were tested with 100.0% of targets sequenced at >20x coverage. For more gene-specific information and assistance with residual risk calculation, see the SUPPLEMENTAL TABLE.

SLC6A19

## METHODS:

---

Genomic DNA was isolated from the submitted specimen indicated above (if cellular material was submitted). DNA was barcoded, and enriched for the coding exons of targeted genes using hybrid capture technology. Prepared DNA libraries were then sequenced using a Next Generation Sequencing technology. Following alignment to the human genome reference sequence (assembly GRCh37), variants were detected in regions of at least 10x coverage. For this specimen, 100.00% and 100.00% of coding regions and splicing junctions of genes listed had been sequenced with coverage of at least 10x and 20x, respectively, by NGS or by Sanger sequencing. The remaining regions did not have 10x coverage, and were not evaluated. Variants were interpreted manually using locus specific databases, literature searches, and other molecular biological principles. To minimize false positive results, any variants that do not meet internal quality standards are confirmed by Sanger sequencing. Variants classified as pathogenic, likely pathogenic, or risk allele which are located in the coding regions and nearby intronic regions (+/- 20bp) of the genes listed above are reported. Variants outside these intervals may be reported but are typically not guaranteed. When a single pathogenic or likely pathogenic variant is identified in a clinically relevant gene with autosomal recessive inheritance, the laboratory will attempt to ensure 100% coverage of coding sequences either through NGS or Sanger sequencing technologies ("fill-in"). All genes listed were evaluated for large deletions and/or duplications. However, single exon deletions or duplications will not be detected in this assay, nor will copy number alterations in regions of genes with significant pseudogenes. Putative deletions or duplications are analyzed using Fulgent Germline proprietary pipeline for this specimen. Bioinformatics: The Fulgent Germline v2019.2 pipeline was used to analyze this specimen.

## LIMITATIONS:

---

### General Limitations

These test results and variant interpretation are based on the proper identification of the submitted specimen, accuracy of any stated familial relationships, and use of the correct human reference sequences at the queried loci. In very rare instances, errors may result due to mix-up or co-mingling of specimens. Positive results do not imply that there are no other contributors, genetic or otherwise, to future pregnancies, and negative results do not rule out the genetic risk to a pregnancy. Official gene names change over time. Fulgent uses the most up to date gene names based on HUGO Gene Nomenclature Committee (<https://www.genenames.org>) recommendations. If the gene name on report does not match that of ordered gene, please contact the laboratory and details can be provided. Result interpretation is based on the available clinical and family history information for this individual, collected published information, and Alamut annotation available at the time of reporting. This assay is not designed or validated for the detection of low-level mosaicism or somatic mutations. This assay will not detect certain types of genomic aberrations such as translocations, inversions, or repeat expansions other than specified genes. DNA alterations in regulatory regions or deep intronic regions (greater than 20bp from an exon) may not be detected by this test. Unless otherwise indicated, no additional assays have been performed to evaluate genetic changes in this specimen. There are technical limitations on the ability of DNA sequencing to detect small insertions and deletions. Our laboratory uses a sensitive detection algorithm, however these types of alterations are not detected as reliably as single nucleotide variants. Rarely, due to systematic chemical, computational, or human error, DNA variants may be missed. Although next generation sequencing technologies and our bioinformatics analysis significantly reduce the confounding contribution of pseudogene sequences or other highly-homologous sequences, sometimes these may still interfere with the technical ability of the assay to identify pathogenic alterations in both sequencing and deletion/duplication analyses. Deletion/duplication analysis can identify alterations of genomic regions which include one whole gene (buccal swab specimens and whole blood specimens) and are two or more contiguous exons in size (whole blood specimens only); single exon deletions or duplications may occasionally be identified, but are not routinely detected by this test. When novel DNA duplications are identified, it is not possible to discern the genomic location or orientation of the duplicated segment, hence the effect of the duplication cannot be predicted. Where deletions are detected, it is not always possible to determine whether the predicted product will remain in-frame or not. Unless otherwise indicated, deletion/duplication analysis has not been performed in regions that have been sequenced by Sanger.



### Gene Specific Notes and Limitations

No gene specific limitations apply to the genes on the tested panel.

### SIGNATURE:

---



A handwritten signature in black ink that reads "Harry Gao".

**Dr. Harry Gao, DABMG, FACMG** on 7/28/2024  
Laboratory Director, Fulgent

### DISCLAIMER:

---

This test was developed and its performance characteristics determined by **Fulgent Therapeutics LLC**. It has not been cleared or approved by the FDA. The laboratory is regulated under CLIA as qualified to perform high-complexity testing. This test is used for clinical purposes. It should not be regarded as investigational or for research. Since genetic variation, as well as systematic and technical factors, can affect the accuracy of testing, the results of testing should always be interpreted in the context of clinical and familial data. For assistance with interpretation of these results, healthcare professionals may contact us directly at **(626) 350-0537** or [info@fulgentgenetics.com](mailto:info@fulgentgenetics.com). It is recommended that patients receive appropriate genetic counseling to explain the implications of the test result, including its residual risks, uncertainties and reproductive or medical options.

4399 Santa Anita Ave.  
El Monte, CA, 91731  
(p) 626-350-0537 (f) 626-454-1667  
[info@fulgentgenetics.com](mailto:info@fulgentgenetics.com)  
[www.fulgentgenetics.com](http://www.fulgentgenetics.com)



To view the supplemental table describing the carrier frequencies, detection rates, and residual risks associated with the genes on this test please visit the following link:

[Beacon Expanded Carrier Screening Supplemental Table](#)

