



Donor 7164

Genetic Testing Summary

Fairfax Cryobank recommends reviewing this genetic testing summary with your healthcare provider to determine suitability.

Last Updated: 03/13/25

Donor Reported Ancestry: Cuban, Japanese

Jewish Ancestry: No

Genetic Test*	Result	Comments/Donor's Residual Risk**
Chromosome analysis (karyotype)	Normal male karyotype	No evidence of clinically significant chromosome abnormalities
Hemoglobin evaluation	Normal hemoglobin fractionation and MCV/MCH results	Reduced risk to be a carrier for sickle cell anemia, beta thalassemia, alpha thalassemia trait (aa/-- and a-/a-) and other hemoglobinopathies
Expanded Genetic Disease Carrier Screening Panel attached- 549 diseases by gene sequencing.	Carrier: Aicardi-Goutieres Syndrome, RNASEH2A-Related Carrier: Primary Congenital Glaucoma/ Peters Anomaly (CYP1B1) Carrier: Retinitis Pigmentosa 25 (EYS) Carrier: Xeroderma Pigmentosum, Group A (XPA) Negative for other genes sequenced.	Partner testing is recommended before using this donor.

*No single test can screen for all genetic disorders. A negative screening result significantly reduces, but cannot eliminate, the risk for these conditions in a pregnancy.

**Donor residual risk is the chance the donor is still a carrier after testing negative.

Patient Information

Patient Name: Donor 7164

Date Of Birth: [REDACTED]

Gender: Male

Ethnicity: Other

Patient ID: N/A

Medical Record #: N/A

Collection Kit: [REDACTED]

Accession ID: N/A

Case File ID: [REDACTED]

Test Information

Ordering Physician: [REDACTED]

Clinic Information: Fairfax Cryobank

Phone: [REDACTED]

Report Date: 12/17/2024

Sample Collected: 12/03/2024

Sample Received: 12/04/2024

Sample Type: Blood

**CARRIER SCREENING REPORT**

ABOUT THIS SCREEN: Horizon™ is a carrier screen for specific autosomal recessive and X-linked diseases. This information can help patients learn their risk of having a child with specific genetic conditions.

ORDER SELECTED: The Horizon Custom panel was ordered for this patient. Males are not screened for X-linked diseases

FINAL RESULTS SUMMARY:**CARRIER for Aicardi-Goutieres Syndrome, RNASEH2A-Related**

Positive for the likely pathogenic variant c.199+1G>C in the RNASEH2A gene. If this individual's partner is a carrier for AICARDI-GOUTIERES SYNDROME, RNASEH2A-RELATED, their chance to have a child with this condition may be as high as 1 in 4 (25%). Carrier screening for this individual's partner is suggested.

CARRIER for Primary Congenital Glaucoma/Peters Anomaly

Positive for the pathogenic variant c.535del (p.A179Rfs*18) in the CYP1B1 gene. If this individual's partner is a carrier for PRIMARY CONGENITAL GLAUCOMA/PETERS ANOMALY, their chance to have a child with this condition is 1 in 4 (25%). Carrier screening for this individual's partner is suggested.

CARRIER for Retinitis Pigmentosa 25

Positive for the pathogenic variant c.2380C>T (p.R794*) in the EYS gene. If this individual's partner is a carrier for RETINITIS PIGMENTOSA 25, their chance to have a child with this condition is 1 in 4 (25%). Carrier screening for this individual's partner is suggested.

CARRIER for Xeroderma Pigmentosum, Group A

Positive for the pathogenic variant c.390-1G>C in the XPA gene. If this individual's partner is a carrier for XERODERMA PIGMENTOSUM, GROUP A, their chance to have a child with this condition is 1 in 4 (25%). Carrier screening for this individual's partner is suggested.

Negative for 545 out of 549 diseases

No other pathogenic variants were detected in the genes that were screened. The patient's remaining carrier risk after the negative screening results is listed for each disease/gene on the Horizon website at <https://www.natera.com/panel-option/h-all/>. Please see the following pages of this report for a comprehensive list of all conditions included on this individual's screen.

Carrier screening is not diagnostic and may not detect all possible pathogenic variants in a given gene.

RECOMMENDATIONS

Individuals who would like to review their Horizon report with a Natera Laboratory Genetic Counselor may schedule a telephone genetic information session by calling 650-249-9090 or visiting naterasession.com. Clinicians with questions may contact Natera at 650-249-9090 or email support@natera.com. Individuals with positive results may wish to discuss these results with family members to allow them the option to be screened. Comprehensive genetic counseling to discuss the implications of these test results and possible associated reproductive risk is recommended.

Christine M. Eng, M.D.
Medical Director, Baylor Genetics

J. Dianne Keen-Kim, Ph.D., FACMG
Senior Laboratory Director, Natera

Xiaoyan Ge, Ph.D., FACMG
Associate Laboratory Director, Baylor Genetics

Yang Wang, Ph.D., FACMG
Laboratory Director, Natera

Patient Information

Patient Name: Donor 7164

Test Information

Ordering Physician: [REDACTED]



Clinic Information: Fairfax Cryobank

Date Of Birth: [REDACTED]

Case File ID: [REDACTED]

Report Date: 12/17/2024

AICARDI-GOUTIERES SYNDROME, RNASEH2A-RELATED**Understanding Your Horizon Carrier Screen Results****What is Aicardi-Goutieres syndrome, RNASEH2A-Related?**

Aicardi-Goutieres Syndrome, RNASEH2A-Related is an inherited disorder that affects the brain, immune system, and skin. Symptoms often first appear in infancy and include severe irritability, poor feeding, vomiting, fever, and seizures. Episodes of brain inflammation (encephalopathy) cause slowed brain growth and small head size (microcephaly), delayed development, loss of skills, and inability to walk. Most children with Aicardi-Goutieres Syndrome, RNASEH2A-Related have severe intellectual disability, muscle stiffness (spasticity), involuntary muscle spasms called dystonia, and abnormal eye movements. They may also have painful or itchy skin patches called chilblains on their fingers, toes and ears. Children with this condition often do not survive past childhood, although people with milder symptoms may live into adulthood. Currently there is no cure or specific treatment for this condition. Clinical trials involving potential new treatments for this condition may be available (see www.clinicaltrials.gov).

What causes Aicardi-Goutieres syndrome, RNASEH2A-Related?

Aicardi-Goutieres Syndrome, RNASEH2A-Related is caused by gene changes, or mutations, in both copies of the RNASEH2A gene pair. These mutations cause the gene to not work properly or not work at all. When both copies of the RNASEH2A gene do not work correctly, an abnormal immune and inflammatory response in the brain and skin occurs and leads to the symptoms described above. Aicardi-Goutieres Syndrome, RNASEH2A-Related is inherited in an autosomal recessive manner. This means that, in most cases, both parents must be carriers of a mutation in one copy of the RNASEH2A gene to have a child with Aicardi-Goutieres Syndrome, RNASEH2A-Related. People who are carriers for Aicardi-Goutieres Syndrome, RNASEH2A-Related are usually healthy and do not have the condition themselves. Usually a child inherits two copies of each gene, one copy from the mother and one copy from the father. If the mother and father are both carriers for Aicardi-Goutieres Syndrome, RNASEH2A-Related there is a 1 in 4, or 25%, chance in each pregnancy for both partners to pass on their RNASEH2A gene mutations to the child, who will then have this condition. Individuals found to carry more than one mutation for Aicardi-Goutieres Syndrome, RNASEH2A-Related should discuss their risk for having an affected child, and any specific risks to their own health, with their health care provider. There are a number of other forms of Aicardi-Goutieres Syndrome, each caused by mutations in different genes. People who are carriers for a mutation in the RNASEH2A gene are not likely to be at increased risk for having children with these other forms of the condition.

What can I do next?

You may wish to speak with a local genetic counselor about your carrier test results. A genetic counselor in your area can be located on the National Society of Genetic Counselors website (www.nsgc.org). Your siblings and other relatives are at increased risk to also have this mutation. You are encouraged to inform your family members of your test results as they may wish to consider being tested themselves. If you are pregnant, your partner can have carrier screening for Aicardi-Goutieres Syndrome, RNASEH2A-Related ordered by a health care professional. If your partner is not found to be a carrier of Aicardi-Goutieres Syndrome, RNASEH2A-Related, your risk of having a child with this condition is greatly reduced. Couples at risk of having a child with Aicardi-Goutieres Syndrome, RNASEH2A-Related can opt to have prenatal diagnostic testing done through chorionic villus sampling (CVS) or amniocentesis during pregnancy or can choose to have the baby tested after birth for this condition. If you are not yet pregnant, your partner can have carrier screening for Aicardi-Goutieres Syndrome, RNASEH2A-Related ordered by a health care professional. If your partner is found to be a carrier for Aicardi-Goutieres Syndrome, RNASEH2A-Related you have several reproductive options to consider:

- Natural pregnancy with or without prenatal diagnostic testing of the fetus or testing the baby after birth for Aicardi-Goutieres Syndrome, RNASEH2A-Related
- Preimplantation genetic diagnosis (PGD) with in vitro fertilization (IVF) to test embryos for Aicardi-Goutieres Syndrome, RNASEH2A-Related
- Adoption or use of a sperm or egg donor who is not a carrier for Aicardi-Goutieres Syndrome, RNASEH2A-Related

What resources are available?

- Genetics Home Reference: <http://ghr.nlm.nih.gov/condition/aicardi-goutieres-syndrome>
- United Leukodystrophy Foundation: www.ulf.org/aicardi-goutieres-syndrome
- Prenatal diagnosis by CVS: <http://www.marchofdimes.org/chorionic-villus-sampling.aspx>
- Prenatal diagnosis by amniocentesis: <http://www.marchofdimes.org/amniocentesis.aspx>
- PGD with IVF: <http://www.natera.com/spectrum>

Patient Information

Patient Name:

Test Information

Ordering Physician:



Clinic Information:

Date Of Birth:



Case File ID:



Report Date:

PRIMARY CONGENITAL GLAUCOMA/PETERS ANOMALY

Understanding Your Horizon Carrier Screen Results

What is Primary Congenital Glaucoma/Peters Anomaly?

Primary Congenital Glaucoma and Peters Anomaly are related inherited disorders that affect the eyes. Primary Congenital Glaucoma causes increased pressure in the eye (glaucoma), which usually starts before 1 year of age, although some children have symptoms that start later but usually before 5 years. The increasing pressure in the eye damages the optic nerves which causes worsening vision loss, and may cause bulging of the eyes, light sensitivity (photophobia) and excess tears. Early treatment with surgery and/or medications may prevent or lessen the vision loss. Peters Anomaly is less common and causes problems in the parts at the front of the eye: the cornea, lens, and iris. Children with Peters Anomaly have corneas that are partially or completely opaque instead of clear, which causes blurred and decreased vision. Some affected children also have lazy eye (amblyopia) and/or eyes that don't point in the same direction (strabismus). Other symptoms seen in some affected children include glaucoma, cataracts, and/or very small eyeballs (microphthalmia). There is no cure for either of these disorders and treatment is based on symptoms. Clinical trials involving potential new treatments for these conditions may be available (see www.clinicaltrials.gov).

What causes Primary Congenital Glaucoma/Peters Anomaly?

Primary Congenital Glaucoma and Peters Anomaly are caused by a gene change, or mutation, in both copies of the CYP1B1 gene pair. These mutations cause the genes to not work properly or not work at all. When both copies of the CYP1B1 gene do not work correctly, it leads to the symptoms of one of the two conditions described above. It is sometimes, but not always, possible to tell whether a specific mutation in the CYP1B1 gene will cause Primary Congenital Glaucoma or Peters Anomaly. Primary Congenital Glaucoma and Peters Anomaly are inherited in an autosomal recessive manner. This means that, in most cases, both parents must be carriers of a mutation in one copy of the CYP1B1 gene to have a child with one of these disorders. People who are carriers for Primary Congenital Glaucoma/Peters Anomaly are usually healthy and do not have symptoms, nor do they have either disorder themselves. Usually a child inherits two copies of each gene, one copy from the mother and one copy from the father. If the mother and father are both carriers for Primary Congenital Glaucoma/Peters Anomaly, there is a 1 in 4, or 25%, chance in each pregnancy for both partners to pass on their CYP1B1 gene mutations to the child, who will then have one of these disorders. Individuals found to carry more than one mutation for Primary Congenital Glaucoma/Peters Anomaly should discuss their risk for having an affected child with their health care provider. There are a number of other forms of both inherited Glaucoma and Peters Anomaly, each caused by mutations in different genes. A person who carries a mutation in the CYP1B1 gene is not likely to be at increased risk for having children with the other forms of these disorders.

What can I do next?

You may wish to speak with a local genetic counselor about your carrier test results. A genetic counselor in your area can be located on the National Society of Genetic Counselors website (www.nsgc.org). Your siblings and other relatives are at increased risk to also have this mutation. You are encouraged to inform your family members of your test results as they may wish to consider being tested themselves. If you are pregnant, your partner can have carrier screening for Primary Congenital Glaucoma/Peters Anomaly ordered by a health care professional. If your partner is not found to be a carrier for Primary Congenital Glaucoma/Peters Anomaly, your risk of having an affected child is greatly reduced. Couples at risk for having a baby with Primary Congenital Glaucoma/Peters Anomaly can opt to have prenatal diagnosis done through chorionic villus sampling (CVS) or amniocentesis during pregnancy or can choose to have the baby tested after birth. If you are not yet pregnant, your partner can have carrier screening for Primary Congenital Glaucoma/Peters Anomaly ordered by a health care professional. If your partner is found to be a carrier for Primary Congenital Glaucoma/Peters Anomaly, you have several reproductive options to consider:

- Natural pregnancy with or without prenatal diagnosis of the fetus or testing the baby after birth for Primary Congenital Glaucoma/Peters Anomaly
- Preimplantation genetic diagnosis (PGD) with in vitro fertilization (IVF) to test embryos for Primary Congenital Glaucoma/Peters Anomaly
- Adoption or use of a sperm or egg donor who is not a carrier for Primary Congenital Glaucoma/Peters Anomaly

What resources are available?

- Genetics Home Reference: <https://ghr.nlm.nih.gov/condition/early-onset-glaucoma>
- Genetics Home Reference: <https://ghr.nlm.nih.gov/condition/peters-anomaly>
- Prenatal diagnosis done through CVS: <http://www.marchofdimes.org/chorionic-villus-sampling.aspx>
- Prenatal diagnosis done through Amniocentesis: <http://www.marchofdimes.org/amniocentesis.aspx>
- PGD with IVF: <http://www.natera.com/spectrum>

Patient Information

Patient Name: [REDACTED]

Test Information

Ordering Physician: [REDACTED]



Date Of Birth: [REDACTED]

Case File ID: [REDACTED]

Clinic Information:

Report Date:

RETINITIS PIGMENTOSA 25**Understanding Your Horizon Carrier Screen Results****What is Retinitis Pigmentosa 25?**

Retinitis Pigmentosa 25 is one of a group of inherited eye disorders in which the retina, the area at the back of the eye that allows you to see, gradually stops working. Retinitis Pigmentosa 25 causes progressive vision loss. The age at which symptoms begin and the severity of the condition varies from person to person. The first symptom is usually loss of night vision. Over time, loss of peripheral vision (tunnel vision) develops. Then, loss of central vision occurs. RP affects only the vision. Currently there is no cure or specific treatment to prevent the vision loss. Clinical trials involving potential new treatments for this condition may be available (see www.clinicaltrials.gov).

What causes Retinitis Pigmentosa 25?

Retinitis Pigmentosa can be caused by mutations in one of a number of different genes with different inheritance patterns. Retinitis Pigmentosa 25 is caused by a gene change, or mutation, in both copies of the EYS gene pair. These mutations cause the genes to not work properly or not work at all. When both copies of this gene do not work correctly, it results in the progressive vision loss described above. Retinitis Pigmentosa 25 is inherited in an autosomal recessive manner. This means that, in most cases, both parents must be carriers of a mutation in one copy of the EYS gene to have a child with Retinitis Pigmentosa 25. People who are carriers for Retinitis Pigmentosa 25 are usually healthy and do not have symptoms nor do they have Retinitis Pigmentosa themselves. Usually a child inherits two copies of each gene, one copy from the mother and one copy from the father. If the mother and father are both carriers for Retinitis Pigmentosa 25, there is a 1 in 4, or 25%, chance in each pregnancy for both partners to pass on their EYS gene mutations to the child, who will then have this condition. Individuals found to carry more than one mutation for Retinitis Pigmentosa 25 should discuss any potential effects to their own vision and their risk for having an affected child with their health care provider.

What can I do next?

You may wish to speak with a local genetic counselor about your carrier test results. A genetic counselor in your area can be located on the National Society of Genetic Counselors website (www.nsgc.org). Your siblings and other relatives are at increased risk to also have this mutation. You are encouraged to inform your family members of your test results as they may wish to consider being tested themselves. If you are pregnant, your partner can have carrier screening for Retinitis Pigmentosa 25 ordered by a health care professional. If your partner is not found to be a carrier for Retinitis Pigmentosa 25, your risk of having a child with Retinitis Pigmentosa 25 is greatly reduced. Couples at risk of having a baby with Retinitis Pigmentosa 25 can opt to have prenatal diagnosis done through chorionic villus sampling (CVS) or amniocentesis during pregnancy or can choose to have the baby tested after birth for this condition. If you are not yet pregnant, your partner can have carrier screening for Retinitis Pigmentosa 25 ordered by a health care professional. If your partner is found to be a carrier for Retinitis Pigmentosa 25, you have several reproductive options to consider:

- Natural pregnancy with or without prenatal diagnosis of the fetus or testing the baby after birth for Retinitis Pigmentosa 25
- Preimplantation genetic diagnosis (PGD) with in vitro fertilization (IVF) to test embryos for Retinitis Pigmentosa 25
- Adoption or use of a sperm or egg donor who is not a carrier for Retinitis Pigmentosa 25

What resources are available?

- Genetics Home Reference: <http://ghr.nlm.nih.gov/condition/retinitis-pigmentosa>
- Prenatal diagnosis done through CVS: <http://www.marchofdimes.org/chorionic-villus-sampling.aspx>
- Prenatal diagnosis done through Amniocentesis: <http://www.marchofdimes.org/amniocentesis.aspx>
- PGD with IVF: <http://www.natera.com/spectrum>

Patient Information

Patient Name:

Test Information

Ordering Physician:



Clinic Information:

Date Of Birth:



Case File ID:



Report Date:

XERODERMA PIGMENTOSUM, GROUP A

Understanding Your Horizon Carrier Screen Results

What is Xeroderma Pigmentosum, Group A?

Xeroderma Pigmentosum, Group A is an inherited condition that causes the body to have extreme sensitivity to ultraviolet (UV) rays from sunlight. This condition affects the eyes and areas of skin exposed to the sun, and in some cases affects the brain and nervous system. Signs and symptoms of Xeroderma Pigmentosum, Group A usually appear in infancy or early childhood. Affected children can develop a severe sunburn after spending only a few minutes in the sun. By age 2, almost all children with Xeroderma Pigmentosum, Group A have freckling in sun-exposed areas, dry skin, and patchy changes in the color of the skin. The eyes are very sensitive to the sun, leading to irritation and other eye problems that can affect vision. People with Xeroderma Pigmentosum, Group A have a greatly increased risk of developing skin cancer. Unless protected from the sun, about half of children will develop a skin cancer by age 10. Most people with Xeroderma Pigmentosum, Group A will have a number of skin cancers during their life. There may also be an increased risk for other types of cancer, including eye cancer and brain tumors, although more studies need to be done to know for sure. Lifelong medical care is important for both prevention and early detection of skin cancers. Some people with Xeroderma Pigmentosum, Group A develop neurological problems that worsen over time. These problems may include hearing loss, poor coordination, difficulty walking, movement problems, loss of intellectual abilities, difficulty swallowing and talking, and seizures. Currently there is no cure for this disorder and treatment is based on symptoms. Clinical trials involving potential new treatments for this condition may be available (see www.clinicaltrials.gov).

What causes Xeroderma Pigmentosum, Group A?

Xeroderma Pigmentosum, Group A is caused by a change, or mutation, in both copies of the XPA gene pair. These mutations cause the genes to not work properly or not work at all. The normal function of the XPA gene pair is to help repair the body's DNA. When both copies of the XPA gene do not work correctly, DNA mistakes are left uncorrected, leading to the symptoms described above. Xeroderma Pigmentosum, Group A is inherited in an autosomal recessive manner. This means that, in most cases, both parents must be carriers of a mutation in one copy of the XPA gene pair to have a child with Xeroderma Pigmentosum, Group A. People who are carriers for Xeroderma Pigmentosum, Group A are usually healthy and do not have symptoms nor do they have Xeroderma Pigmentosum, Group A themselves. Usually a child inherits two copies of each gene, one copy from the mother and one copy from the father. If the mother and father are both carriers for Xeroderma Pigmentosum, Group A there is a 1 in 4, or 25%, chance in each pregnancy for both partners to pass on their XPA gene mutations to the child, who will then have Xeroderma Pigmentosum, Group A. Individuals found to carry more than one mutation for Xeroderma Pigmentosum, Group A should discuss their risk for having an affected child with their health care provider. There are several forms of Xeroderma Pigmentosum, each caused by mutations in different genes. Carriers for Xeroderma Pigmentosum, Group A are not likely to be at increased risk of having a child with these other forms of this group of disorders.

What can I do next?

You may wish to speak with a local genetic counselor about your carrier test results. A genetic counselor in your area can be located on the National Society of Genetic Counselors website (www.nsgc.org). Your siblings and other relatives are at increased risk to also have this mutation. You are encouraged to inform your family members of your test results as they may wish to consider being tested themselves. If you are pregnant, your partner can have carrier screening for Xeroderma Pigmentosum, Group A ordered by a health care professional. If your partner is not found to be a carrier for Xeroderma Pigmentosum, Group A, your risk of having a child with this condition is greatly reduced. Couples at risk of having a baby with Xeroderma Pigmentosum, Group A can opt to have prenatal diagnosis done through chorionic villus sampling (CVS) or amniocentesis during pregnancy or can choose to have the baby tested after birth. If you are not yet pregnant, your partner can have carrier screening for Xeroderma Pigmentosum, Group A ordered by a health care professional. If your partner is found to be a carrier for Xeroderma Pigmentosum, Group A, you have several reproductive options to consider:

- Natural pregnancy with or without prenatal diagnosis of the fetus or testing the baby after birth for Xeroderma Pigmentosum, Group A
- Preimplantation genetic diagnosis (PGD) with in vitro fertilization (IVF) to test embryos for Xeroderma Pigmentosum, Group A
- Adoption or use of a sperm or egg donor who is not a carrier for Xeroderma Pigmentosum, Group A

What resources are available?

- Genetics Home Reference: <https://ghr.nlm.nih.gov/condition/xeroderma-pigmentosum>
- Prenatal diagnosis done through CVS: <http://www.marchofdimes.org/chorionic-villus-sampling.aspx>
- Prenatal diagnosis done through Amniocentesis: <http://www.marchofdimes.org/amniocentesis.aspx> PGD with IVF: <http://www.natera.com/spectrum>

Patient Information

Patient Name:

Test Information

Ordering Physician:



Date Of Birth:



Case File ID:



Clinic Information:

Report Date:

VARIANT DETAILS**CYP1B1, c.535del (p.A179Rfs*18), pathogenic**

- The c.535del (p.A179Rfs*18) variant in the CYP1B1 gene has been observed at a frequency of 0.0048% in the gnomAD v2.1.1 dataset.
- This variant has been reported in a homozygous state or in conjunction with another variant in individual(s) with primary congenital glaucoma (PMID: 12372064).
- This premature termination variant is predicted to cause nonsense-mediated decay (NMD) in a gene where loss-of-function is a known mechanism of disease.
- This variant has been reported in ClinVar [ID: 523943].

EYS, c.2380C>T (p.R794*), pathogenic

- The c.2380C>T (p.R794*) variant in the EYS gene has been observed at a frequency of 0.0007% in the gnomAD v2.1.1 dataset.
- This variant has been reported in a homozygous state or in conjunction with another variant in individual(s) with retinitis pigmentosa 25 (PMID: 26161267, 29159838).
- This premature termination variant is predicted to cause nonsense-mediated decay (NMD) in a gene where loss-of-function is a known mechanism of disease.
- This variant has been reported in ClinVar [ID: 650440].

RNASEH2A, c.199+1G>C, likely pathogenic

- The c.199+1G>C variant in the RNASEH2A gene has been observed at a frequency of 0.0004% in the gnomAD v2.1.1 dataset.
- This canonical splicing variant is predicted to maintain the reading frame but disrupt a significant percentage of the protein length or a critical region of the protein, potentially disrupting normal protein function.
- This variant has been reported in ClinVar [ID: 2202408].

XPA, c.390-1G>C, pathogenic

- The c.390-1G>C variant in the XPA gene has been observed at a frequency of 0.0012% in the gnomAD v2.1.1 dataset.
- This variant has been reported in a homozygous state or in conjunction with another variant in individual(s) with Xeroderma pigmentosum (XP) (PMID: 27607234).
- This canonical splicing variant is predicted to alter the reading frame and cause nonsense-mediated decay (NMD) in a gene where loss-of-function is a known mechanism of disease.
- This variant has been described in ClinVar [ID: 264684].

Patient Information

Patient Name:

Test Information

Ordering Physician:



Clinic Information:

Date Of Birth:



Case File ID:



Report Date:

DISEASES SCREENED

Below is a list of all diseases screened and the result. Certain conditions have unique patient-specific numerical values, therefore, results for those conditions are formatted differently.

Autosomal Recessive**1**17-BETA HYDROXYSTEROID DEHYDROGENASE 3 DEFICIENCY (*HSD17B3*) **negative****3**

3-BETA-HYDROXYSTEROID DEHYDROGENASE TYPE II DEFICIENCY (*HSD3B2*) **negative**
3-HYDROXY-3-METHYLGLUTARYL-COENZYME A LYASE DEFICIENCY (*HMGCL*) **negative**
3-HYDROXYACYL-COA DEHYDROGENASE DEFICIENCY (*HADH*) **negative**
3-METHYLCROTONYL-CoA CARBOXYLASE 2 DEFICIENCY (*MCCC2*) **negative**
3-PHOSPHOGLYCERATE DEHYDROGENASE DEFICIENCY (*PHGDH*) **negative**

55-ALPHA-REDUCTASE DEFICIENCY (*SRD5A2*) **negative****6**6-PYRUVYL-TETRAHYDROPTERIN SYNTHASE (*PTPS*) DEFICIENCY (*PTS*) **negative****A**

ABCA4-RELATED CONDITIONS (*ABCA4*) **negative**
ABETALIPOPROTEINEMIA (*MTTP*) **negative**
ACHONDROGENESIS, TYPE 1B (*SLC26A2*) **negative**
ACHROMATOPSIA, CNGB3-RELATED (*CNGB3*) **negative**
ACRODERMATITIS ENTEROPATHICA (*SLC39A4*) **negative**
ACTION MYOCLONUS-RENAL FAILURE (AMRF) SYNDROME (*SCARB2*) **negative**
ACUTE INFANTILE LIVER FAILURE, TRMU-RELATED (*TRMU*) **negative**
ACYL-COA OXIDASE I DEFICIENCY (*ACOX1*) **negative**
AICARDI-GOUTIÈRES SYNDROME (*SAMHD1*) **negative**
AICARDI-GOUTIÈRES SYNDROME, RNASEH2A-RELATED (*RNASEH2A*) **see first page**
AICARDI-GOUTIÈRES SYNDROME, RNASEH2B-RELATED (*RNASEH2B*) **negative**
AICARDI-GOUTIÈRES SYNDROME, RNASEH2C-RELATED (*RNASEH2C*) **negative**
AICARDI-GOUTIÈRES SYNDROME, TREX1-RELATED (*TREX1*) **negative**
ALPHA-MANNOSIDOSIS (*MAN2B1*) **negative**
ALPHA-THALASSEMIA (*HBA1/HBA2*) **negative**
ALPORT SYNDROME, COL4A3-RELATED (*COL4A3*) **negative**
ALPORT SYNDROME, COL4A4-RELATED (*COL4A4*) **negative**
ALSTROM SYNDROME (*ALMS1*) **negative**
AMISH INFANTILE EPILEPSY SYNDROME (*ST3GAL5*) **negative**
ANDERMANN SYNDROME (*SLC12A6*) **negative**
ARGININE:GLYCINE AMIDINOTRANSFERASE DEFICIENCY (AGAT DEFICIENCY) (*GATM*) **negative**
ARGININEMIA (*ARG1*) **negative**
ARGININOSUCCINATE LYASE DEFICIENCY (*ASL*) **negative**
AROMATASE DEFICIENCY (*CYP19A1*) **negative**
ASPARAGINE SYNTHETASE DEFICIENCY (*ASNS*) **negative**
ASPARTYLGLYCOSAMINURIA (AGA) **negative**
ATAXIA WITH VITAMIN E DEFICIENCY (*TTPA*) **negative**
ATAXIA-TELANGEICTASIA (*ATM*) **negative**
ATAXIA-TELANGEICTASIA-LIKE DISORDER 1 (*MRE11*) **negative**
ATANSFERRINEMIA (*TF*) **negative**
AUTISM SPECTRUM, EPILEPSY AND ARTHROGRYPOSIS (*SLC35A3*) **negative**
AUTOIMMUNE POLYGLANDULAR SYNDROME, TYPE 1 (*AIRE*) **negative**
AUTOSOMAL RECESSIVE CONGENITAL ICHTHYOSIS (ARCI), SLC27A4-RELATED (*SLC27A4*) **negative**
AUTOSOMAL RECESSIVE SPASTIC ATAXIA OF CHARLEVOIX-SAGUENAY (SACS) **negative**

B

BARDET-BIEDL SYNDROME, ARL6-RELATED (*ARL6*) **negative**
BARDET-BIEDL SYNDROME, BBS10-RELATED (*BBS10*) **negative**
BARDET-BIEDL SYNDROME, BBS12-RELATED (*BBS12*) **negative**
BARDET-BIEDL SYNDROME, BBS1-RELATED (*BBS1*) **negative**
BARDET-BIEDL SYNDROME, BBS2-RELATED (*BBS2*) **negative**
BARDET-BIEDL SYNDROME, BBS4-RELATED (*BBS4*) **negative**
BARDET-BIEDL SYNDROME, BBS5-RELATED (*BBS5*) **negative**
BARDET-BIEDL SYNDROME, BBS7-RELATED (*BBS7*) **negative**
BARDET-BIEDL SYNDROME, BBS9-RELATED (*BBS9*) **negative**
BARDET-BIEDL SYNDROME, TTC8-RELATED (*TTC8*) **negative**
BARE LYMPHOCYTE SYNDROME, CIITA-RELATED (*CIITA*) **negative**
BARTTER SYNDROME, BSND-RELATED (*BSND*) **negative**
BARTTER SYNDROME, KCNJ1-RELATED (*KCNJ1*) **negative**
BARTTER SYNDROME, SLC12A1-RELATED (*SLC12A1*) **negative**
BATTEN DISEASE, CLN3-RELATED (*CLN3*) **negative**
BETA-HEMOGLOBINOPATHIES (*HBB*) **negative**
BETA-KETOTHIOLASE DEFICIENCY (*ACAT1*) **negative**
BETA-MANNOSIDOSIS (*MANBA*) **negative**
BETA-UREIDOPROPIONASE DEFICIENCY (*UPB1*) **negative**
BILATERAL FRONTOPIRIETAL POLYMICROGYRIA (*GPR56*) **negative**

BIOTINIDASE DEFICIENCY (*BTD*) **negative**BIOTIN-THIAMINE-RESPONSIVE BASAL GANGLIA DISEASE (BTBGD) (*SLC19A3*) **negative**BLOOM SYNDROME (*BLM*) **negative**BRITTLE CORNEA SYNDROME 1 (*ZNF469*) **negative**BRITTLE CORNEA SYNDROME 2 (*PRDM5*) **negative****C**

CANAVAN DISEASE (*ASPA*) **negative**
CARBAMOYL PHOSPHATE SYNTHETASE I DEFICIENCY (*CPS1*) **negative**
CARNITINE DEFICIENCY (*SLC22A5*) **negative**
CARNITINE PALMITOYLTRANSFERASE IA DEFICIENCY (*CPT1A*) **negative**
CARNITINE PALMITOYLTRANSFERASE II DEFICIENCY (*CPT2*) **negative**
CARNITINE-ACYLCARNITINE TRANSLOCASE DEFICIENCY (*SLC25A20*) **negative**
CARPENTER SYNDROME (*RAB23*) **negative**
CARILAGE-HAIR HYPOPLASIA (*RMRP*) **negative**
CATECHOLAMINERGIC POLYMORPHIC VENTRICULAR TACHYCARDIA (*CASQ2*) **negative**
CD59-MEDIATED HEMOLYTIC ANEMIA (*CD59*) **negative**
CEP152-RELATED MICROCEPHALY (*CEP152*) **negative**
CEREBRAL DYSGENESIS, NEUROPATHY, ICHTHYOSIS, AND PALMOPLANTAR KERATODERMA (CEDNIK) SYNDROME (*SNAP29*) **negative**
CEREBROTENDINOUS XANTHOMATOSIS (*CYP27A1*) **negative**
CHARCOT-MARIE-TOOTH DISEASE, RECESSIVE INTERMEDIATE C (*PLEKHG5*) **negative**
CHARCOT-MARIE-TOOTH-DISEASE, TYPE 4D (*NDRG1*) **negative**
CHEDIAK-HIGASHI SYNDROME (*LYST*) **negative**
CHOREOACANTHOCYTOSIS (*VPS13A*) **negative**
CHRONIC GRANULOMATOUS DISEASE, CYBA-RELATED (*CYBA*) **negative**
CHRONIC GRANULOMATOUS DISEASE, NCF2-RELATED (*NCF2*) **negative**
CILIOPATHIES, RPGRIP1L-RELATED (*RPGRIP1L*) **negative**
CITRIN DEFICIENCY (*SLC25A13*) **negative**
CITRULLINEMIA, TYPE 1 (*ASS1*) **negative**
CLN10 DISEASE (*CTSD*) **negative**
COHEN SYNDROME (*VPS13B*) **negative**
COL11A2-RELATED CONDITIONS (*COL11A2*) **negative**
COMBINED MALONIC AND METHYLMALONIC ACIDURIA (*ACSF3*) **negative**
COMBINED OXIDATIVE PHOSPHORYLATION DEFICIENCY 1 (*GFM1*) **negative**
COMBINED OXIDATIVE PHOSPHORYLATION DEFICIENCY 3 (*TSFM*) **negative**
COMBINED PITUITARY HORMONE DEFICIENCY 1 (*POU1F1*) **negative**
COMBINED PITUITARY HORMONE DEFICIENCY-2 (*PROP1*) **negative**
CONGENITAL ADRENAL HYPERPLASIA, 11-BETA-HYDROXYLASE DEFICIENCY (*CYP11B1*) **negative**
CONGENITAL ADRENAL HYPERPLASIA, 17-ALPHA-HYDROXYLASE DEFICIENCY (*CYP17A1*) **negative**
CONGENITAL ADRENAL HYPERPLASIA, 21-HYDROXYLASE DEFICIENCY (*CYP21A2*) **negative**
CONGENITAL ADRENAL INSUFFICIENCY, CYP11A1-RELATED (*CYP11A1*) **negative**
CONGENITAL AMEGAKARYOCYTIC THROMBOCYTOPENIA (*MPL*) **negative**
CONGENITAL CHRONIC DIARRHEA (*DGAT1*) **negative**
CONGENITAL DISORDER OF GLYCOSYLATION TYPE 1, ALG1-RELATED (*ALG1*) **negative**
CONGENITAL DISORDER OF GLYCOSYLATION, TYPE 1A, PMM2-Related (*PMM2*) **negative**
CONGENITAL DISORDER OF GLYCOSYLATION, TYPE 1B (*MPL*) **negative**
CONGENITAL DISORDER OF GLYCOSYLATION, TYPE 1C (*ALG6*) **negative**
CONGENITAL DYSERYTHROPOIETIC ANEMIA TYPE 2 (*SEC23B*) **negative**
CONGENITAL FINNISH NEPHROSIS (*NPHS1*) **negative**
CONGENITAL HYDROCEPHALUS 1 (*CCDC88C*) **negative**
CONGENITAL HYPERINSULINISM, KCNJ11-Related (*KCNJ11*) **negative**
CONGENITAL INSENSITIVITY TO PAIN WITH ANHIDROSIS (CIPA) (*NTRK1*) **negative**
CONGENITAL MYASTHENIC SYNDROME, CHAT-RELATED (*CHAT*) **negative**
CONGENITAL MYASTHENIC SYNDROME, CHRNE-RELATED (*CHRNE*) **negative**
CONGENITAL MYASTHENIC SYNDROME, COLQ-RELATED (*COLQ*) **negative**
CONGENITAL MYASTHENIC SYNDROME, DOK7-RELATED (*DOK7*) **negative**
CONGENITAL MYASTHENIC SYNDROME, RAPSIN-RELATED (*RAPSIN*) **negative**
CONGENITAL NEPHROTIC SYNDROME, PLCE1-RELATED (*PLCE1*) **negative**
CONGENITAL NEUTROPENIA, G6PC3-RELATED (*G6PC3*) **negative**
CONGENITAL NEUTROPENIA, HAX1-RELATED (*HAX1*) **negative**
CONGENITAL NEUTROPENIA, VPS45-RELATED (*VPS45*) **negative**
CONGENITAL SECRETORY CHLORIDE DIARRHEA 1 (*SLC26A3*) **negative**
CORNEAL DYSTROPHY AND PERCEPTIVE DEAFNESS (*SLC4A11*) **negative**
CORTICOSTERONE METHYLOXIDASE DEFICIENCY (*CYP11B2*) **negative**
COSTEFF SYNDROME (3-METHYLGLUTACONIC ACIDURIA, TYPE 3) (*OPA3*) **negative**
CRB1-RELATED RETINAL DYSTROPHIES (*CRB1*) **negative**
CYSTIC FIBROSIS (*CFTR*) **negative**
CYSTINOSIS (*CTNS*) **negative**
CYTOCHROME C OXIDASE DEFICIENCY, PET100-RELATED (*PET100*) **negative**
CYTOCHROME P450 OXIDOREDUCTASE DEFICIENCY (*POR*) **negative**

DD-BIFUNCTIONAL PROTEIN DEFICIENCY (*HSD17B4*) **negative**

Patient Information

Patient Name:

Test Information

Ordering Physician:



Clinic Information:

Date Of Birth:



Case File ID:



Report Date:

D

DEAFNESS, AUTOSOMAL RECESSIVE 77 (LOXHD1) **negative**
DIHYDROPTERIDINE REDUCTASE (DHPR) DEFICIENCY (QDPR) **negative**
DONNAI-BARROW SYNDROME (LRP2) **negative**
DUBIN-JOHNSON SYNDROME (ABCC2) **negative**
DYSKERATOSIS CONGENITA SPECTRUM DISORDERS (TERT) **negative**
DYSKERATOSIS CONGENITA, RTTEL1-RELATED (RTTEL1) **negative**
DYSTROPHIC EPIDERMOLYSIS BULLOSA, COL7A1-Related (COL7A1) **negative**

E

EARLY INFANTILE EPILEPTIC ENCEPHALOPATHY, CAD-RELATED (CAD) **negative**
EHLERS-DANLOS SYNDROME TYPE VI (PLOD1) **negative**
EHLERS-DANLOS SYNDROME, CLASSIC-LIKE, TNXB-RELATED (TNXB) **negative**
EHLERS-DANLOS SYNDROME, TYPE VII C (ADAMTS2) **negative**
ELLIS-VAN CREVELD SYNDROME, EVC2-RELATED (EVC2) **negative**
ELLIS-VAN CREVELD SYNDROME, EVC-RELATED (EVC) **negative**
ENHANCED S-CONE SYNDROME (NR2E3) **negative**
EPIMERASE DEFICIENCY (GALACTOSEMIA TYPE III) (GALE) **negative**
EPIPHYSEAL DYSPLASIA, MULTIPLE, 7/DESBUQUOIS DYSPLASIA 1 (CANT1) **negative**
ERCC6-RELATED DISORDERS (ERCC6) **negative**
ERCC8-RELATED DISORDERS (ERCC8) **negative**
ETHYLMALONIC ENCEPHALOPATHY (ETHE1) **negative**

F

FACTOR XI DEFICIENCY (F11) **negative**
FAMILIAL DYSAUTONOMIA (IKBKAP) **negative**
FAMILIAL HEMOPHAGOCYTIC LYMPHOHISTIOCYTOSIS, PRF1-RELATED (PRF1) **negative**
FAMILIAL HEMOPHAGOCYTIC LYMPHOHISTIOCYTOSIS, STX11-RELATED (STX11) **negative**
FAMILIAL HEMOPHAGOCYTIC LYMPHOHISTIOCYTOSIS, STXBP2-RELATED (STXBP2) **negative**
FAMILIAL HEMOPHAGOCYTIC LYMPHOHISTIOCYTOSIS, UNC13D-RELATED (UNC13D) **negative**
FAMILIAL HYPERCHOLESTEROLEMIA, LDLRAP1-RELATED (LDLRAP1) **negative**
FAMILIAL HYPERCHOLESTEROLEMIA, LDLR-RELATED (LDLR) **negative**
FAMILIAL HYPERINSULINISM, ABCC8-RELATED (ABCC8) **negative**
FAMILIAL NEPHROGENIC DIABETES INSIPIDUS, AQP2-RELATED (AQP2) **negative**
FANCONI ANEMIA, GROUP A (FANCA) **negative**
FANCONI ANEMIA, GROUP C (FANCC) **negative**
FANCONI ANEMIA, GROUP D2 (FANCD2) **negative**
FANCONI ANEMIA, GROUP E (FANCE) **negative**
FANCONI ANEMIA, GROUP F (FANCF) **negative**
FANCONI ANEMIA, GROUP G (FANCG) **negative**
FANCONI ANEMIA, GROUP I (FANCI) **negative**
FANCONI ANEMIA, GROUP J (BRIP1) **negative**
FANCONI ANEMIA, GROUP L (FANCL) **negative**
FARBER LIPOGRANULOMATOSIS (ASAHI1) **negative**
FOVEAL HYPOPLASIA (SLC38A8) **negative**
FRASER SYNDROME 3, GRIP1-RELATED (GRIP1) **negative**
FRASER SYNDROME, FRAS1-RELATED (FRAS1) **negative**
FRASER SYNDROME, FREM2-RELATED (FREM2) **negative**
FRIEDREICH ATAXIA (FXN) **negative**
FRUCTOSE-1,6-BISPHOSPHATASE DEFICIENCY (FBP1) **negative**
FUCOSIDOSIS, FUCA1-RELATED (FUCA1) **negative**
FUMARASE DEFICIENCY (FH) **negative**

G

GABA-TRANSAMINASE DEFICIENCY (ABAT) **negative**
GALACTOKINASE DEFICIENCY (GALACTOSEMIA, TYPE II) (GALK1) **negative**
GALACTOSEMIA (GALT) **negative**
GALACTOSIALIDOSIS (CTSA) **negative**
GAUCHER DISEASE (GBA) **negative**
GCH1-RELATED CONDITIONS (GCH1) **negative**
GDF5-RELATED CONDITIONS (GDF5) **negative**
GERODERMA OSTEODYSPLASTICA (GORAB) **negative**
GITELMAN SYNDROME (SLC12A3) **negative**
GLANZMANN THROMBASTHENIA (ITGB3) **negative**
GLUTARIC ACIDEMIA, TYPE 1 (GCDH) **negative**
GLUTARIC ACIDEMIA, TYPE 2A (ETFA) **negative**
GLUTARIC ACIDEMIA, TYPE 2B (ETFB) **negative**
GLUTARIC ACIDEMIA, TYPE 2C (ETFDH) **negative**
GLUTATHIONE SYNTHETASE DEFICIENCY (GSS) **negative**
GLYCINE ENCEPHALOPATHY, AMT-RELATED (AMT) **negative**
GLYCINE ENCEPHALOPATHY, GLDC-RELATED (GLDC) **negative**
GLYCOGEN STORAGE DISEASE TYPE 5 (McArdle Disease) (PYGM) **negative**
GLYCOGEN STORAGE DISEASE TYPE IXB (PHKB) **negative**
GLYCOGEN STORAGE DISEASE TYPE IXC (PHKG2) **negative**
GLYCOGEN STORAGE DISEASE, TYPE 1a (G6PC) **negative**
GLYCOGEN STORAGE DISEASE, TYPE 1b (SLC37A4) **negative**
GLYCOGEN STORAGE DISEASE, TYPE 2 (POMPE DISEASE) (GAA) **negative**
GLYCOGEN STORAGE DISEASE, TYPE 3 (AGL) **negative**
GLYCOGEN STORAGE DISEASE, TYPE 4 (GBE1) **negative**
GLYCOGEN STORAGE DISEASE, TYPE 7 (PFKM) **negative**

GRACILE SYNDROME (BCS1L) **negative**
GUANIDINOACETATE METHYLTRANSFERASE DEFICIENCY (GAMT) **negative**

H

HARLEQUIN ICHTHYOSIS (ABCA12) **negative**
HEME OXYGENASE 1 DEFICIENCY (HMOX1) **negative**
HEMOCHROMATOSIS TYPE 2A (HFE2) **negative**
HEMOCHROMATOSIS, TYPE 3, TFR2-Related (TFR2) **negative**
HEPATOCEREBRAL MITOCHONDRIAL DNA DEPLETION SYNDROME, MPV17-RELATED (MPV17) **negative**
HEREDITARY FRUCTOSE INTOLERANCE (ALDOB) **negative**
HEREDITARY HEMOCHROMATOSIS TYPE 2B (HAMP) **negative**
HEREDITARY SPASTIC PARAPARESIS, TYPE 49 (TECPR2) **negative**
HEREDITARY SPASTIC PARAPLEGIA, CYP7B1-RELATED (CYP7B1) **negative**
HERMANSKY-PUDLAK SYNDROME, AP3B1-RELATED (AP3B1) **negative**
HERMANSKY-PUDLAK SYNDROME, BLOC1S3-RELATED (BLOC1S3) **negative**
HERMANSKY-PUDLAK SYNDROME, BLOC1S6-RELATED (BLOC1S6) **negative**
HERMANSKY-PUDLAK SYNDROME, HPS1-RELATED (HPS1) **negative**
HERMANSKY-PUDLAK SYNDROME, HPS3-RELATED (HPS3) **negative**
HERMANSKY-PUDLAK SYNDROME, HPS4-RELATED (HPS4) **negative**
HERMANSKY-PUDLAK SYNDROME, HPS5-RELATED (HPS5) **negative**
HERMANSKY-PUDLAK SYNDROME, HPS6-RELATED (HPS6) **negative**
HOLOCARBOXYLASE SYNTHETASE DEFICIENCY (HLCS) **negative**
HOMOCYSTINURIA AND MEGALOBlastic ANEMIA TYPE CBLG (MTR) **negative**
HOMOCYSTINURIA DUE TO DEFICIENCY OF MTHFR (MTHFR) **negative**
HOMOCYSTINURIA, CBS-RELATED (CBS) **negative**
HOMOCYSTINURIA, Type cblE (MTRR) **negative**
HYDROLETHALUS SYNDROME (HLYS1) **negative**
HYPER-IGM IMMUNODEFICIENCY (CD40) **negative**
HYPERORNITHINEMIA-HYPERAMMONEMIA-HOMOCITRULLINURIA (HHH SYNDROME) (SLC25A15) **negative**
HYPERPHOSPHATEMIC FAMILIAL TUMORAL CALCINOSIS, GALNT3-RELATED (GALNT3) **negative**
HYPOMYELINATING LEUKODYSTROPHY 12 (VPS11) **negative**
HYPOPHOSPHATASIA, ALPL-RELATED (ALPL) **negative**

I

IMERSLUND-GRÄSBECK SYNDROME 2 (AMN) **negative**
IMMUNODEFICIENCY-CENTROMERIC INSTABILITY-FACIAL ANOMALIES (ICF) SYNDROME, DNMT3B-RELATED (DNMT3B) **negative**
IMMUNODEFICIENCY-CENTROMERIC INSTABILITY-FACIAL ANOMALIES (ICF) SYNDROME, ZBTB24-RELATED (ZBTB24) **negative**
INCLUSION BODY MYOPATHY 2 (GNE) **negative**
INFANTILE CEREBRAL AND CEREBELLAR ATROPHY (MED17) **negative**
INFANTILE NEPHRONOPHTHISIS (INVS) **negative**
INFANTILE NEUROAXONAL DYSTROPHY (PLA2G6) **negative**
ISOLATED ECTOPIA LENTIS (ADAMTSL4) **negative**
ISOLATED SULFITE OXIDASE DEFICIENCY (SUOX) **negative**
ISOLATED THYROID-STIMULATING HORMONE DEFICIENCY (TSHB) **negative**
ISOVALERIC ACIDEMIA (IVD) **negative**

J

JOHANSON-BLIZZARD SYNDROME (UBR1) **negative**
JOUBERT SYNDROME 2 / MECKEL SYNDROME 2 (TMEM216) **negative**
JOUBERT SYNDROME AND RELATED DISORDERS (JSRD), TMEM67-RELATED (TMEM67) **negative**
JOUBERT SYNDROME, AHI1-RELATED (AHI1) **negative**
JOUBERT SYNDROME, ARL13B-RELATED (ARL13B) **negative**
JOUBERT SYNDROME, B9D1-RELATED (B9D1) **negative**
JOUBERT SYNDROME, B9D2-RELATED (B9D2) **negative**
JOUBERT SYNDROME, C2CD3-RELATED/OROFACIODIGITAL SYNDROME 14 (C2CD3) **negative**
JOUBERT SYNDROME, CC2D2A-RELATED/COACH SYNDROME (CC2D2A) **negative**
JOUBERT SYNDROME, CEP104-RELATED (CEP104) **negative**
JOUBERT SYNDROME, CEP120-RELATED/SHORT-RIB THORACIC DYSPLASIA 13 WITH OR WITHOUT POLYDACTYLY (CEP120) **negative**
JOUBERT SYNDROME, CEP41-RELATED (CEP41) **negative**
JOUBERT SYNDROME, CPLANE1-RELATED / OROFACIODIGITAL SYNDROME 6 (CPLANE1) **negative**
JOUBERT SYNDROME, CSPP1-RELATED (CSPP1) **negative**
JOUBERT SYNDROME, INPP5E-RELATED (INPP5E) **negative**
JUNCTIONAL EPIDERMOLYSIS BULLOSA, COL17A1-RELATED (COL17A1) **negative**
JUNCTIONAL EPIDERMOLYSIS BULLOSA, ITGA6-RELATED (ITGA6) **negative**
JUNCTIONAL EPIDERMOLYSIS BULLOSA, ITGB4-RELATED (ITGB4) **negative**
JUNCTIONAL EPIDERMOLYSIS BULLOSA, LAMB3-RELATED (LAMB3) **negative**
JUNCTIONAL EPIDERMOLYSIS BULLOSA, LAMC2-RELATED (LAMC2) **negative**
JUNCTIONAL EPIDERMOLYSIS BULLOSA/LARYNGOONYCHOCUTANEOUS SYNDROME, LAMA3-RELATED (LAMA3) **negative**

K

KRABBE DISEASE (GALC) **negative**

L

LAMELLAR ICHTHYOSIS, TYPE 1 (TGM1) **negative**

Patient Information

Patient Name:

Test Information

Ordering Physician:

Clinic Information:

Date Of Birth:

Case File ID:

Report Date:



L
LARON SYNDROME (*GHR*) **negative**
LEBER CONGENITAL AMAUROSIS 2 (*RPE65*) **negative**
LEBER CONGENITAL AMAUROSIS TYPE APL1 (*APL1*) **negative**
LEBER CONGENITAL AMAUROSIS TYPE GUCY2D (*GUCY2D*) **negative**
LEBER CONGENITAL AMAUROSIS TYPE TULP1 (*TULP1*) **negative**
LEBER CONGENITAL AMAUROSIS, IQCB1-RELATED/SENIOR-LOKEN SYNDROME 5 (*IQCB1*) **negative**
LEBER CONGENITAL AMAUROSIS, TYPE CEP290 (*CEP290*) **negative**
LEBER CONGENITAL AMAUROSIS, TYPE LCA5 (*LCA5*) **negative**
LEBER CONGENITAL AMAUROSIS, TYPE RDH12 (*RDH12*) **negative**
LEIGH SYNDROME, FRENCH-CANADIAN TYPE (*LRPPRC*) **negative**
LETHAL CONGENITAL CONTRACTURE SYNDROME 1 (*GLE1*) **negative**
LEUKOENCEPHALOPATHY WITH VANISHING WHITE MATTER (*EIF2B5*) **negative**
LEUKOENCEPHALOPATHY WITH VANISHING WHITE MATTER, EIF2B1-RELATED (*EIF2B1*) **negative**
LEUKOENCEPHALOPATHY WITH VANISHING WHITE MATTER, EIF2B2-RELATED (*EIF2B2*) **negative**
LEUKOENCEPHALOPATHY WITH VANISHING WHITE MATTER, EIF2B3-RELATED (*EIF2B3*) **negative**
LEUKOENCEPHALOPATHY WITH VANISHING WHITE MATTER, EIF2B4-RELATED (*EIF2B4*) **negative**
LIG4 SYNDROME (*LIG4*) **negative**
LIMB-GIRDLE MUSCULAR DYSTROPHY TYPE 8 (*TRIM32*) **negative**
LIMB-GIRDLE MUSCULAR DYSTROPHY, TYPE 2A (*CAPN3*) **negative**
LIMB-GIRDLE MUSCULAR DYSTROPHY, TYPE 2B (*DYSF*) **negative**
LIMB-GIRDLE MUSCULAR DYSTROPHY, TYPE 2C (*SGCG*) **negative**
LIMB-GIRDLE MUSCULAR DYSTROPHY, TYPE 2D (*SGCA*) **negative**
LIMB-GIRDLE MUSCULAR DYSTROPHY, TYPE 2E (*SGCB*) **negative**
LIMB-GIRDLE MUSCULAR DYSTROPHY, TYPE 2F (*SGCD*) **negative**
LIMB-GIRDLE MUSCULAR DYSTROPHY, TYPE 2I (*FKRP*) **negative**
LIPOAMIDE DEHYDROGENASE DEFICIENCY (DIHYDROLIPOAMIDE DEHYDROGENASE DEFICIENCY) (*DLD*) **negative**
LIPOID ADRENAL HYPERPLASIA (*STAR*) **negative**
LIPOPROTEIN LIPASE DEFICIENCY (*LPL*) **negative**
LONG CHAIN 3-HYDROXYACYL-COA DEHYDROGENASE DEFICIENCY (*HADHA*) **negative**
LRAT-RELATED CONDITIONS (*LRAT*) **negative**
LUNG DISEASE, IMMUNODEFICIENCY, AND CHROMOSOME BREAKAGE SYNDROME (LICS) (*NSMCE3*) **negative**
LYSINURIC PROTEIN INTOLERANCE (*SLC7A7*) **negative**

M
MALONYL-COA DECARBOXYLASE DEFICIENCY (*MLYCD*) **negative**
MAPLE SYRUP URINE DISEASE, TYPE 1A (*BCKDHA*) **negative**
MAPLE SYRUP URINE DISEASE, TYPE 1B (*BCKDHB*) **negative**
MAPLE SYRUP URINE DISEASE, TYPE 2 (*DBT*) **negative**
MCKUSICK-KAUFMAN SYNDROME (*MKKS*) **negative**
MECKEL SYNDROME 7/NEPHRONOPHTHISIS 3 (*NPHP3*) **negative**
MECKEL-GRUBER SYNDROME, TYPE 1 (*MKS1*) **negative**
MECR-RELATED NEUROLOGIC DISORDER (*MECR*) **negative**
MEDIUM CHAIN ACYL-CoA DEHYDROGENASE DEFICIENCY (*ACADM*) **negative**
MEDNIK SYNDROME (*AP1S1*) **negative**
MEGALENCEPHALIC LEUKOENCEPHALOPATHY WITH SUBCORTICAL CYSTS (*MLC1*) **negative**
MEROSIN-DEFICIENT MUSCULAR DYSTROPHY (*LAMA2*) **negative**
METABOLIC ENCEPHALOPATHY AND ARRHYTHMIAS, TANGO2-RELATED (*TANGO2*) **negative**
METACHROMATIC LEUKODYSTROPHY, ARSA-RELATED (*ARSA*) **negative**
METACHROMATIC LEUKODYSTROPHY, PSAP-RELATED (*PSAP*) **negative**
METHYLMALONIC ACIDEMIA AND HOMOCYSTINURIA TYPE CBLF (*LMBRD1*) **negative**
METHYLMALONIC ACIDEMIA, MCEE-RELATED (*MCEE*) **negative**
METHYLMALONIC ACIDURIA AND HOMOCYSTINURIA, TYPE CBLF (*MMACHC*) **negative**
METHYLMALONIC ACIDURIA AND HOMOCYSTINURIA, TYPE CblD (*MMADHC*) **negative**
METHYLMALONIC ACIDURIA, MMAA-RELATED (*MMAA*) **negative**
METHYLMALONIC ACIDURIA, MMAB-RELATED (*MMAB*) **negative**
METHYLMALONIC ACIDURIA, TYPE MUT(0) (*MUT*) **negative**
MEVALONIC KINASE DEFICIENCY (*MVK*) **negative**
MICROCEPHALIC OSTEODYSPLASTIC PRIMORDIAL DWARFISM TYPE II (*PCNT*) **negative**
MICROPHTHALMIA / ANOPHTHALMIA, VSX2-RELATED (*VSX2*) **negative**
MITOCHONDRIAL COMPLEX 1 DEFICIENCY, ACAD9-RELATED (*ACAD9*) **negative**
MITOCHONDRIAL COMPLEX 1 DEFICIENCY, NDUF5-RELATED (*NDUF5*) **negative**
MITOCHONDRIAL COMPLEX 1 DEFICIENCY, NDUF56-RELATED (*NDUF56*) **negative**
MITOCHONDRIAL COMPLEX I DEFICIENCY, NUCLEAR TYPE 1 (*NDUF54*) **negative**
MITOCHONDRIAL COMPLEX I DEFICIENCY, NUCLEAR TYPE 10 (*NDUF52*) **negative**
MITOCHONDRIAL COMPLEX I DEFICIENCY, NUCLEAR TYPE 17 (*NDUF56*) **negative**
MITOCHONDRIAL COMPLEX I DEFICIENCY, NUCLEAR TYPE 19 (*FOXRED1*) **negative**
MITOCHONDRIAL COMPLEX I DEFICIENCY, NUCLEAR TYPE 3 (*NDUF57*) **negative**
MITOCHONDRIAL COMPLEX I DEFICIENCY, NUCLEAR TYPE 4 (*NDUFV1*) **negative**
MITOCHONDRIAL COMPLEX IV DEFICIENCY, NUCLEAR TYPE 2, SCO2-RELATED (*SCO2*) **negative**
MITOCHONDRIAL COMPLEX IV DEFICIENCY, NUCLEAR TYPE 6 (*COX15*) **negative**
MITOCHONDRIAL DNA DEPLETION SYNDROME 2 (*TK2*) **negative**

MITOCHONDRIAL DNA DEPLETION SYNDROME 3 (*DGUOK*) **negative**
MITOCHONDRIAL MYOPATHY AND SIDEROBLASTIC ANEMIA (MLASA1) (*PUS1*) **negative**
MITOCHONDRIAL TRIFUNCTIONAL PROTEIN DEFICIENCY, HADHB-RELATED (*HADHB*) **negative**
MOLYBDENUM COFACTOR DEFICIENCY TYPE B (*MOC52*) **negative**
MOLYBDENUM COFACTOR DEFICIENCY, TYPE A (*MOC51*) **negative**
MUCOLIPIDOSIS II/III A (*GNPTAB*) **negative**
MUCOLIPIDOSIS III GAMMA (*GNPTG*) **negative**
MUCOLIPIDOSIS, TYPE IV (*MCOLN1*) **negative**
MUCOPOLYSACCHARIDOSIS, TYPE I (HURLER SYNDROME) (*IDUA*) **negative**
MUCOPOLYSACCHARIDOSIS, TYPE III A (SANFILIPPO A) (*SGSH*) **negative**
MUCOPOLYSACCHARIDOSIS, TYPE III B (SANFILIPPO B) (*NAGLU*) **negative**
MUCOPOLYSACCHARIDOSIS, TYPE III C (SANFILIPPO C) (*HGSNAT*) **negative**
MUCOPOLYSACCHARIDOSIS, TYPE III D (SANFILIPPO D) (*GNS*) **negative**
MUCOPOLYSACCHARIDOSIS, TYPE IV A (MORQUIO SYNDROME) (*GALNS*) **negative**
MUCOPOLYSACCHARIDOSIS, TYPE IV B/GM1 GANGLIOSIDOSIS (*GLB1*) **negative**
MUCOPOLYSACCHARIDOSIS, TYPE IX (*HYAL1*) **negative**
MUCOPOLYSACCHARIDOSIS, TYPE VI (MAROTEAUX-LAMY) (*ARSB*) **negative**
MUCOPOLYSACCHARIDOSIS, TYPE VII (*GUSB*) **negative**
MULIBREY NANISM (*TRIM37*) **negative**
MULTIPLE PTERYGIUM SYNDROME, CHRNG-RELATED/ESCOBAR SYNDROME (*CHNRG*) **negative**
MULTIPLE SULFATASE DEFICIENCY (*SUMF1*) **negative**
MUSCLE-EYE-BRAIN DISEASE, POMGNT1-RELATED (*POMGNT1*) **negative**
MUSCULAR DYSTROPHY-DYSTROGLYCANOPATHY (*RXYLT1*) **negative**
MUSK-RELATED CONGENITAL MYASTHENIC SYNDROME (*MUSK*) **negative**
MYONEUROGASTROINTESTINAL ENCEPHALOPATHY (MNGIE) (*TYMP*) **negative**
MYOTONIA CONGENITA (*CLCN1*) **negative**

N
N-ACETYLGLUTAMATE SYNTHASE DEFICIENCY (*NAGS*) **negative**
NEMALINE MYOPATHY, NEB-RELATED (*NEB*) **negative**
NEPHRONOPHTHISIS 1 (*NPHP1*) **negative**
NEURONAL CEROID LIPOFUSCINOSIS, CLN5-RELATED (*CLN5*) **negative**
NEURONAL CEROID LIPOFUSCINOSIS, CLN6-RELATED (*CLN6*) **negative**
NEURONAL CEROID LIPOFUSCINOSIS, CLN8-RELATED (*CLN8*) **negative**
NEURONAL CEROID LIPOFUSCINOSIS, MFSD8-RELATED (*MFSD8*) **negative**
NEURONAL CEROID LIPOFUSCINOSIS, PPT1-RELATED (*PPT1*) **negative**
NEURONAL CEROID LIPOFUSCINOSIS, TPP1-RELATED (*TPP1*) **negative**
NGLY1-CONGENITAL DISORDER OF GLYCOSYLATION (*NGLY1*) **negative**
NIEMANN-PICK DISEASE, TYPE C1 / D (*NPC1*) **negative**
NIEMANN-PICK DISEASE, TYPE C2 (*NPC2*) **negative**
NIEMANN-PICK DISEASE, TYPES A / B (*SMPD1*) **negative**
NIJMEGEN BREAKAGE SYNDROME (*NBN*) **negative**
NON-SYNDROMIC HEARING LOSS, GJB2-RELATED (*GJB2*) **negative**
NON-SYNDROMIC HEARING LOSS, MYO15A-RELATED (*MYO15A*) **negative**
NONSYNDROMIC HEARING LOSS, OTOA-RELATED (*OTOA*) **negative**
NONSYNDROMIC HEARING LOSS, OTOF-RELATED (*OTOF*) **negative**
NONSYNDROMIC HEARING LOSS, PJK-RELATED (*PJK*) **negative**
NONSYNDROMIC HEARING LOSS, SYNE4-RELATED (*SYNE4*) **negative**
NONSYNDROMIC HEARING LOSS, TMC1-RELATED (*TMC1*) **negative**
NONSYNDROMIC HEARING LOSS, TMPS53-RELATED (*TMPS53*) **negative**
NONSYNDROMIC INTELLECTUAL DISABILITY (*CC2D1A*) **negative**
NORMOPHOSPHATEMIC TUMORAL CALCINOSIS (*SAMD9*) **negative**

O
OCULOCUTANEOUS ALBINISM TYPE III (*TYRP1*) **negative**
OCULOCUTANEOUS ALBINISM TYPE IV (*SLC45A2*) **negative**
OCULOCUTANEOUS ALBINISM, OCA2-RELATED (*OCA2*) **negative**
OCULOCUTANEOUS ALBINISM, TYPES 1A AND 1B (*TYR*) **negative**
ODONTO-ONYCHO-DERMAL DYSPLASIA / SCHOPF-SCHULZ-PASSARGE SYNDROME (*WNT10A*) **negative**
OMENN SYNDROME, RAG2-RELATED (*RAG2*) **negative**
ORNITHINE AMINOTRANSFERASE DEFICIENCY (*OAT*) **negative**
OSTEOGENESIS IMPERFECTA TYPE VII (*CRTAP*) **negative**
OSTEOGENESIS IMPERFECTA TYPE VIII (*P3H1*) **negative**
OSTEOGENESIS IMPERFECTA TYPE XI (*FKBP10*) **negative**
OSTEOGENESIS IMPERFECTA TYPE XIII (*BMP1*) **negative**
OSTEOPETROSIS, INFANTILE MALIGNANT, TCIRG1-RELATED (*TCIRG1*) **negative**
OSTEOPETROSIS, OSTM1-RELATED (*OSTM1*) **negative**

P
PANTOTHENATE KINASE-ASSOCIATED NEURODEGENERATION (*PANK2*) **negative**
PAPILLON LEFÈVRE SYNDROME (*CTSC*) **negative**
PARKINSON DISEASE 15 (*FBXO7*) **negative**
PENDRED SYNDROME (*SLC26A4*) **negative**
PERLMAN SYNDROME (*DIS3L2*) **negative**
PGM3-CONGENITAL DISORDER OF GLYCOSYLATION (*PGM3*) **negative**
PHENYLKETONURIA (*PAH*) **negative**
PIGN-CONGENITAL DISORDER OF GLYCOSYLATION (*PIGN*) **negative**
PITUITARY HORMONE DEFICIENCY, COMBINED 3 (*LHX3*) **negative**
POLG-RELATED DISORDERS (*POLG*) **negative**

Patient Information

Patient Name:

Test Information

Ordering Physician:



Clinic Information:

Date Of Birth:



Case File ID:



Report Date:

P

POLYCYSTIC KIDNEY DISEASE, AUTOSOMAL RECESSIVE (*PKHD1*) **negative**
PONTOCEREBELLAR HYPOPLASIA, EXOSC3-RELATED (*EXOSC3*) **negative**
PONTOCEREBELLAR HYPOPLASIA, RARS2-RELATED (*RARS2*) **negative**
PONTOCEREBELLAR HYPOPLASIA, TSEN2-RELATED (*TSEN2*) **negative**
PONTOCEREBELLAR HYPOPLASIA, TSEN54-RELATED (*TSEN54*) **negative**
PONTOCEREBELLAR HYPOPLASIA, TYPE 1A (*VRK1*) **negative**
PONTOCEREBELLAR HYPOPLASIA, TYPE 2D (*SEPSECS*) **negative**
PONTOCEREBELLAR HYPOPLASIA, VPS53-RELATED (*VPS53*) **negative**
PRIMARY CILIARY DYSKINESIA, CCDC103-RELATED (*CCDC103*) **negative**
PRIMARY CILIARY DYSKINESIA, CCDC39-RELATED (*CCDC39*) **negative**
PRIMARY CILIARY DYSKINESIA, DNAH11-RELATED (*DNAH11*) **negative**
PRIMARY CILIARY DYSKINESIA, DNAH5-RELATED (*DNAH5*) **negative**
PRIMARY CILIARY DYSKINESIA, DNAI1-RELATED (*DNAI1*) **negative**
PRIMARY CILIARY DYSKINESIA, DNAI2-RELATED (*DNAI2*) **negative**
PRIMARY CONGENITAL GLAUCOMA/PETERS ANOMALY (*CYP1B1*) **see first page**
PRIMARY HYPEROXALURIA, TYPE 1 (*AGXT*) **negative**
PRIMARY HYPEROXALURIA, TYPE 2 (*GRHPR*) **negative**
PRIMARY HYPEROXALURIA, TYPE 3 (*HOGA1*) **negative**
PRIMARY MICROCEPHALY 1, AUTOSOMAL RECESSIVE (*MCPH1*) **negative**
PROGRESSIVE EARLY-ONSET ENCEPHALOPATHY WITH BRAIN ATROPHY AND THIN CORPUS CALLOSUM (*TBCD*) **negative**
PROGRESSIVE FAMILIAL INTRAHEPATIC CHOLESTASIS, ABCB4-RELATED (*ABCB4*) **negative**
PROGRESSIVE FAMILIAL INTRAHEPATIC CHOLESTASIS, TYPE 1 (*PFIC1*) (*ATP8B1*) **negative**
PROGRESSIVE FAMILIAL INTRAHEPATIC CHOLESTASIS, TYPE 2 (*ABCB11*) **negative**
PROGRESSIVE FAMILIAL INTRAHEPATIC CHOLESTASIS, TYPE 4 (*PFIC4*) (*TJP2*) **negative**
PROGRESSIVE PSEUDORHEUMATOID DYSPLASIA (*CCN6*) **negative**
PROLIDASE DEFICIENCY (*PEPD*) **negative**
PROPIONIC ACIDEMIA, PCCA-RELATED (*PCCA*) **negative**
PROPIONIC ACIDEMIA, PCCB-RELATED (*PCCB*) **negative**
PSEUDOXANTHOMA ELASTICUM (*ABCC6*) **negative**
PTERIN-4 ALPHA-CARBINOLAMINE DEHYDRATASE (*PCD*) DEFICIENCY (*PCBD1*) **negative**
PYCNODYSTOSIS (*CTSK*) **negative**
PYRIDOXAL 5'-PHOSPHATE-DEPENDENT EPILEPSY (*PNPO*) **negative**
PYRIDOXINE-DEPENDENT EPILEPSY (*ALDH7A1*) **negative**
PYRUVATE CARBOXYLASE DEFICIENCY (*PC*) **negative**
PYRUVATE DEHYDROGENASE DEFICIENCY, PDHB-RELATED (*PDHB*) **negative**

R

REFSUM DISEASE, PHYH-RELATED (*PHYH*) **negative**
RENAL TUBULAR ACIDOSIS AND DEAFNESS, ATP6V1B1-RELATED (*ATP6V1B1*) **negative**
RENAL TUBULAR ACIDOSIS, PROXIMAL, WITH OCULAR ABNORMALITIES AND MENTAL RETARDATION (*SLC4A4*) **negative**
RETINITIS PIGMENTOSA 25 (*EYS*) **see first page**
RETINITIS PIGMENTOSA 26 (*CERKL*) **negative**
RETINITIS PIGMENTOSA 28 (*FAM161A*) **negative**
RETINITIS PIGMENTOSA 36 (*PRCD*) **negative**
RETINITIS PIGMENTOSA 59 (*DHDDS*) **negative**
RETINITIS PIGMENTOSA 62 (*MAK*) **negative**
RHIZOMELIC CHONDRODYSPLASIA PUNCTATA, TYPE 1 (*PEX7*) **negative**
RHIZOMELIC CHONDRODYSPLASIA PUNCTATA, TYPE 2 (*GNPAT*) **negative**
RHIZOMELIC CHONDRODYSPLASIA PUNCTATA, TYPE 3 (*AGPS*) **negative**
RLBP1-RELATED RETINOPATHY (*RLBP1*) **negative**
ROBERTS SYNDROME (*ESCO2*) **negative**
RYYR1-RELATED CONDITIONS (*RYYR1*) **negative**

S

SALLA DISEASE (*SLC17A5*) **negative**
SANDHOFF DISEASE (*HEXB*) **negative**
SCHIMKE IMMUNOSKELETAL DYSPLASIA (*SMARCA1*) **negative**
SCHINDLER DISEASE (*NAGA*) **negative**
SEGAWA SYNDROME, TH-RELATED (*TH*) **negative**
SENIOR-LOKEN SYNDROME 4/NEPHRONOPHTHISIS 4 (*NPHP4*) **negative**
SEPIAPTERIN REDUCTASE DEFICIENCY (*SPR*) **negative**
SEVERE COMBINED IMMUNODEFICIENCY (*SCID*), CD3D-RELATED (*CD3D*) **negative**
SEVERE COMBINED IMMUNODEFICIENCY (*SCID*), CD3E-RELATED (*CD3E*) **negative**
SEVERE COMBINED IMMUNODEFICIENCY (*SCID*), FOXP1-RELATED (*FOXP1*) **negative**
SEVERE COMBINED IMMUNODEFICIENCY (*SCID*), IKKB-RELATED (*IKKB*) **negative**
SEVERE COMBINED IMMUNODEFICIENCY (*SCID*), IL7R-RELATED (*IL7R*) **negative**
SEVERE COMBINED IMMUNODEFICIENCY (*SCID*), JAK3-RELATED (*JAK3*) **negative**
SEVERE COMBINED IMMUNODEFICIENCY (*SCID*), PTPRC-RELATED (*PTPRC*) **negative**
SEVERE COMBINED IMMUNODEFICIENCY (*SCID*), RAG1-RELATED (*RAG1*) **negative**
SEVERE COMBINED IMMUNODEFICIENCY, ADA-Related (*ADA*) **negative**
SEVERE COMBINED IMMUNODEFICIENCY, TYPE ATHABASKAN (*DCLRE1C*) **negative**
SHORT-RIB THORACIC DYSPLASIA 3 WITH OR WITHOUT POLYDACTYL (*DYNC2H1*) **negative**
SHWACHMAN-DIAMOND SYNDROME, SBDS-RELATED (*SBDS*) **negative**
SIALIDOSIS (*NEU1*) **negative**
SJÖGREN-LARSSON SYNDROME (*ALDH3A2*) **negative**
SMITH-LEMLI-OPITZ SYNDROME (*DHCR7*) **negative**
SPASTIC PARAPLEGIA, TYPE 15 (*ZFYVE26*) **negative**

SPASTIC TETRAPLEGIA, THIN CORPUS CALLOSUM, AND PROGRESSIVE MICROCEPHALY (*SPATCCM*) (*SLC1A4*) **negative**
SPG11-RELATED CONDITIONS (*SPG11*) **negative**
SPINAL MUSCULAR ATROPHY (*SMN1*) **negative** **SMN1: Two copies; g.27134T>G: absent; the absence of the g.27134T>G variant decreases the chance to be a silent (2+0) carrier.**
SPINAL MUSCULAR ATROPHY WITH RESPIRATORY DISTRESS TYPE 1 (*IGHMBP2*) **negative**
SPINOCEREBELLAR ATAXIA, AUTOSOMAL RECESSIVE 10 (*ANO10*) **negative**
SPINOCEREBELLAR ATAXIA, AUTOSOMAL RECESSIVE 12 (*WWOX*) **negative**
SPONDYLOCOSTAL DYSOSTOSIS 1 (*DLL3*) **negative**
SPONDYLOTHORACIC DYSOSTOSIS, MESP2-Related (*MESP2*) **negative**
STEEL SYNDROME (*COL27A1*) **negative**
STEROID-RESISTANT NEPHROTIC SYNDROME (*NPHS2*) **negative**
STUVE-WIEDEMANN SYNDROME (*LIFR*) **negative**
SURF1-RELATED CONDITIONS (*SURF1*) **negative**
SURFACTANT DYSFUNCTION, ABCA3-RELATED (*ABCA3*) **negative**

T

TAY-SACHS DISEASE (*HEXA*) **negative**
TBCE-RELATED CONDITIONS (*TBCE*) **negative**
THIAMINE-RESPONSIVE MEGALOBlastic ANEMIA SYNDROME (*SLC19A2*) **negative**
THYROID DYSHORMONOGENESIS 1 (*SLC5A5*) **negative**
THYROID DYSHORMONOGENESIS 2A (*TPO*) **negative**
THYROID DYSHORMONOGENESIS 3 (*TG*) **negative**
THYROID DYSHORMONOGENESIS 6 (*DUOX2*) **negative**
TRANSCOBALAMIN II DEFICIENCY (*TCN2*) **negative**
TRICHOHEPATOENTERIC SYNDROME, SKIC2-RELATED (*SKIC2*) **negative**
TRICHOHEPATOENTERIC SYNDROME, TTC37-RELATED (*TTC37*) **negative**
TRICHOHYDROTIC SYNDROME 1/XERODERMA PIGMENTOSUM, GROUP D (*ERCC2*) **negative**
TRIMETHYLAMINURIA (*FMO3*) **negative**
TRIPLE A SYNDROME (*AAAS*) **negative**
TSHR-RELATED CONDITIONS (*TSHR*) **negative**
TYROSINEMIA TYPE III (*HPD*) **negative**
TYROSINEMIA, TYPE 1 (*FAH*) **negative**
TYROSINEMIA, TYPE 2 (*TAT*) **negative**

U

USHER SYNDROME, TYPE 1B (*MYO7A*) **negative**
USHER SYNDROME, TYPE 1C (*USH1C*) **negative**
USHER SYNDROME, TYPE 1D (*CDH23*) **negative**
USHER SYNDROME, TYPE 1F (*PCDH15*) **negative**
USHER SYNDROME, TYPE 1J/DEAFNESS, AUTOSOMAL RECESSIVE, 48 (*CIB2*) **negative**
USHER SYNDROME, TYPE 2A (*USH2A*) **negative**
USHER SYNDROME, TYPE 2C (*ADGRV1*) **negative**
USHER SYNDROME, TYPE 3 (*CLRN1*) **negative**

V

VERY LONG-CHAIN ACYL-CoA DEHYDROGENASE DEFICIENCY (*ACADVL*) **negative**
VICI SYNDROME (*EPG5*) **negative**
VITAMIN D-DEPENDENT RICKETS, TYPE 1A (*CYP27B1*) **negative**
VITAMIN D-RESISTANT RICKETS TYPE 2A (*VDR*) **negative**
VLDLR-ASSOCIATED CEREBELLAR HYPOPLASIA (*VLDLR*) **negative**

W

WALKER-WARBURG SYNDROME, CRPPA-RELATED (*CRPPA*) **negative**
WALKER-WARBURG SYNDROME, FKTN-RELATED (*FKTN*) **negative**
WALKER-WARBURG SYNDROME, LARGE1-RELATED (*LARGE1*) **negative**
WALKER-WARBURG SYNDROME, POMT1-RELATED (*POMT1*) **negative**
WALKER-WARBURG SYNDROME, POMT2-RELATED (*POMT2*) **negative**
WARSAW BREAKAGE SYNDROME (*DDX11*) **negative**
WERNER SYNDROME (*WRN*) **negative**
WILSON DISEASE (*ATP7B*) **negative**
WOLCOTT-RALLISON SYNDROME (*EIF2AK3*) **negative**
WOLMAN DISEASE (*LIPA*) **negative**
WOODHOUSE-SAKATI SYNDROME (*DCAF17*) **negative**

X

XERODERMA PIGMENTOSUM VARIANT TYPE (*POLH*) **negative**
XERODERMA PIGMENTOSUM, GROUP A (*XPA*) **see first page**
XERODERMA PIGMENTOSUM, GROUP C (*XPC*) **negative**

Z

ZELLWEGER SPECTRUM DISORDER, PEX13-RELATED (*PEX13*) **negative**
ZELLWEGER SPECTRUM DISORDER, PEX16-RELATED (*PEX16*) **negative**
ZELLWEGER SPECTRUM DISORDER, PEX5-RELATED (*PEX5*) **negative**
ZELLWEGER SPECTRUM DISORDERS, PEX10-RELATED (*PEX10*) **negative**
ZELLWEGER SPECTRUM DISORDERS, PEX12-RELATED (*PEX12*) **negative**
ZELLWEGER SPECTRUM DISORDERS, PEX1-RELATED (*PEX1*) **negative**
ZELLWEGER SPECTRUM DISORDERS, PEX26-RELATED (*PEX26*) **negative**
ZELLWEGER SPECTRUM DISORDERS, PEX2-RELATED (*PEX2*) **negative**

Patient Information

Patient Name:

Test Information

Ordering Physician:



Date Of Birth:



Case File ID:



Clinic Information:

Report Date:

Z

ZELLWEGER SPECTRUM DISORDERS, PEX6-RELATED (PEX6) **negative**

Patient Information

Patient Name: [REDACTED]

Test Information

Ordering Physician: [REDACTED]



Date Of Birth: [REDACTED]

Clinic Information:

Case File ID: [REDACTED]

Report Date:

Testing Methodology, Limitations, and Comments:**Next-generation sequencing (NGS)**

Sequencing library prepared from genomic DNA isolated from a patient sample is enriched for targets of interest using standard hybridization capture protocols and PCR amplification (for targets specified below). NGS is then performed to achieve the standards of quality control metrics, including a minimum coverage of 99% of targeted regions at 20X sequencing depth. Sequencing data is aligned to human reference sequence, followed by deduplication, metric collection and variant calling (coding region +/- 20bp). Variants are then classified according to ACMGG/AMP standards of interpretation using publicly available databases including but not limited to ENSEMBL, HGMD Pro, ClinGen, ClinVar, 1000G, ESP and gnomAD. Variants predicted to be pathogenic or likely pathogenic for the specified diseases are reported. It should be noted that the data interpretation is based on our current understanding of the genes and variants at the time of reporting. Putative positive sequencing variants that do not meet internal quality standards or are within highly homologous regions are confirmed by Sanger sequencing or gene-specific long-range PCR as needed prior to reporting.

Copy Number Variant (CNV) analysis is limited to deletions involving two or more exons for all genes on the panel, in addition to specific known recurrent single-exon deletions. CNVs of small size may have reduced detection rate. This method does not detect gene inversions, single-exonic and sub-exonic deletions (unless otherwise specified), and duplications of all sizes (unless otherwise specified). Additionally, this method does not define the exact breakpoints of detected CNV events. Confirmation testing for copy number variation is performed by specific PCR, Multiplex Ligation-dependent Probe Amplification (MLPA), next generation sequencing, or other methodology.

This test may not detect certain variants due to local sequence characteristics, high/low genomic complexity, homologous sequence, or allele dropout (PCR-based assays). Variants within noncoding regions (promoter, 5'UTR, 3'UTR, deep intronic regions, unless otherwise specified), small deletions or insertions larger than 25bp, low-level mosaic variants, structural variants such as inversions, and/or balanced translocations may not be detected with this technology.

SPECIAL NOTES

For ABCC6, sequencing variants in exons 1-7 are not detected due to the presence of regions of high homology.

For CFTR, when the CFTR R117H variant is detected, reflex analysis of the polythymidine variations (5T, 7T and 9T) at the intron 9 branch/acceptor site of the CFTR gene will be performed. Multi-exon duplication analysis is included.

For CYP21A2, targets were enriched using long-range PCR amplification, followed by next generation sequencing. Duplication analysis will only be performed and reported when c.955C>T (p.Q319*) is detected. Sequencing and CNV analysis may have reduced sensitivity, if variants result from complex rearrangements, in trans with a gene deletion, or CYP21A2 gene duplication on one chromosome and deletion on the other chromosome. This analysis cannot detect sequencing variants located on the CYP21A2 duplicated copy.

For DDX11, sequencing variants in exons 7-11 and CNV for the entire gene are not analyzed due to high sequence homology.

For GJB2, CNV analysis of upstream deletions of GJB6-D13S1830 (309kb deletion) and GJB6-D13S1854 (232kb deletion) is included.

For HBA1/HBA2, CNV analysis is offered to detect common deletions of -alpha3.7, -alpha4.2, --MED, --SEA, --FIL, --THAI, --alpha20.5, and/or HS-40.

For OTOA, sequencing variants in exons 25-29 and CNV in exons 21-29 are not analyzed due to high sequence homology.

For RPGRIP1L, variants in exon 23 are not detected due to assay limitation.

For SAMD9, only p.K1495E variant will be analyzed and reported.

Friedreich Ataxia (FXN)

The GAA repeat region of the FXN gene is assessed by trinucleotide PCR assay and capillary electrophoresis. Variances of +/-1 repeat for normal alleles and up to +/-3 repeats for premutation alleles may occur. For fully penetrant expanded alleles, the precise repeat size cannot be determined, therefore the approximate allele size is reported. Sequencing and copy number variants are analyzed by next-generation sequencing analysis.

Friedreich Ataxia Repeat Categories

Categories	GAA Repeat Sizes
Normal	<34
Premutation	34 - 65
Full	>65

Patient Information

Patient Name:

Test Information

Ordering Physician:



Date Of Birth:

Case File ID:

Clinic Information:

Report Date:

Spinal Muscular Atrophy (SMN1)

The total combined copy number of SMN1 and SMN2 exon 7 is quantified based on NGS read depth. The ratio of SMN1 to SMN2 is calculated based on the read depth of a single nucleotide that distinguishes these two genes in exon 7. In addition to copy number analysis, testing for the presence or absence of a single nucleotide polymorphism (g.27134T>G in intron 7 of SMN1) associated with the presence of a SMN1 duplication allele is performed using NGS.

Ethnicity	Two SMN1 copies carrier risk before g.27134T>G testing	Carrier risk after g.27134T>G testing	
		g.27134T>G ABSENT	g.27134T>G PRESENT
Caucasian	1 in 632	1 in 769	1 in 29
Ashkenazi Jewish	1 in 350	1 in 580	LIKELY CARRIER
Asian	1 in 628	1 in 702	LIKELY CARRIER
African-American	1 in 121	1 in 396	1 in 34
Hispanic	1 in 1061	1 in 1762	1 in 140

Variant Classification

Only pathogenic or likely pathogenic variants are reported. Other variants including benign variants, likely benign variants, variants of uncertain significance, or inconclusive variants identified during this analysis may be reported in certain circumstances. Our laboratory's variant classification criteria are based on the ACMG and internal guidelines and our current understanding of the specific genes. This interpretation may change over time as more information about a gene and/or variant becomes available. Natera and its lab partner(s) may reclassify variants at certain intervals but may not release updated reports without a specific request made to Natera by the ordering provider. Natera may disclose incidental findings if deemed clinically pertinent to the test performed.

Negative Results

A negative carrier screening result reduces the risk for a patient to be a carrier of a specific disease but does not completely rule out carrier status. Please visit <https://www.natera.com/panel-option/h-all/> for a table of carrier rates, detection rates, residual risks and promised variants/exons per gene. Carrier rates before and after testing vary by ethnicity and assume a negative family history for each disease screened and the absence of clinical symptoms in the patient. Any patient with a family history for a specific genetic disease will have a higher carrier risk prior to testing and, if the disease-causing mutation in their family is not included on the test, their carrier risk would remain unchanged. Genetic counseling is recommended for patients with a family history of genetic disease so that risk figures based on actual family history can be determined and discussed along with potential implications for reproduction. Horizon carrier screening has been developed to identify the reproductive risks for monogenic inherited conditions. Even when one or both members of a couple screen negative for pathogenic variants in a specific gene, the disease risk for their offspring is not zero. There is still a low risk for the condition in their offspring due to a number of different mechanisms that are not detected by Horizon including, but not limited to, pathogenic variant(s) in the tested gene or in a different gene not included on Horizon, pathogenic variant(s) in an upstream regulator, uniparental disomy, de novo mutation(s), or digenic or polygenic inheritance.

Additional Comments

These analyses generally provide highly accurate information regarding the patient's carrier status. Despite this high level of accuracy, it should be kept in mind that there are many potential sources of diagnostic error, including misidentification of samples, polymorphisms, or other rare genetic variants that interfere with analysis. Families should understand that rare diagnostic errors may occur for these reasons.