



Donor 7173

Genetic Testing Summary

Fairfax Cryobank recommends reviewing this genetic testing summary with your healthcare provider to determine suitability.

Last Updated: 04/30/25

Donor Reported Ancestry: Italian, Chilean

Jewish Ancestry: No

Genetic Test*	Result	Comments/Donor's Residual Risk**
Chromosome analysis (karyotype)	Normal male karyotype	No evidence of clinically significant chromosome abnormalities
Hemoglobin evaluation	Normal hemoglobin fractionation and MCV/MCH results	Reduced risk to be a carrier for sickle cell anemia, beta thalassemia, alpha thalassemia trait (aa/-- and a-/a-) and other hemoglobinopathies
Expanded Genetic Disease Carrier Screening Panel attached- 549 diseases by gene sequencing.	Carrier: Carnitine Deficiency (SLC22A5) Carrier: Very Long Chain Acyl-CoA Dehydrogenase Deficiency (ACADVL) Negative for other genes sequenced.	Partner testing is recommended before using this donor.

*No single test can screen for all genetic disorders. A negative screening result significantly reduces, but cannot eliminate, the risk for these conditions in a pregnancy.

**Donor residual risk is the chance the donor is still a carrier after testing negative.

Patient Information

Patient Name: Donor 7173

Date Of Birth: [REDACTED]

Gender: Male

Ethnicity: Hispanic/Latin American

Patient ID: N/A

Medical Record #: N/A

Collection Kit: [REDACTED]

Accession ID: N/A

Case File ID: [REDACTED]

Test Information

Ordering Physician: [REDACTED]

Clinic Information: Fairfax Cryobank

Phone: [REDACTED]

Report Date: 03/11/2025

Sample Collected: 02/25/2025

Sample Received: 02/26/2025

Sample Type: Blood

**CARRIER SCREENING REPORT**

ABOUT THIS SCREEN: Horizon™ is a carrier screen for specific autosomal recessive and X-linked diseases. This information can help patients learn their risk of having a child with specific genetic conditions.

ORDER SELECTED: The Horizon Custom panel was ordered for this patient. Males are not screened for X-linked diseases

FINAL RESULTS SUMMARY:**CARRIER for Carnitine Deficiency**

Positive for the likely pathogenic variant c.1354G>A (p.E452K) in the SLC22A5 gene. If this individual's partner is a carrier for CARNITINE DEFICIENCY, their chance to have a child with this condition may be as high as 1 in 4 (25%). Carrier screening for this individual's partner is suggested.

CARRIER for VERY LONG-CHAIN ACYL-CoA DEHYDROGENASE DEFICIENCY

Positive for the likely pathogenic variant c.947G>A (p.R316Q) in the ACADVL gene. This variant has been reported in a homozygous state or in conjunction with another variant in individual(s) with positive newborn screening (PMID: 29519241, internal data). If this individual's partner is a carrier for VERY LONG-CHAIN ACYL-CoA DEHYDROGENASE DEFICIENCY, their chance to have a child with this condition may be as high as 1 in 4 (25%). Carrier screening for this individual's partner is suggested.

Negative for 547 out of 549 diseases

No other pathogenic variants were detected in the genes that were screened. The patient's remaining carrier risk after the negative screening results is listed for each disease/gene on the Horizon website at <https://www.natera.com/panel-option/h-all/>. Please see the following pages of this report for a comprehensive list of all conditions included on this individual's screen.

Carrier screening is not diagnostic and may not detect all possible pathogenic variants in a given gene.

RECOMMENDATIONS

Individuals who would like to review their Horizon report with a Natera Laboratory Genetic Counselor may schedule a telephone genetic information session by calling 650-249-9090 or visiting naterasession.com. Clinicians with questions may contact Natera at 650-249-9090 or email support@natera.com. Individuals with positive results may wish to discuss these results with family members to allow them the option to be screened. Comprehensive genetic counseling to discuss the implications of these test results and possible associated reproductive risk is recommended.

Christine M. Eng, M.D.
Medical Director, Baylor Genetics

J. Dianne Keen-Kim, Ph.D., FACMG
Senior Laboratory Director, Natera

Linyan Meng, Ph.D.
Laboratory Director, Baylor Genetics

Yang Wang, Ph.D., FACMG
Laboratory Director, Natera

Patient Information

Patient Name: Donor 7173

Test Information

Ordering Physician: [REDACTED]



Date Of Birth: [REDACTED]

Case File ID: [REDACTED]

Clinic Information: Fairfax Cryobank

Report Date: 03/11/2025

CARNITINE DEFICIENCY**Understanding Your Horizon Carrier Screen Results****What is Carnitine Deficiency?**

Carnitine Deficiency (also called Carnitine Uptake Defect, Primary Carnitine Deficiency, or Carnitine Transporter Deficiency) is an inherited disorder in which certain fats cannot be broken down and used for energy because the body cannot process carnitine. Carnitine is a substance that is found in food and helps the body turn fat into energy. Signs and symptoms of Carnitine Deficiency may begin shortly after birth or in childhood. People with this condition may have low blood sugar (hypoglycemia), lack of energy, poor appetite, breathing problems, vomiting, diarrhea, and confusion. Symptoms often appear during an illness or after going a long time without food, and can be life-threatening if not treated. Children with this disorder who do not receive treatment may develop an enlarged heart, muscle weakness, and liver disease. Some people with Carnitine Deficiency never have symptoms of this condition. Treatment with carnitine can help prevent or reverse the signs and symptoms of Carnitine Deficiency. Children with this disorder who receive treatment can have healthy growth and development. Clinical trials involving potential new treatments for this condition may be available (see www.clinicaltrials.gov).

What causes Carnitine Deficiency?

Carnitine Deficiency is caused by a gene change, or mutation, in both copies of the SLC22A5 gene pair. These mutations cause the genes to not work properly or not work at all. When both copies of the SLC22A5 gene do not work correctly, it leads to the symptoms described above. Carnitine Deficiency is inherited in an autosomal recessive manner. This means that, in most cases, both parents must be carriers of a mutation in one copy of the SLC22A5 gene to have a child with Carnitine Deficiency. People who are carriers for Carnitine Deficiency are usually healthy and do not typically have symptoms nor do they have Carnitine Deficiency themselves. Usually a child inherits two copies of each gene, one copy from the mother and one copy from the father. If the mother and father are both carriers for Carnitine Deficiency there is a 1 in 4, or 25%, chance in each pregnancy for both partners to pass on their SLC22A5 gene mutations to the child, who will then have Carnitine Deficiency. Individuals found to carry more than one mutation for Carnitine Deficiency should discuss their risk for having an affected child, and any potential effects to their own health, with their health care provider.

What can I do next?

You may wish to speak with a local genetic counselor about your carrier test results. A genetic counselor in your area can be located on the National Society of Genetic Counselors website (www.nsgc.org). Your siblings and other relatives are at increased risk to also have this mutation. You are encouraged to inform your family members of your test results as they may wish to consider being tested themselves. If you are pregnant, your partner can have carrier screening for Carnitine Deficiency ordered by a health care professional. If your partner is not found to be a carrier for Carnitine Deficiency, your risk of having a child with Carnitine Deficiency is greatly reduced. Although not requested routinely, couples at risk of having a baby with Carnitine Deficiency can opt to have prenatal diagnosis done through chorionic villus sampling (CVS) or amniocentesis during pregnancy to test the fetus for Carnitine Deficiency. The baby can also be tested after birth for this condition. Although Carnitine Deficiency is screened for as part of the Newborn Screening program in some US states, babies at 25% for this condition may need diagnostic testing in addition to newborn screening. If you are not yet pregnant, your partner can have carrier screening for Carnitine Deficiency ordered by a health care professional. If your partner is found to be a carrier for Carnitine Deficiency you have several reproductive options to consider:

- Natural pregnancy with or without prenatal diagnosis of the fetus or testing the baby after birth for Carnitine Deficiency
- Preimplantation genetic diagnosis (PGD) with in vitro fertilization (IVF) to test embryos for Carnitine Deficiency
- Adoption or use of a sperm or egg donor who is not a carrier for Carnitine Deficiency Please note: prenatal diagnosis, PGD, and use of sperm or egg donors are not routinely requested because Carnitine Deficiency is considered a treatable condition.

What resources are available?

- Genetics Home Reference: <http://ghr.nlm.nih.gov/condition/primary-carnitine-deficiency>
- Muscular Dystrophy Association: <https://www.mda.org/disease/metabolic-diseases-of-muscle/carnitine-deficiency>
- Prenatal diagnosis done through CVS: <http://www.marchofdimes.org/chorionic-villus-sampling.aspx>
- Prenatal diagnosis done through Amniocentesis: <http://www.marchofdimes.org/amniocentesis.aspx>
- PGD with IVF: <http://www.natera.com/spectrum>

Patient Information

Patient Name: [REDACTED]

Test Information

Ordering Physician: [REDACTED]



Clinic Information: [REDACTED]

Date Of Birth: [REDACTED]

Case File ID: [REDACTED]

Report Date: [REDACTED]

VERY LONG-CHAIN ACYL-CoA DEHYDROGENASE DEFICIENCY**Understanding Your Horizon Carrier Screen Results****What is Very Long-Chain Acyl-CoA Dehydrogenase Deficiency?**

Very Long-Chain Acyl-CoA Dehydrogenase Deficiency (VLCAD Deficiency) is an inherited disorder that prevents the body from breaking down certain types of fat into energy, particularly during periods of fasting, illness, or exercise. Signs and symptoms can begin anytime between infancy and adulthood. Infants with the most severe form of VLCAD Deficiency develop symptoms in the first few months of life. VLCAD Deficiency causes a thickening of the heart muscle (cardiomyopathy) which causes the heart not to work properly. It can also cause an abnormal heart rhythm and/or fluid around the heart. Infants with VLCAD Deficiency can have poor muscle tone, lack of energy, an enlarged liver, and periods of low blood sugar (hypoglycemia). If not treated, affected infants can die. With early diagnosis and lifelong treatment, infants with VLCAD Deficiency can survive and may have healthy growth and development. Affected individuals who develop symptoms later in childhood may not have heart disease. People with the childhood-onset form typically have low blood sugar, an enlarged liver, and muscle weakness, especially after exercise. Some individuals with VLCAD Deficiency have signs and symptoms that do not begin until adulthood. This milder form typically does not affect the heart and may or may not cause low blood sugar. Individuals with VLCAD Deficiency that begins in adulthood may have muscle cramps and pain, often following exercise. If untreated, the muscles in the body start breaking down and kidney damage can occur. With careful treatment, people with the childhood and adult forms of VLCAD Deficiency can live healthy lives with typical growth and development. Clinical trials involving potential new treatments for this condition may be available (see www.clinicaltrials.gov).

What causes Very Long-Chain Acyl-CoA Dehydrogenase Deficiency?

VLCAD Deficiency is caused by a gene change, or mutation, in both copies of the ACADVL gene pair. These mutations cause the genes to not work properly or not work at all. The job of the ACADVL genes is to help break down certain types of fat in the body to use for energy. When both copies of this gene do not work correctly, the body cannot break down these fats and they build up, leading to the symptoms described above. VLCAD Deficiency is inherited in an autosomal recessive manner. This means that, in most cases, both parents must be carriers of a mutation in one copy of the ACADVL gene to have a child with this condition. People who are carriers for VLCAD Deficiency are usually healthy and do not have symptoms nor do they have VLCAD Deficiency themselves. Usually a child inherits two copies of each gene, one copy from the mother and one copy from the father. If the mother and father are both carriers for VLCAD Deficiency, there is a 1 in 4, or 25%, chance in each pregnancy for both partners to pass on their ACADVL gene mutations to the child, who will then have this condition. Women who have a pregnancy in which the fetus has VLCAD Deficiency are also at risk for pregnancy complications such as HELLP syndrome (Hemolysis, Elevated Liver enzymes and Low Platelets) and AFLP (Acute Fatty Liver of Pregnancy). Individuals found to carry more than one mutation for VLCAD Deficiency should discuss any potential risks to their own health and their risk for having an affected child with their health care provider.

What can I do next?

You may wish to speak with a local genetic counselor about your carrier test results. A genetic counselor in your area can be located on the National Society of Genetic Counselors website (www.nsgc.org). Your siblings and other relatives are at increased risk to also have this mutation. You are encouraged to inform your family members of your test results as they may wish to consider being tested themselves. If you are pregnant, your partner can have carrier screening for VLCAD Deficiency ordered by a health care professional. If your partner is not found to be a carrier for this condition your risk of having a child with VLCAD Deficiency is greatly reduced. Couples at risk of having a baby with VLCAD Deficiency can opt to have prenatal diagnosis done through chorionic villus sampling (CVS) or amniocentesis during pregnancy or can choose to have the baby tested after birth for this condition. Although VLCAD Deficiency is screened for as part of the newborn screening program in all U.S. states, babies at 25% risk for this condition may need diagnostic testing in addition to newborn screening. If you are pregnant with a fetus that has VLCAD Deficiency, you should discuss pregnancy management options for women at risk for HELLP syndrome/AFLP with your health care providers. If you are not yet pregnant, your partner can have carrier screening for VLCAD Deficiency ordered by a health care professional. If your partner is found to be a carrier for VLCAD Deficiency, you have several reproductive options to consider:

- Natural pregnancy with or without prenatal diagnostic testing of the fetus or testing the baby after birth for VLCAD Deficiency
- Preimplantation genetic diagnosis (PGD) with in vitro fertilization (IVF) to test embryos for VLCAD Deficiency
- Adoption or use of a sperm or egg donor who is not a carrier for VLCAD Deficiency

What resources are available?

- Baby's First Test "Very Long-Chain Acyl-CoA Dehydrogenase Deficiency": <http://www.babysfirsttest.org/newborn-screening/conditions/very-long-chain-acyl-coa-dehydrogenase-deficiency>
- Prenatal diagnosis by CVS: <http://www.marchofdimes.org/chorionic-villus-sampling.aspx>
- Prenatal diagnosis by amniocentesis: <http://www.marchofdimes.org/amniocentesis.aspx>
- Preimplantation genetic diagnosis (PGD) with IVF: <http://www.natera.com/spectrum>

Patient Information

Patient Name: [REDACTED]

Test Information

Ordering Physician: [REDACTED]



Date Of Birth: [REDACTED]

Case File ID: [REDACTED]

Clinic Information:

Report Date:

VARIANT DETAILS**ACADVL, c.947G>A (p.R316Q), likely pathogenic**

- The c.947G>A (p.R316Q) variant in the ACADVL gene has been observed at a frequency of 0.0032% in the gnomAD v2.1.1 dataset.
- This variant has been reported in a homozygous state or in conjunction with another variant in individual(s) with positive newborn screening (PMID: 29519241, internal data).
- This variant has been reported in ClinVar [ID: 203577].

SLC22A5, c.1354G>A (p.E452K), likely pathogenic

- The c.1354G>A (p.E452K) variant in the SLC22A5 gene has been observed at a frequency of 0.0028% in the gnomAD v2.1.1 dataset.
- This variant has been reported in a homozygous state or in conjunction with another variant in individual(s) with carnitine deficiency, systemic primary (PMID: 10679939).
- Functional studies demonstrated that this variant causes reduced transport activity (PMID: 10679939).
- This variant has been reported in ClinVar [ID: 25421].

Patient Information

Patient Name:

Test Information

Ordering Physician:



Clinic Information:

Date Of Birth:



Case File ID:



Report Date:

DISEASES SCREENED

Below is a list of all diseases screened and the result. Certain conditions have unique patient-specific numerical values, therefore, results for those conditions are formatted differently.

Autosomal Recessive**1**17-BETA HYDROXYSTEROID DEHYDROGENASE 3 DEFICIENCY (*HSD17B3*) **negative****3**

3-BETA-HYDROXYSTEROID DEHYDROGENASE TYPE II DEFICIENCY (*HSD3B2*) **negative**
3-HYDROXY-3-METHYLGLUTARYL-COENZYME A LYASE DEFICIENCY (*HMGCL*) **negative**
3-HYDROXYACYL-COA DEHYDROGENASE DEFICIENCY (*HADH*) **negative**
3-METHYLCROTONYL-CoA CARBOXYLASE 2 DEFICIENCY (*MCCC2*) **negative**
3-PHOSPHOGLYCERATE DEHYDROGENASE DEFICIENCY (*PHGDH*) **negative**

55-ALPHA-REDUCTASE DEFICIENCY (*SRD5A2*) **negative****6**6-PYRUVYL-TETRAHYDROPTERIN SYNTHASE (*PTPS*) DEFICIENCY (*PTS*) **negative****A**

ABCA4-RELATED CONDITIONS (*ABCA4*) **negative**
ABETALIPOPROTEINEMIA (*MTTP*) **negative**
ACHONDROGENESIS, TYPE 1B (*SLC26A2*) **negative**
ACHROMATOPSIA, CNGB3-RELATED (*CNGB3*) **negative**
ACRODERMATITIS ENTEROPATHICA (*SLC39A4*) **negative**
ACTION MYOCLONUS-RENAL FAILURE (AMRF) SYNDROME (*SCARB2*) **negative**
ACUTE INFANTILE LIVER FAILURE, TRMU-RELATED (*TRMU*) **negative**
ACYL-COA OXIDASE I DEFICIENCY (*ACOX1*) **negative**
AICARDI-GOUTIERES SYNDROME (*SAMHD1*) **negative**
AICARDI-GOUTIERES SYNDROME, RNASEH2A-RELATED (*RNASEH2A*) **negative**
AICARDI-GOUTIERES SYNDROME, RNASEH2B-RELATED (*RNASEH2B*) **negative**
AICARDI-GOUTIERES SYNDROME, RNASEH2C-RELATED (*RNASEH2C*) **negative**
AICARDI-GOUTIERES SYNDROME, TREX1-RELATED (*TREX1*) **negative**
ALPHA-MANNOSIDOSIS (*MAN2B1*) **negative**
ALPHA-THALASSEMIA (*HBA1/HBA2*) **negative**
ALPORT SYNDROME, COL4A3-RELATED (*COL4A3*) **negative**
ALPORT SYNDROME, COL4A4-RELATED (*COL4A4*) **negative**
ALSTROM SYNDROME (*ALMS1*) **negative**
AMISH INFANTILE EPILEPSY SYNDROME (*ST3GAL5*) **negative**
ANDERMANN SYNDROME (*SLC12A6*) **negative**
ARGININE:GLYCINE AMIDINOTRANSFERASE DEFICIENCY (AGAT DEFICIENCY) (*GATM*) **negative**
ARGININEMIA (*ARG1*) **negative**
ARGININOSUCCINATE LYASE DEFICIENCY (*ASL*) **negative**
AROMATASE DEFICIENCY (*CYP19A1*) **negative**
ASPARAGINE SYNTHETASE DEFICIENCY (*ASNS*) **negative**
ASPARTYLGLYCOSAMINURIA (AGA) **negative**
ATAXIA WITH VITAMIN E DEFICIENCY (*TTPA*) **negative**
ATAXIA-TELANGIECTASIA (*ATM*) **negative**
ATAXIA-TELANGIECTASIA-LIKE DISORDER 1 (*MRE11*) **negative**
ATRAANSFERRINEMIA (*TF*) **negative**
AUTISM SPECTRUM, EPILEPSY AND ARTHROGRYPOSIS (*SLC35A3*) **negative**
AUTOIMMUNE POLYGLANDULAR SYNDROME, TYPE 1 (*AIRE*) **negative**
AUTOSOMAL RECESSIVE CONGENITAL ICHTHYOSIS (ARCI), SLC27A4-RELATED (*SLC27A4*) **negative**
AUTOSOMAL RECESSIVE SPASTIC ATAXIA OF CHARLEVOIX-SAGUENAY (SACS) **negative**

B

BARDET-BIEDL SYNDROME, ARL6-RELATED (*ARL6*) **negative**
BARDET-BIEDL SYNDROME, BBS10-RELATED (*BBS10*) **negative**
BARDET-BIEDL SYNDROME, BBS12-RELATED (*BBS12*) **negative**
BARDET-BIEDL SYNDROME, BBS1-RELATED (*BBS1*) **negative**
BARDET-BIEDL SYNDROME, BBS2-RELATED (*BBS2*) **negative**
BARDET-BIEDL SYNDROME, BBS4-RELATED (*BBS4*) **negative**
BARDET-BIEDL SYNDROME, BBS5-RELATED (*BBS5*) **negative**
BARDET-BIEDL SYNDROME, BBS7-RELATED (*BBS7*) **negative**
BARDET-BIEDL SYNDROME, BBS9-RELATED (*BBS9*) **negative**
BARDET-BIEDL SYNDROME, TTC8-RELATED (*TTC8*) **negative**
BARE LYMPHOCYTE SYNDROME, CIITA-RELATED (*CIITA*) **negative**
BARTTER SYNDROME, BSND-RELATED (*BSND*) **negative**
BARTTER SYNDROME, KCNJ1-RELATED (*KCNJ1*) **negative**
BARTTER SYNDROME, SLC12A1-RELATED (*SLC12A1*) **negative**
BATTEN DISEASE, CLN3-RELATED (*CLN3*) **negative**
BETA-HEMOGLOBINOPATHIES (*HBB*) **negative**
BETA-KETOTHIOLASE DEFICIENCY (*ACAT1*) **negative**
BETA-MANNOSIDOSIS (*MANBA*) **negative**
BETA-UREIDOPROPIONASE DEFICIENCY (*UPB1*) **negative**
BILATERAL FRONTOPIRIETAL POLYMICROGYRIA (*GPR56*) **negative**

BIOTINIDASE DEFICIENCY (*BTD*) **negative**BIOTIN-THIAMINE-RESPONSIVE BASAL GANGLIA DISEASE (BTBGD) (*SLC19A3*) **negative**BLOOM SYNDROME (*BLM*) **negative**BRITTLE CORNEA SYNDROME 1 (*ZNF469*) **negative**BRITTLE CORNEA SYNDROME 2 (*PRDM5*) **negative****C**

CANAVAN DISEASE (*ASPA*) **negative**
CARBAMOYL PHOSPHATE SYNTHETASE I DEFICIENCY (*CPS1*) **negative**
CARNITINE DEFICIENCY (*SLC22A5*) **see first page**
CARNITINE PALMITOYLTRANSFERASE IA DEFICIENCY (*CPT1A*) **negative**
CARNITINE PALMITOYLTRANSFERASE II DEFICIENCY (*CPT2*) **negative**
CARNITINE-ACYLCARNITINE TRANSLOCASE DEFICIENCY (*SLC25A20*) **negative**
CARPENTER SYNDROME (*RAB23*) **negative**
CARTILAGE-HAIR HYPOPLASIA (*RMRP*) **negative**
CATECHOLAMINERGIC POLYMORPHIC VENTRICULAR TACHYCARDIA (*CASQ2*) **negative**
CD59-MEDIATED HEMOLYTIC ANEMIA (*CD59*) **negative**
CEP152-RELATED MICROCEPHALY (*CEP152*) **negative**
CEREBRAL DYSGENESIS, NEUROPATHY, ICHTHYOSIS, AND PALMOPLANTAR KERATODERMA (CEDNIK) SYNDROME (*SNAP29*) **negative**
CEREBROTENDINOUS XANTHOMATOSIS (*CYP27A1*) **negative**
CHARCOT-MARIE-TOOTH DISEASE, RECESSIVE INTERMEDIATE C (*PLEKHG5*) **negative**
CHARCOT-MARIE-TOOTH-DISEASE, TYPE 4D (*NDRG1*) **negative**
CHEDIAK-HIGASHI SYNDROME (*LYST*) **negative**
CHOREOACANTHOCYTOSIS (*VPS13A*) **negative**
CHRONIC GRANULOMATOUS DISEASE, CYBA-RELATED (*CYBA*) **negative**
CHRONIC GRANULOMATOUS DISEASE, NCF2-RELATED (*NCF2*) **negative**
CILIOPATHIES, RPGRIP1L-RELATED (*RPGRIP1L*) **negative**
CITRIN DEFICIENCY (*SLC25A13*) **negative**
CITRULLINEMIA, TYPE 1 (*ASS1*) **negative**
CLN10 DISEASE (*CTSD*) **negative**
COHEN SYNDROME (*VPS13B*) **negative**
COL11A2-RELATED CONDITIONS (*COL11A2*) **negative**
COMBINED MALONIC AND METHYLMALONIC ACIDURIA (*ACSF3*) **negative**
COMBINED OXIDATIVE PHOSPHORYLATION DEFICIENCY 1 (*GFM1*) **negative**
COMBINED OXIDATIVE PHOSPHORYLATION DEFICIENCY 3 (*TSFM*) **negative**
COMBINED PITUITARY HORMONE DEFICIENCY 1 (*POU1F1*) **negative**
COMBINED PITUITARY HORMONE DEFICIENCY-2 (*PROP1*) **negative**
CONGENITAL ADRENAL HYPERPLASIA, 11-BETA-HYDROXYLASE DEFICIENCY (*CYP11B1*) **negative**
CONGENITAL ADRENAL HYPERPLASIA, 17-ALPHA-HYDROXYLASE DEFICIENCY (*CYP17A1*) **negative**
CONGENITAL ADRENAL HYPERPLASIA, 21-HYDROXYLASE DEFICIENCY (*CYP21A2*) **negative**
CONGENITAL ADRENAL INSUFFICIENCY, CYP11A1-RELATED (*CYP11A1*) **negative**
CONGENITAL AMEGAKARYOCYTIC THROMBOCYTOPENIA (*MPL*) **negative**
CONGENITAL CHRONIC DIARRHEA (*DGAT1*) **negative**
CONGENITAL DISORDER OF GLYCOSYLATION TYPE 1, ALG1-RELATED (*ALG1*) **negative**
CONGENITAL DISORDER OF GLYCOSYLATION, TYPE 1A, PMM2-Related (*PMM2*) **negative**
CONGENITAL DISORDER OF GLYCOSYLATION, TYPE 1B (*MPL*) **negative**
CONGENITAL DISORDER OF GLYCOSYLATION, TYPE 1C (*ALG6*) **negative**
CONGENITAL DYSERYTHROPOIETIC ANEMIA TYPE 2 (*SEC23B*) **negative**
CONGENITAL FINNISH NEPHROSIS (*NPHS1*) **negative**
CONGENITAL HYDROCEPHALUS 1 (*CCDC88C*) **negative**
CONGENITAL HYPERINSULINISM, KCNJ11-Related (*KCNJ11*) **negative**
CONGENITAL INSENSITIVITY TO PAIN WITH ANHIDROSIS (CIPA) (*NTRK1*) **negative**
CONGENITAL MYASTHENIC SYNDROME, CHAT-RELATED (*CHAT*) **negative**
CONGENITAL MYASTHENIC SYNDROME, CHRNE-RELATED (*CHRNE*) **negative**
CONGENITAL MYASTHENIC SYNDROME, COLQ-RELATED (*COLQ*) **negative**
CONGENITAL MYASTHENIC SYNDROME, DOK7-RELATED (*DOK7*) **negative**
CONGENITAL MYASTHENIC SYNDROME, RAPSN-RELATED (*RAPSN*) **negative**
CONGENITAL NEPHROTIC SYNDROME, PLCE1-RELATED (*PLCE1*) **negative**
CONGENITAL NEUTROPENIA, G6PC3-RELATED (*G6PC3*) **negative**
CONGENITAL NEUTROPENIA, HAX1-RELATED (*HAX1*) **negative**
CONGENITAL NEUTROPENIA, VPS45-RELATED (*VPS45*) **negative**
CONGENITAL SECRETORY CHLORIDE DIARRHEA 1 (*SLC26A3*) **negative**
CORNEAL DYSTROPHY AND PERCEPTIVE DEAFNESS (*SLC4A11*) **negative**
CORTICOSTERONE METHYLOXIDASE DEFICIENCY (*CYP11B2*) **negative**
COSTEFF SYNDROME (3-METHYLGLOUTACONIC ACIDURIA, TYPE 3) (*OPA3*) **negative**
CRB1-RELATED RETINAL DYSTROPHIES (*CRB1*) **negative**
CYSTIC FIBROSIS (*CFTR*) **negative**
CYSTINOSIS (*CTNS*) **negative**
CYTOCHROME C OXIDASE DEFICIENCY, PET100-RELATED (*PET100*) **negative**
CYTOCHROME P450 OXIDOREDUCTASE DEFICIENCY (*POR*) **negative**

DD-BIFUNCTIONAL PROTEIN DEFICIENCY (*HSD17B4*) **negative**

Patient Information

Patient Name:

Test Information

Ordering Physician:



Clinic Information:

Date Of Birth:



Case File ID:



Report Date:

D

DEAFNESS, AUTOSOMAL RECESSIVE 77 (LOXHD1) **negative**
DIHYDROPTERIDINE REDUCTASE (DHPR) DEFICIENCY (QDPR) **negative**
DONNAI-BARROW SYNDROME (LRP2) **negative**
DUBIN-JOHNSON SYNDROME (ABCC2) **negative**
DYSKERATOSIS CONGENITA SPECTRUM DISORDERS (TERT) **negative**
DYSKERATOSIS CONGENITA, RTKL1-RELATED (RTKL1) **negative**
DYSTROPHIC EPIDERMOLYSIS BULLOSA, COL7A1-Related (COL7A1) **negative**

E

EARLY INFANTILE EPILEPTIC ENCEPHALOPATHY, CAD-RELATED (CAD) **negative**
EHLERS-DANLOS SYNDROME TYPE VI (PLOD1) **negative**
EHLERS-DANLOS SYNDROME, CLASSIC-LIKE, TNXB-RELATED (TNXB) **negative**
EHLERS-DANLOS SYNDROME, TYPE VII C (ADAMTS2) **negative**
ELLIS-VAN CREVELD SYNDROME, EVC2-RELATED (EVC2) **negative**
ELLIS-VAN CREVELD SYNDROME, EVC-RELATED (EVC) **negative**
ENHANCED S-CONE SYNDROME (NR2E3) **negative**
EPIMERASE DEFICIENCY (GALACTOSEMIA TYPE III) (GALE) **negative**
EPIPHYSEAL DYSPLASIA, MULTIPLE, 7/DESBUQUOIS DYSPLASIA 1 (CANT1) **negative**
ERCC6-RELATED DISORDERS (ERCC6) **negative**
ERCC8-RELATED DISORDERS (ERCC8) **negative**
ETHYLMALONIC ENCEPHALOPATHY (ETHE1) **negative**

F

FACTOR XI DEFICIENCY (F11) **negative**
FAMILIAL DYSAUTONOMIA (IKBKAP) **negative**
FAMILIAL HEMOPHAGOCYTIC LYMPHOHISTIOCYTOSIS, PRF1-RELATED (PRF1) **negative**
FAMILIAL HEMOPHAGOCYTIC LYMPHOHISTIOCYTOSIS, STX11-RELATED (STX11) **negative**
FAMILIAL HEMOPHAGOCYTIC LYMPHOHISTIOCYTOSIS, STXBP2-RELATED (STXBP2) **negative**
FAMILIAL HEMOPHAGOCYTIC LYMPHOHISTIOCYTOSIS, UNC13D-RELATED (UNC13D) **negative**
FAMILIAL HYPERCHOLESTEROLEMIA, LDLRAP1-RELATED (LDLRAP1) **negative**
FAMILIAL HYPERCHOLESTEROLEMIA, LDLR-RELATED (LDLR) **negative**
FAMILIAL HYPERINSULINISM, ABC8-RELATED (ABCC8) **negative**
FAMILIAL NEPHROGENIC DIABETES INSIPIDUS, AQP2-RELATED (AQP2) **negative**
FANCONI ANEMIA, GROUP A (FANCA) **negative**
FANCONI ANEMIA, GROUP C (FANCC) **negative**
FANCONI ANEMIA, GROUP D2 (FANCD2) **negative**
FANCONI ANEMIA, GROUP E (FANCE) **negative**
FANCONI ANEMIA, GROUP F (FANCF) **negative**
FANCONI ANEMIA, GROUP G (FANGG) **negative**
FANCONI ANEMIA, GROUP I (FANCI) **negative**
FANCONI ANEMIA, GROUP J (BRIP1) **negative**
FANCONI ANEMIA, GROUP L (FANCL) **negative**
FARBER LIPOGRANULOMATOSIS (ASAH1) **negative**
FOVEAL HYPOPLASIA (SLC38A8) **negative**
FRASER SYNDROME 3, GRIP1-RELATED (GRIP1) **negative**
FRASER SYNDROME, FRAS1-RELATED (FRAS1) **negative**
FRASER SYNDROME, FREM2-RELATED (FREM2) **negative**
FRIEDREICH ATAXIA (FXN) **negative**
FRUCTOSE-1,6-BISPHOSPHATASE DEFICIENCY (FBP1) **negative**
FUCOSIDOSIS, FUCA1-RELATED (FUCA1) **negative**
FUMARASE DEFICIENCY (FH) **negative**

G

GABA-TRANSAMINASE DEFICIENCY (ABAT) **negative**
GALACTOKINASE DEFICIENCY (GALACTOSEMIA, TYPE II) (GALK1) **negative**
GALACTOSEMIA (GALT) **negative**
GALACTOSIALIDOSIS (CTSA) **negative**
GAUCHER DISEASE (GBA) **negative**
GCH1-RELATED CONDITIONS (GCH1) **negative**
GDF5-RELATED CONDITIONS (GDF5) **negative**
GERODERMA OSTEODYSPLASTICA (GORAB) **negative**
GITELMAN SYNDROME (SLC12A3) **negative**
GLANZMANN THROMBASTHENIA (ITGB3) **negative**
GLUTARIC ACIDEMIA, TYPE 1 (GCDH) **negative**
GLUTARIC ACIDEMIA, TYPE 2A (ETFA) **negative**
GLUTARIC ACIDEMIA, TYPE 2B (ETFB) **negative**
GLUTARIC ACIDEMIA, TYPE 2C (ETFDH) **negative**
GLUTATHIONE SYNTHETASE DEFICIENCY (GSS) **negative**
GLYCINE ENCEPHALOPATHY, AMT-RELATED (AMT) **negative**
GLYCINE ENCEPHALOPATHY, GLDC-RELATED (GLDC) **negative**
GLYCOGEN STORAGE DISEASE TYPE 5 (McArdle Disease) (PYGM) **negative**
GLYCOGEN STORAGE DISEASE TYPE IXB (PHKB) **negative**
GLYCOGEN STORAGE DISEASE TYPE IXC (PHKG2) **negative**
GLYCOGEN STORAGE DISEASE, TYPE 1a (G6PC) **negative**
GLYCOGEN STORAGE DISEASE, TYPE 1b (SLC37A4) **negative**
GLYCOGEN STORAGE DISEASE, TYPE 2 (POMPE DISEASE) (GAA) **negative**
GLYCOGEN STORAGE DISEASE, TYPE 3 (AGL) **negative**
GLYCOGEN STORAGE DISEASE, TYPE 4 (GBE1) **negative**
GLYCOGEN STORAGE DISEASE, TYPE 7 (PFKM) **negative**

GRACILE SYNDROME (BCS1L) **negative**GUANIDINOACETATE METHYLTRANSFERASE DEFICIENCY (GAMT) **negative****H**

HARLEQUIN ICHTHYOSIS (ABCA12) **negative**
HEME OXYGENASE 1 DEFICIENCY (HMOX1) **negative**
HEMOCHROMATOSIS TYPE 2A (HFE2) **negative**
HEMOCHROMATOSIS, TYPE 3, TFR2-Related (TFR2) **negative**
HEPATOCEREBRAL MITOCHONDRIAL DNA DEPLETION SYNDROME, MPV17-RELATED (MPV17) **negative**
HEREDITARY FRUCTOSE INTOLERANCE (ALDOB) **negative**
HEREDITARY HEMOCHROMATOSIS TYPE 2B (HAMP) **negative**
HEREDITARY SPASTIC PARAPARESIS, TYPE 49 (TECPR2) **negative**
HEREDITARY SPASTIC PARAPLEGIA, CYP7B1-RELATED (CYP7B1) **negative**
HERMANSKY-PUDLAK SYNDROME, AP3B1-RELATED (AP3B1) **negative**
HERMANSKY-PUDLAK SYNDROME, BLOC1S3-RELATED (BLOC1S3) **negative**
HERMANSKY-PUDLAK SYNDROME, BLOC1S6-RELATED (BLOC1S6) **negative**
HERMANSKY-PUDLAK SYNDROME, HPS1-RELATED (HPS1) **negative**
HERMANSKY-PUDLAK SYNDROME, HPS3-RELATED (HPS3) **negative**
HERMANSKY-PUDLAK SYNDROME, HPS4-RELATED (HPS4) **negative**
HERMANSKY-PUDLAK SYNDROME, HPS5-RELATED (HPS5) **negative**
HERMANSKY-PUDLAK SYNDROME, HPS6-RELATED (HPS6) **negative**
HOLOCARBOXYLASE SYNTHETASE DEFICIENCY (HLCS) **negative**
HOMOCYSTINURIA AND MEGALOBlastic ANEMIA TYPE CBLG (MTR) **negative**
HOMOCYSTINURIA DUE TO DEFICIENCY OF MTHFR (MTHFR) **negative**
HOMOCYSTINURIA, CBS-RELATED (CBS) **negative**
HOMOCYSTINURIA, Type cblE (MTRR) **negative**
HYDROLETHALUS SYNDROME (HYS1) **negative**
HYPER-IGM IMMUNODEFICIENCY (CD40) **negative**
HYPERORNITHINEMIA-HYPERAMMONEMIA-HOMOCITRULLINURIA (HHH SYNDROME) (SLC25A15) **negative**
HYPERPHOSPHATEMIC FAMILIAL TUMORAL CALCINOSIS, GALNT3-RELATED (GALNT3) **negative**
HYPOMYELINATING LEUKODYSTROPHY 12 (VPS11) **negative**
HYPOPHOSPHATASIA, ALPL-RELATED (ALPL) **negative**

I

IMERSLUND-GRÄSBECK SYNDROME 2 (AMN) **negative**
IMMUNODEFICIENCY-CENTROMERIC INSTABILITY-FACIAL ANOMALIES (ICF) SYNDROME, DNMT3B-RELATED (DNMT3B) **negative**
IMMUNODEFICIENCY-CENTROMERIC INSTABILITY-FACIAL ANOMALIES (ICF) SYNDROME, ZBTB24-RELATED (ZBTB24) **negative**
INCLUSION BODY MYOPATHY 2 (GNE) **negative**
INFANTILE CEREBRAL AND CEREBELLAR ATROPHY (MED17) **negative**
INFANTILE NEPHRONOPHTHISIS (INVS) **negative**
INFANTILE NEUROAXONAL DYSTROPHY (PLA2G6) **negative**
ISOLATED ECTOPIA LENTIS (ADAMTSL4) **negative**
ISOLATED SULFITE OXIDASE DEFICIENCY (SUOX) **negative**
ISOLATED THYROID-STIMULATING HORMONE DEFICIENCY (TSHB) **negative**
ISOVALERIC ACIDEMIA (IVD) **negative**

J

JOHANSON-BLIZZARD SYNDROME (UBR1) **negative**
JOUBERT SYNDROME 2 / MECKEL SYNDROME 2 (TMEM216) **negative**
JOUBERT SYNDROME AND RELATED DISORDERS (JSRD), TMEM67-RELATED (TMEM67) **negative**
JOUBERT SYNDROME, AHI1-RELATED (AHI1) **negative**
JOUBERT SYNDROME, ARL13B-RELATED (ARL13B) **negative**
JOUBERT SYNDROME, B9D1-RELATED (B9D1) **negative**
JOUBERT SYNDROME, B9D2-RELATED (B9D2) **negative**
JOUBERT SYNDROME, C2CD3-RELATED/OROFACIODIGITAL SYNDROME 14 (C2CD3) **negative**
JOUBERT SYNDROME, CC2D2A-RELATED/COACH SYNDROME (CC2D2A) **negative**
JOUBERT SYNDROME, CEP104-RELATED (CEP104) **negative**
JOUBERT SYNDROME, CEP120-RELATED/SHORT-RIB THORACIC DYSPLASIA 13 WITH OR WITHOUT POLYDACTYLY (CEP120) **negative**
JOUBERT SYNDROME, CEP41-RELATED (CEP41) **negative**
JOUBERT SYNDROME, CPLANE1-RELATED / OROFACIODIGITAL SYNDROME 6 (CPLANE1) **negative**
JOUBERT SYNDROME, CSPP1-RELATED (CSPP1) **negative**
JOUBERT SYNDROME, INPP5E-RELATED (INPP5E) **negative**
JUNCTIONAL EPIDERMOLYSIS BULLOSA, COL17A1-RELATED (COL17A1) **negative**
JUNCTIONAL EPIDERMOLYSIS BULLOSA, ITGA6-RELATED (ITGA6) **negative**
JUNCTIONAL EPIDERMOLYSIS BULLOSA, ITGB4-RELATED (ITGB4) **negative**
JUNCTIONAL EPIDERMOLYSIS BULLOSA, LAMB3-RELATED (LAMB3) **negative**
JUNCTIONAL EPIDERMOLYSIS BULLOSA, LAMC2-RELATED (LAMC2) **negative**
JUNCTIONAL EPIDERMOLYSIS BULLOSA/LARYNGOONYCHOCUTANEOUS SYNDROME, LAMA3-RELATED (LAMA3) **negative**

KKRABBE DISEASE (GALC) **negative****L**LAMELLAR ICHTHYOSIS, TYPE 1 (TGM1) **negative**

Patient Information

Patient Name:

Test Information

Ordering Physician:

Clinic Information:

Date Of Birth:

Case File ID:

Report Date:

**L**

LARON SYNDROME (*GHR*) **negative**
LEBER CONGENITAL AMAUROSIS 2 (*RPE65*) **negative**
LEBER CONGENITAL AMAUROSIS TYPE AIP1 (*AIP1*) **negative**
LEBER CONGENITAL AMAUROSIS TYPE GUCY2D (*GUCY2D*) **negative**
LEBER CONGENITAL AMAUROSIS TYPE TULP1 (*TULP1*) **negative**
LEBER CONGENITAL AMAUROSIS, IQCB1-RELATED/SENIOR-LOKEN SYNDROME 5 (*IQCB1*) **negative**
LEBER CONGENITAL AMAUROSIS, TYPE CEP290 (*CEP290*) **negative**
LEBER CONGENITAL AMAUROSIS, TYPE LCA5 (*LCA5*) **negative**
LEBER CONGENITAL AMAUROSIS, TYPE RDH12 (*RDH12*) **negative**
LEIGH SYNDROME, FRENCH-CANADIAN TYPE (*LRPPRC*) **negative**
LETHAL CONGENITAL CONTRACTURE SYNDROME 1 (*GLE1*) **negative**
LEUKOENCEPHALOPATHY WITH VANISHING WHITE MATTER (*EIF2B5*) **negative**
LEUKOENCEPHALOPATHY WITH VANISHING WHITE MATTER, EIF2B1-RELATED (*EIF2B1*) **negative**
LEUKOENCEPHALOPATHY WITH VANISHING WHITE MATTER, EIF2B2-RELATED (*EIF2B2*) **negative**
LEUKOENCEPHALOPATHY WITH VANISHING WHITE MATTER, EIF2B3-RELATED (*EIF2B3*) **negative**
LEUKOENCEPHALOPATHY WITH VANISHING WHITE MATTER, EIF2B4-RELATED (*EIF2B4*) **negative**
LIG4 SYNDROME (*LIG4*) **negative**
LIMB-GIRDLE MUSCULAR DYSTROPHY TYPE 8 (*TRIM32*) **negative**
LIMB-GIRDLE MUSCULAR DYSTROPHY, TYPE 2A (*CAPN3*) **negative**
LIMB-GIRDLE MUSCULAR DYSTROPHY, TYPE 2B (*DYSF*) **negative**
LIMB-GIRDLE MUSCULAR DYSTROPHY, TYPE 2C (*SGCG*) **negative**
LIMB-GIRDLE MUSCULAR DYSTROPHY, TYPE 2D (*SGCA*) **negative**
LIMB-GIRDLE MUSCULAR DYSTROPHY, TYPE 2E (*SGCB*) **negative**
LIMB-GIRDLE MUSCULAR DYSTROPHY, TYPE 2F (*SGCD*) **negative**
LIMB-GIRDLE MUSCULAR DYSTROPHY, TYPE 2I (*FKRP*) **negative**
LIPOAMIDE DEHYDROGENASE DEFICIENCY (DIHYDROLIPOAMIDE DEHYDROGENASE DEFICIENCY) (*DLD*) **negative**
LIPOID ADRENAL HYPERPLASIA (*STAR*) **negative**
LIPOPROTEIN LIPASE DEFICIENCY (*LPL*) **negative**
LONG CHAIN 3-HYDROXYACYL-CoA DEHYDROGENASE DEFICIENCY (*HADHA*) **negative**
LRAT-RELATED CONDITIONS (*LRAT*) **negative**
LUNG DISEASE, IMMUNODEFICIENCY, AND CHROMOSOME BREAKAGE SYNDROME (*LICS*) (*NSMCE3*) **negative**
LYSINURIC PROTEIN INTOLERANCE (*SLC7A7*) **negative**

M

MALONYL-CoA DECARBOXYLASE DEFICIENCY (*MLYCD*) **negative**
MAPLE SYRUP URINE DISEASE, TYPE 1A (*BCKDHA*) **negative**
MAPLE SYRUP URINE DISEASE, TYPE 1B (*BCKDHB*) **negative**
MAPLE SYRUP URINE DISEASE, TYPE 2 (*DBT*) **negative**
MCKUSICK-KAUFMAN SYNDROME (*MKKS*) **negative**
MECKEL SYNDROME 7/NEPHRONOPHTHISIS 3 (*NPHP3*) **negative**
MECKEL-GRUBER SYNDROME, TYPE 1 (*MKS1*) **negative**
MECR-RELATED NEUROLOGIC DISORDER (*MECR*) **negative**
MEDIUM CHAIN ACYL-CoA DEHYDROGENASE DEFICIENCY (*ACADM*) **negative**
MEDNIK SYNDROME (*AP1S1*) **negative**
MEGALENCEPHALIC LEUKOENCEPHALOPATHY WITH SUBCORTICAL CYSTS (*MLC1*) **negative**
MEROSIN-DEFICIENT MUSCULAR DYSTROPHY (*LAMA2*) **negative**
METABOLIC ENCEPHALOPATHY AND ARRHYTHMIAS, TANGO2-RELATED (*TANGO2*) **negative**
METACHROMATIC LEUKODYSTROPHY, ARSA-RELATED (*ARSA*) **negative**
METACHROMATIC LEUKODYSTROPHY, PSAP-RELATED (*PSAP*) **negative**
METHYLMALONIC ACIDEMIA AND HOMOCYSTEINURIA TYPE CBLF (*LMBRD1*) **negative**
METHYLMALONIC ACIDEMIA, MCEE-RELATED (*MCEE*) **negative**
METHYLMALONIC ACIDURIA AND HOMOCYSTEINURIA, TYPE CBLF (*MMACHC*) **negative**
METHYLMALONIC ACIDURIA AND HOMOCYSTEINURIA, TYPE CblD (*MMADHC*) **negative**
METHYLMALONIC ACIDURIA, MMAA-RELATED (*MMAA*) **negative**
METHYLMALONIC ACIDURIA, MMAB-RELATED (*MMAB*) **negative**
METHYLMALONIC ACIDURIA, TYPE MUT (0) (*MUT*) **negative**
MEVALONIC KINASE DEFICIENCY (*MVK*) **negative**
MICROCEPHALIC OSTEODYSPLASTIC PRIMORDIAL DWARFISM TYPE II (*PCNT*) **negative**
MICROPHTHALMIA / ANOPHTHALMIA, VSX2-RELATED (*VSX2*) **negative**
MITOCHONDRIAL COMPLEX 1 DEFICIENCY, ACAD9-RELATED (*ACAD9*) **negative**
MITOCHONDRIAL COMPLEX 1 DEFICIENCY, NDUF5-RELATED (*NDUF5*) **negative**
MITOCHONDRIAL COMPLEX 1 DEFICIENCY, NDUF56-RELATED (*NDUF56*) **negative**
MITOCHONDRIAL COMPLEX I DEFICIENCY, NUCLEAR TYPE 1 (*NDUF54*) **negative**
MITOCHONDRIAL COMPLEX I DEFICIENCY, NUCLEAR TYPE 10 (*NDUF52*) **negative**
MITOCHONDRIAL COMPLEX I DEFICIENCY, NUCLEAR TYPE 17 (*NDUF56*) **negative**
MITOCHONDRIAL COMPLEX I DEFICIENCY, NUCLEAR TYPE 19 (*FOXRED1*) **negative**
MITOCHONDRIAL COMPLEX I DEFICIENCY, NUCLEAR TYPE 3 (*NDUF57*) **negative**
MITOCHONDRIAL COMPLEX I DEFICIENCY, NUCLEAR TYPE 4 (*NDUFV1*) **negative**
MITOCHONDRIAL COMPLEX IV DEFICIENCY, NUCLEAR TYPE 2, SCO2-RELATED (*SCO2*) **negative**
MITOCHONDRIAL COMPLEX IV DEFICIENCY, NUCLEAR TYPE 6 (*COX15*) **negative**
MITOCHONDRIAL DNA DEPLETION SYNDROME 2 (*TK2*) **negative**

MITOCHONDRIAL DNA DEPLETION SYNDROME 3 (*DGUOK*) **negative**
MITOCHONDRIAL MYOPATHY AND SIDEROBLASTIC ANEMIA (MLASA1) (*PUS1*) **negative**
MITOCHONDRIAL TRIFUNCTIONAL PROTEIN DEFICIENCY, HADHB-RELATED (*HADHB*) **negative**
MOLYBDENUM COFACTOR DEFICIENCY TYPE B (*MOCS2*) **negative**
MOLYBDENUM COFACTOR DEFICIENCY, TYPE A (*MOCS1*) **negative**
MUCOLIPIDOSIS II/III A (*GNPTAB*) **negative**
MUCOLIPIDOSIS III GAMMA (*GNPTG*) **negative**
MUCOLIPIDOSIS, TYPE IV (*MCOLN1*) **negative**
MUCOPOLYSACCHARIDOSIS, TYPE I (HURLER SYNDROME) (*IDUA*) **negative**
MUCOPOLYSACCHARIDOSIS, TYPE III A (SANFILIPPO A) (*SGSH*) **negative**
MUCOPOLYSACCHARIDOSIS, TYPE III B (SANFILIPPO B) (*NAGLU*) **negative**
MUCOPOLYSACCHARIDOSIS, TYPE III C (SANFILIPPO C) (*HGSNAT*) **negative**
MUCOPOLYSACCHARIDOSIS, TYPE III D (SANFILIPPO D) (*GNS*) **negative**
MUCOPOLYSACCHARIDOSIS, TYPE IV A (MORQUIO SYNDROME) (*GALNS*) **negative**
MUCOPOLYSACCHARIDOSIS, TYPE IV B/GM1 GANGLIOSIDOSIS (*GLB1*) **negative**
MUCOPOLYSACCHARIDOSIS, TYPE IX (*HYAL1*) **negative**
MUCOPOLYSACCHARIDOSIS, TYPE VI (MAROTEAUX-LAMY) (*ARSB*) **negative**
MUCOPOLYSACCHARIDOSIS, TYPE VII (*GUSB*) **negative**
MULIBREY NANISM (*TRIM37*) **negative**
MULTIPLE PTERYGIUM SYNDROME, CHRNG-RELATED/ESCOBAR SYNDROME (*CHNRG*) **negative**
MULTIPLE SULFATASE DEFICIENCY (*SUMF1*) **negative**
MUSCLE-EYE-BRAIN DISEASE, POMGNT1-RELATED (*POMGNT1*) **negative**
MUSCULAR DYSTROPHY-DYSTROGLYCANOPATHY (*RXYLT1*) **negative**
MUSK-RELATED CONGENITAL MYASTHENIC SYNDROME (*MUSK*) **negative**
MYONEUROGASTROINTESTINAL ENCEPHALOPATHY (MNGIE) (*TYMP*) **negative**
MYOTONIA CONGENITA (*CLCN1*) **negative**

N

N-ACETYLGLUTAMATE SYNTHASE DEFICIENCY (*NAGS*) **negative**
NEMALINE MYOPATHY, NEB-RELATED (*NEB*) **negative**
NEPHRONOPHTHISIS 1 (*NPHP1*) **negative**
NEURONAL CEROID LIPOFUSCINOSIS, CLN5-RELATED (*CLN5*) **negative**
NEURONAL CEROID LIPOFUSCINOSIS, CLN6-RELATED (*CLN6*) **negative**
NEURONAL CEROID LIPOFUSCINOSIS, CLN8-RELATED (*CLN8*) **negative**
NEURONAL CEROID LIPOFUSCINOSIS, MFSD8-RELATED (*MFSD8*) **negative**
NEURONAL CEROID LIPOFUSCINOSIS, PPT1-RELATED (*PPT1*) **negative**
NEURONAL CEROID LIPOFUSCINOSIS, TPP1-RELATED (*TPP1*) **negative**
NGLY1-CONGENITAL DISORDER OF GLYCOSYLATION (*NGLY1*) **negative**
NIEMANN-PICK DISEASE, TYPE C1 / D (*NPC1*) **negative**
NIEMANN-PICK DISEASE, TYPE C2 (*NPC2*) **negative**
NIEMANN-PICK DISEASE, TYPES A / B (*SMPD1*) **negative**
NIJMEGEN BREAKAGE SYNDROME (*NBN*) **negative**
NON-SYNDROMIC HEARING LOSS, GJB2-RELATED (*GJB2*) **negative**
NON-SYNDROMIC HEARING LOSS, MYO15A-RELATED (*MYO15A*) **negative**
NONSYNDROMIC HEARING LOSS, OTOA-RELATED (*OTOA*) **negative**
NONSYNDROMIC HEARING LOSS, OTOF-RELATED (*OTOF*) **negative**
NONSYNDROMIC HEARING LOSS, PJK-RELATED (*PJK*) **negative**
NONSYNDROMIC HEARING LOSS, SYNE4-RELATED (*SYNE4*) **negative**
NONSYNDROMIC HEARING LOSS, TMC1-RELATED (*TMC1*) **negative**
NONSYNDROMIC HEARING LOSS, TMPS3-RELATED (*TMPS3*) **negative**
NONSYNDROMIC INTELLECTUAL DISABILITY (*CC2D1A*) **negative**
NORMOPHOSPHATEMIC TUMORAL CALCINOSIS (*SAMD9*) **negative**

O

OCULOCUTANEOUS ALBINISM TYPE III (*TYRP1*) **negative**
OCULOCUTANEOUS ALBINISM TYPE IV (*SLC45A2*) **negative**
OCULOCUTANEOUS ALBINISM, OCA2-RELATED (*OCA2*) **negative**
OCULOCUTANEOUS ALBINISM, TYPES 1A AND 1B (*TYR*) **negative**
ODONTO-ONYCHO-DERMAL DYSPLASIA / SCHOPF-SCHULZ-PASSARGE SYNDROME (*WNT10A*) **negative**
OMENN SYNDROME, RAG2-RELATED (*RAG2*) **negative**
ORNITHINE AMINOTRANSFERASE DEFICIENCY (*OAT*) **negative**
OSTEOGENESIS IMPERFECTA TYPE VII (*CRTAP*) **negative**
OSTEOGENESIS IMPERFECTA TYPE VIII (*P3H1*) **negative**
OSTEOGENESIS IMPERFECTA TYPE XI (*FKBP10*) **negative**
OSTEOGENESIS IMPERFECTA TYPE XIII (*BMP1*) **negative**
OSTEOPETROSIS, INFANTILE MALIGNANT, TCIRG1-RELATED (*TCIRG1*) **negative**
OSTEOPETROSIS, OSTM1-RELATED (*OSTM1*) **negative**

P

PANTOTHENATE KINASE-ASSOCIATED NEURODEGENERATION (*PANK2*) **negative**
PAPILLON LEFÈVRE SYNDROME (*CTSC*) **negative**
PARKINSON DISEASE 15 (*FBXO7*) **negative**
PENDRED SYNDROME (*SLC26A4*) **negative**
PERLMAN SYNDROME (*DIS3L2*) **negative**
PGM3-CONGENITAL DISORDER OF GLYCOSYLATION (*PGM3*) **negative**
PHENYLKETONURIA (*PAH*) **negative**
PIGN-CONGENITAL DISORDER OF GLYCOSYLATION (*PIGN*) **negative**
PITUITARY HORMONE DEFICIENCY, COMBINED 3 (*LHX3*) **negative**
POLG-RELATED DISORDERS (*POLG*) **negative**

Patient Information

Patient Name:

Test Information

Ordering Physician:



Clinic Information:

Date Of Birth:

Case File ID:

Report Date:

P

POLYCYSTIC KIDNEY DISEASE, AUTOSOMAL RECESSIVE (*PKHD1*) **negative**
 PONTOCEREBELLAR HYPOPLASIA, EXOSC3-RELATED (*EXOSC3*) **negative**
 PONTOCEREBELLAR HYPOPLASIA, RARS2-RELATED (*RARS2*) **negative**
 PONTOCEREBELLAR HYPOPLASIA, TSEN2-RELATED (*TSEN2*) **negative**
 PONTOCEREBELLAR HYPOPLASIA, TSEN54-RELATED (*TSEN54*) **negative**
 PONTOCEREBELLAR HYPOPLASIA, TYPE 1A (*VRK1*) **negative**
 PONTOCEREBELLAR HYPOPLASIA, TYPE 2D (*SEPSECS*) **negative**
 PONTOCEREBELLAR HYPOPLASIA, VPS53-RELATED (*VPS53*) **negative**
 PRIMARY CILIARY DYSKINESIA, CCDC103-RELATED (*CCDC103*) **negative**
 PRIMARY CILIARY DYSKINESIA, CCDC39-RELATED (*CCDC39*) **negative**
 PRIMARY CILIARY DYSKINESIA, DNAH11-RELATED (*DNAH11*) **negative**
 PRIMARY CILIARY DYSKINESIA, DNAH5-RELATED (*DNAH5*) **negative**
 PRIMARY CILIARY DYSKINESIA, DNAI1-RELATED (*DNAI1*) **negative**
 PRIMARY CILIARY DYSKINESIA, DNAI2-RELATED (*DNAI2*) **negative**
 PRIMARY CONGENITAL GLAUCOMA/PETERS ANOMALY (*CYP1B1*) **negative**
 PRIMARY HYPEROXALURIA, TYPE 1 (*AGXT*) **negative**
 PRIMARY HYPEROXALURIA, TYPE 2 (*GRHPR*) **negative**
 PRIMARY HYPEROXALURIA, TYPE 3 (*HOGA1*) **negative**
 PRIMARY MICROCEPHALY 1, AUTOSOMAL RECESSIVE (*MCPH1*) **negative**
 PROGRESSIVE EARLY-ONSET ENCEPHALOPATHY WITH BRAIN ATROPHY AND THIN CORPUS CALLOSUM (*TBCD*) **negative**
 PROGRESSIVE FAMILIAL INTRAHEPATIC CHOLESTASIS, ABCB4-RELATED (*ABCB4*) **negative**
 PROGRESSIVE FAMILIAL INTRAHEPATIC CHOLESTASIS, TYPE 1 (*PFIC1*) (*ATP8B1*) **negative**
 PROGRESSIVE FAMILIAL INTRAHEPATIC CHOLESTASIS, TYPE 2 (*ABCB11*) **negative**
 PROGRESSIVE FAMILIAL INTRAHEPATIC CHOLESTASIS, TYPE 4 (*PFIC4*) (*TJP2*) **negative**
 PROGRESSIVE PSEUDORHEUMATOID DYSPLASIA (*CCN6*) **negative**
 PROLIDASE DEFICIENCY (*PEPD*) **negative**
 PROPIONIC ACIDEMIA, PCCA-RELATED (*PCCA*) **negative**
 PROPIONIC ACIDEMIA, PCCB-RELATED (*PCCB*) **negative**
 PSEUDOXANTHOMA ELASTICUM (*ABCC6*) **negative**
 PTERIN-4 ALPHA-CARBINOLAMINE DEHYDRATASE (*PCD*) DEFICIENCY (*PCBD1*) **negative**
 PYCNODYSTOSIS (*CTSK*) **negative**
 PYRIDOXAL 5'-PHOSPHATE-DEPENDENT EPILEPSY (*PNPO*) **negative**
 PYRIDOXINE-DEPENDENT EPILEPSY (*ALDH7A1*) **negative**
 PYRUVATE CARBOXYLASE DEFICIENCY (*PC*) **negative**
 PYRUVATE DEHYDROGENASE DEFICIENCY, PDHB-RELATED (*PDHB*) **negative**

R

REFSUM DISEASE, PHYH-RELATED (*PHYH*) **negative**
 RENAL TUBULAR ACIDOSIS AND DEAFNESS, ATP6V1B1-RELATED (*ATP6V1B1*) **negative**
 RENAL TUBULAR ACIDOSIS, PROXIMAL, WITH OCULAR ABNORMALITIES AND MENTAL RETARDATION (*SLC4A4*) **negative**
 RETINITIS PIGMENTOSA 25 (*EYS*) **negative**
 RETINITIS PIGMENTOSA 26 (*CERKL*) **negative**
 RETINITIS PIGMENTOSA 28 (*FAM161A*) **negative**
 RETINITIS PIGMENTOSA 36 (*PRCD*) **negative**
 RETINITIS PIGMENTOSA 59 (*DHDDS*) **negative**
 RETINITIS PIGMENTOSA 62 (*MAK*) **negative**
 RHIZOMELIC CHONDRODYSPLASIA PUNCTATA, TYPE 1 (*PEX7*) **negative**
 RHIZOMELIC CHONDRODYSPLASIA PUNCTATA, TYPE 2 (*GNPAT*) **negative**
 RHIZOMELIC CHONDRODYSPLASIA PUNCTATA, TYPE 3 (*AGPS*) **negative**
 RLBP1-RELATED RETINOPATHY (*RLBP1*) **negative**
 ROBERTS SYNDROME (*ESCO2*) **negative**
 RYR1-RELATED CONDITIONS (*RYR1*) **negative**

S

SALLA DISEASE (*SLC17A5*) **negative**
 SANDHOFF DISEASE (*HEXB*) **negative**
 SCHIMKE IMMUNOSKELETAL DYSPLASIA (*SMARCA1*) **negative**
 SCHINDLER DISEASE (*NAGA*) **negative**
 SEGAWA SYNDROME, TH-RELATED (*TH*) **negative**
 SENIOR-LOKEN SYNDROME 4/NEPHRONOPHTHISIS 4 (*NPHP4*) **negative**
 SEPIAPTERIN REDUCTASE DEFICIENCY (*SPR*) **negative**
 SEVERE COMBINED IMMUNODEFICIENCY (*SCID*), CD3D-RELATED (*CD3D*) **negative**
 SEVERE COMBINED IMMUNODEFICIENCY (*SCID*), CD3E-RELATED (*CD3E*) **negative**
 SEVERE COMBINED IMMUNODEFICIENCY (*SCID*), FOXP1-RELATED (*FOXP1*) **negative**
 SEVERE COMBINED IMMUNODEFICIENCY (*SCID*), IKBKB-RELATED (*IKBKB*) **negative**
 SEVERE COMBINED IMMUNODEFICIENCY (*SCID*), IL7R-RELATED (*IL7R*) **negative**
 SEVERE COMBINED IMMUNODEFICIENCY (*SCID*), JAK3-RELATED (*JAK3*) **negative**
 SEVERE COMBINED IMMUNODEFICIENCY (*SCID*), PTPRC-RELATED (*PTPRC*) **negative**
 SEVERE COMBINED IMMUNODEFICIENCY (*SCID*), RAG1-RELATED (*RAG1*) **negative**
 SEVERE COMBINED IMMUNODEFICIENCY, ADA-Related (*ADA*) **negative**
 SEVERE COMBINED IMMUNODEFICIENCY, TYPE ATHABASKAN (*DCLRE1C*) **negative**
 SHORT-RIB THORACIC DYSPLASIA 3 WITH OR WITHOUT POLYDACTYLY (*DYNC2H1*) **negative**
 SHWACHMAN-DIAMOND SYNDROME, SBDS-RELATED (*SBDS*) **negative**
 SIALIDOSIS (*NEU1*) **negative**
 SJÖGREN-LARSSON SYNDROME (*ALDH3A2*) **negative**
 SMITH-LEMLI-OPITZ SYNDROME (*DHCR7*) **negative**
 SPASTIC PARAPLEGIA, TYPE 15 (*ZFYVE26*) **negative**

SPASTIC TETRAPLEGIA, THIN CORPUS CALLOSUM, AND PROGRESSIVE MICROCEPHALY (*SPATCCM*) (*SLC1A4*) **negative**
 SPG11-RELATED CONDITIONS (*SPG11*) **negative**
 SPINAL MUSCULAR ATROPHY (*SMN1*) **negative** **SMN1: Two copies; g.27134T>G: absent; the absence of the g.27134T>G variant decreases the chance to be a silent (2+0) carrier.**
 SPINAL MUSCULAR ATROPHY WITH RESPIRATORY DISTRESS TYPE 1 (*IGHMBP2*) **negative**
 SPINOCEREBELLAR ATAXIA, AUTOSOMAL RECESSIVE 10 (*ANO10*) **negative**
 SPINOCEREBELLAR ATAXIA, AUTOSOMAL RECESSIVE 12 (*WWOX*) **negative**
 SPONDYLOCOSTAL DYSOSTOSIS 1 (*DLL3*) **negative**
 SPONDYLOTHORACIC DYSOSTOSIS, MESP2-Related (*MESP2*) **negative**
 STEEL SYNDROME (*COL27A1*) **negative**
 STEROID-RESISTANT NEPHROTIC SYNDROME (*NPHS2*) **negative**
 STUVE-WIEDEMANN SYNDROME (*LIFR*) **negative**
 SURF1-RELATED CONDITIONS (*SURF1*) **negative**
 SURFACTANT DYSFUNCTION, ABCA3-RELATED (*ABCA3*) **negative**

T

TAY-SACHS DISEASE (*HEXA*) **negative**
 TBCE-RELATED CONDITIONS (*TBCE*) **negative**
 THIAMINE-RESPONSIVE MEGALOBlastic ANEMIA SYNDROME (*SLC19A2*) **negative**
 THYROID DYSHORMONOGENESIS 1 (*SLC5A5*) **negative**
 THYROID DYSHORMONOGENESIS 2A (*TPO*) **negative**
 THYROID DYSHORMONOGENESIS 3 (*TG*) **negative**
 THYROID DYSHORMONOGENESIS 6 (*DUOX2*) **negative**
 TRANSCOBALAMIN II DEFICIENCY (*TCN2*) **negative**
 TRICHOHEPATOENTERIC SYNDROME, SKIC2-RELATED (*SKIC2*) **negative**
 TRICHOHEPATOENTERIC SYNDROME, TTC37-RELATED (*TTC37*) **negative**
 TRICHOHYDROSTROPHY 1/XERODERMA PIGMENTOSUM, GROUP D (*ERCC2*) **negative**
 TRIMETHYLAMINURIA (*FMO3*) **negative**
 TRIPLE A SYNDROME (*AAAS*) **negative**
 TSHR-RELATED CONDITIONS (*TSHR*) **negative**
 TYROSINEMIA TYPE III (*HPD*) **negative**
 TYROSINEMIA, TYPE 1 (*FAH*) **negative**
 TYROSINEMIA, TYPE 2 (*TAT*) **negative**

U

USHER SYNDROME, TYPE 1B (*MYO7A*) **negative**
 USHER SYNDROME, TYPE 1C (*USH1C*) **negative**
 USHER SYNDROME, TYPE 1D (*CDH23*) **negative**
 USHER SYNDROME, TYPE 1F (*PCDH15*) **negative**
 USHER SYNDROME, TYPE 1J/DEAFNESS, AUTOSOMAL RECESSIVE, 48 (*CIB2*) **negative**
 USHER SYNDROME, TYPE 2A (*USH2A*) **negative**
 USHER SYNDROME, TYPE 2C (*ADGRV1*) **negative**
 USHER SYNDROME, TYPE 3 (*CLRN1*) **negative**

V

VERY LONG-CHAIN ACYL-CoA DEHYDROGENASE DEFICIENCY (*ACADVL*) **see first page**
 VICI SYNDROME (*EPG5*) **negative**
 VITAMIN D-DEPENDENT RICKETS, TYPE 1A (*CYP27B1*) **negative**
 VITAMIN D-RESISTANT RICKETS TYPE 2A (*VDR*) **negative**
 VLDLR-ASSOCIATED CEREBELLAR HYPOPLASIA (*VLDLR*) **negative**

W

WALKER-WARBURG SYNDROME, CRPPA-RELATED (*CRPPA*) **negative**
 WALKER-WARBURG SYNDROME, FKTN-RELATED (*FKTN*) **negative**
 WALKER-WARBURG SYNDROME, LARGE1-RELATED (*LARGE1*) **negative**
 WALKER-WARBURG SYNDROME, POMT1-RELATED (*POMT1*) **negative**
 WALKER-WARBURG SYNDROME, POMT2-RELATED (*POMT2*) **negative**
 WARSAW BREAKAGE SYNDROME (*DDX11*) **negative**
 WERNER SYNDROME (*WRN*) **negative**
 WILSON DISEASE (*ATP7B*) **negative**
 WOLCOTT-RALLISON SYNDROME (*EIF2AK3*) **negative**
 WOLMAN DISEASE (*LIPA*) **negative**
 WOODHOUSE-SAKATI SYNDROME (*DCAF17*) **negative**

X

XERODERMA PIGMENTOSUM VARIANT TYPE (*POLH*) **negative**
 XERODERMA PIGMENTOSUM, GROUP A (*XPA*) **negative**
 XERODERMA PIGMENTOSUM, GROUP C (*XPC*) **negative**

Z

ZELLWEGER SPECTRUM DISORDER, PEX13-RELATED (*PEX13*) **negative**
 ZELLWEGER SPECTRUM DISORDER, PEX16-RELATED (*PEX16*) **negative**
 ZELLWEGER SPECTRUM DISORDER, PEX5-RELATED (*PEX5*) **negative**
 ZELLWEGER SPECTRUM DISORDERS, PEX10-RELATED (*PEX10*) **negative**
 ZELLWEGER SPECTRUM DISORDERS, PEX12-RELATED (*PEX12*) **negative**
 ZELLWEGER SPECTRUM DISORDERS, PEX1-RELATED (*PEX1*) **negative**
 ZELLWEGER SPECTRUM DISORDERS, PEX26-RELATED (*PEX26*) **negative**
 ZELLWEGER SPECTRUM DISORDERS, PEX2-RELATED (*PEX2*) **negative**

Patient Information

Patient Name:

Test Information

Ordering Physician:



Date Of Birth:



Case File ID:



Clinic Information:

Report Date:

Z

ZELLWEGER SPECTRUM DISORDERS, PEX6-RELATED (PEX6) **negative**

Patient Information

Patient Name:

Test Information

Ordering Physician:



Date Of Birth:

Clinic Information:

Case File ID:

Report Date:

Testing Methodology, Limitations, and Comments:**Next-generation sequencing (NGS)**

Sequencing library prepared from genomic DNA isolated from a patient sample is enriched for targets of interest using standard hybridization capture protocols and PCR amplification (for targets specified below). NGS is then performed to achieve the standards of quality control metrics, including a minimum coverage of 99% of targeted regions at 20X sequencing depth. Sequencing data is aligned to human reference sequence, followed by deduplication, metric collection and variant calling (coding region +/- 20bp). Variants are then classified according to ACMGG/AMP standards of interpretation using publicly available databases including but not limited to ENSEMBL, HGMD Pro, ClinGen, ClinVar, 1000G, ESP and gnomAD. Variants predicted to be pathogenic or likely pathogenic for the specified diseases are reported. It should be noted that the data interpretation is based on our current understanding of the genes and variants at the time of reporting. Putative positive sequencing variants that do not meet internal quality standards or are within highly homologous regions are confirmed by Sanger sequencing or gene-specific long-range PCR as needed prior to reporting.

Copy Number Variant (CNV) analysis is limited to deletions involving two or more exons for all genes on the panel, in addition to specific known recurrent single-exon deletions. CNVs of small size may have reduced detection rate. This method does not detect gene inversions, single-exonic and sub-exonic deletions (unless otherwise specified), and duplications of all sizes (unless otherwise specified). Additionally, this method does not define the exact breakpoints of detected CNV events. Confirmation testing for copy number variation is performed by specific PCR, Multiplex Ligation-dependent Probe Amplification (MLPA), next generation sequencing, or other methodology.

This test may not detect certain variants due to local sequence characteristics, high/low genomic complexity, homologous sequence, or allele dropout (PCR-based assays). Variants within noncoding regions (promoter, 5'UTR, 3'UTR, deep intronic regions, unless otherwise specified), small deletions or insertions larger than 25bp, low-level mosaic variants, structural variants such as inversions, and/or balanced translocations may not be detected with this technology.

SPECIAL NOTES

For ABCC6, sequencing variants in exons 1-7 are not detected due to the presence of regions of high homology.

For CFTR, when the CFTR R117H variant is detected, reflex analysis of the polythymidine variations (5T, 7T and 9T) at the intron 9 branch/acceptor site of the CFTR gene will be performed. Multi-exon duplication analysis is included.

For CYP21A2, targets were enriched using long-range PCR amplification, followed by next generation sequencing. Duplication analysis will only be performed and reported when c.955C>T (p.Q319*) is detected. Sequencing and CNV analysis may have reduced sensitivity, if variants result from complex rearrangements, in trans with a gene deletion, or CYP21A2 gene duplication on one chromosome and deletion on the other chromosome. This analysis cannot detect sequencing variants located on the CYP21A2 duplicated copy.

For DDX11, sequencing variants in exons 7-11 and CNV for the entire gene are not analyzed due to high sequence homology.

For GJB2, CNV analysis of upstream deletions of GJB6-D13S1830 (309kb deletion) and GJB6-D13S1854 (232kb deletion) is included.

For HBA1/HBA2, CNV analysis is offered to detect common deletions of -alpha3.7, -alpha4.2, --MED, --SEA, --FIL, --THAI, --alpha20.5, and/or HS-40.

For OTOA, sequencing variants in exons 25-29 and CNV in exons 21-29 are not analyzed due to high sequence homology.

For RPGRIP1L, variants in exon 23 are not detected due to assay limitation.

For SAMD9, only p.K1495E variant will be analyzed and reported.

Friedreich Ataxia (FXN)

The GAA repeat region of the FXN gene is assessed by trinucleotide PCR assay and capillary electrophoresis. Variances of +/-1 repeat for normal alleles and up to +/-3 repeats for premutation alleles may occur. For fully penetrant expanded alleles, the precise repeat size cannot be determined, therefore the approximate allele size is reported. Sequencing and copy number variants are analyzed by next-generation sequencing analysis.

Friedreich Ataxia Repeat Categories

Categories	GAA Repeat Sizes
Normal	<34
Premutation	34 - 65
Full	>65

Patient Information

Patient Name: [REDACTED]

Test Information

Ordering Physician: [REDACTED]



Clinic Information: [REDACTED]

Date Of Birth: [REDACTED]

Case File ID: [REDACTED]

Report Date: [REDACTED]

Spinal Muscular Atrophy (SMN1)

The total combined copy number of SMN1 and SMN2 exon 7 is quantified based on NGS read depth. The ratio of SMN1 to SMN2 is calculated based on the read depth of a single nucleotide that distinguishes these two genes in exon 7. In addition to copy number analysis, testing for the presence or absence of a single nucleotide polymorphism (g.27134T>G in intron 7 of SMN1) associated with the presence of a SMN1 duplication allele is performed using NGS.

Ethnicity	Two SMN1 copies carrier risk before g.27134T>G testing	Carrier risk after g.27134T>G testing	
		g.27134T>G ABSENT	g.27134T>G PRESENT
Caucasian	1 in 632	1 in 769	1 in 29
Ashkenazi Jewish	1 in 350	1 in 580	LIKELY CARRIER
Asian	1 in 628	1 in 702	LIKELY CARRIER
African-American	1 in 121	1 in 396	1 in 34
Hispanic	1 in 1061	1 in 1762	1 in 140

Variant Classification

Only pathogenic or likely pathogenic variants are reported. Other variants including benign variants, likely benign variants, variants of uncertain significance, or inconclusive variants identified during this analysis may be reported in certain circumstances. Our laboratory's variant classification criteria are based on the ACMG and internal guidelines and our current understanding of the specific genes. This interpretation may change over time as more information about a gene and/or variant becomes available. Natera and its lab partner(s) may reclassify variants at certain intervals but may not release updated reports without a specific request made to Natera by the ordering provider. Natera may disclose incidental findings if deemed clinically pertinent to the test performed.

Negative Results

A negative carrier screening result reduces the risk for a patient to be a carrier of a specific disease but does not completely rule out carrier status. Please visit <https://www.natera.com/panel-option/h-all/> for a table of carrier rates, detection rates, residual risks and promised variants/exons per gene. Carrier rates before and after testing vary by ethnicity and assume a negative family history for each disease screened and the absence of clinical symptoms in the patient. Any patient with a family history for a specific genetic disease will have a higher carrier risk prior to testing and, if the disease-causing mutation in their family is not included on the test, their carrier risk would remain unchanged. Genetic counseling is recommended for patients with a family history of genetic disease so that risk figures based on actual family history can be determined and discussed along with potential implications for reproduction. Horizon carrier screening has been developed to identify the reproductive risks for monogenic inherited conditions. Even when one or both members of a couple screen negative for pathogenic variants in a specific gene, the disease risk for their offspring is not zero. There is still a low risk for the condition in their offspring due to a number of different mechanisms that are not detected by Horizon including, but not limited to, pathogenic variant(s) in the tested gene or in a different gene not included on Horizon, pathogenic variant(s) in an upstream regulator, uniparental disomy, de novo mutation(s), or digenic or polygenic inheritance.

Additional Comments

These analyses generally provide highly accurate information regarding the patient's carrier status. Despite this high level of accuracy, it should be kept in mind that there are many potential sources of diagnostic error, including misidentification of samples, polymorphisms, or other rare genetic variants that interfere with analysis. Families should understand that rare diagnostic errors may occur for these reasons.