

GENETIC TESTING SUMMARY

Sperm Donor # 8326

This Genetic Testing Summary provides general information regarding a donor's genetic test results. It is not comprehensive and does not replace a clinical genetic counseling consultation or constitute medical advice. Clients should review the donor's complete genetic test results with their medical provider or fertility specialist.

Genetic testing technology and scientific information change over time. Testing is not available for every genetic condition, and Fairfax does not test for every condition for which testing is available. A negative result reduces, but does not eliminate, the risk of having an affected child.

Last Updated: 08/25/2026

Genetic Test	Result	Comments
Chromosome analysis (karyotype)	Normal male karyotype	No evidence of clinically significant chromosome abnormalities
Expanded Genetic Carrier Screening: 549 diseases by gene sequencing and del/dup analysis.	Carrier: Mucopolysaccharidosis, Type III A (Sanfilippo A) (SGSH) Carrier: PIGN-Congenital Disorder of Glycosylation (PIGN) Carrier: Salla Disease (SLC17A5) Carrier: Xeroderma Pigmentosum, Group C (XPC) Negative for other genes tested.	Egg source carrier screening is indicated before using this donor

Patient Information

Patient Name: Donor 8326

Date Of Birth: [REDACTED]

Gender: Male

Patient ID: N/A

Medical Record #: 8326-[REDACTED]

Collection Kit: [REDACTED]

Accession ID: N/A

Case File ID: [REDACTED]

Ethnicity: Other

Test Information

Ordering Physician: [REDACTED]

Clinic Information: Fairfax Cryobank

Phone: N/A

Report Date: 03/04/2026

Sample Collected: 02/18/2026

Sample Received: 02/19/2026

Sample Type: Blood

**Horizon™**
Advanced carrier screening**CARRIER SCREENING REPORT**

ABOUT THIS SCREEN: Horizon™ is a carrier screen for specific autosomal recessive and X-linked diseases. This information can help patients learn their risk of having a child with specific genetic conditions.

ORDER SELECTED: The Horizon Custom panel was ordered for this patient. Males are not screened for X-linked diseases

FINAL RESULTS SUMMARY:**CARRIER for Mucopolysaccharidosis, Type Iii A (Sanfilippo A)**

Positive for the pathogenic variant c.220C>T (p.R74C) in the SGSH gene. If this individual's partner is a carrier for MUCOPOLYSACCHARIDOSIS, TYPE III A (SANFILIPPO A), their chance to have a child with this condition is 1 in 4 (25%). Carrier screening for this individual's partner is suggested.

CARRIER for PIGN-Congenital Disorder Of Glycosylation

Positive for the likely pathogenic variant c.2354G>A (p.R785H) in the PIGN gene. If this individual's partner is a carrier for PIGN-CONGENITAL DISORDER OF GLYCOSYLATION, their chance to have a child with this condition may be as high as 1 in 4 (25%). Carrier screening for this individual's partner is suggested.

CARRIER for Salla Disease

Positive for the pathogenic variant c.1138_1139del (p.V380Sfs*8) in the SLC17A5 gene. If this individual's partner is a carrier for SALLA DISEASE, their chance to have a child with this condition is 1 in 4 (25%). Carrier screening for this individual's partner is suggested.

CARRIER for Xeroderma Pigmentosum, Group C

Positive for the pathogenic variant c.342_343del (p.A116Yfs*4) in the XPC gene. If this individual's partner is a carrier for XERODERMA PIGMENTOSUM, GROUP C, their chance to have a child with this condition is 1 in 4 (25%). Carrier screening for this individual's partner is suggested.

Negative for 545 out of 549 diseases

No other pathogenic variants were detected in the genes that were screened. The patient's remaining carrier risk after the negative screening results is listed for each disease/gene on the Horizon website at <https://www.natera.com/panel-option/h-all/>. Please see the following pages of this report for a comprehensive list of all conditions included on this individual's screen.

Carrier screening is not diagnostic and may not detect all possible pathogenic variants in a given gene.

Christine M. Eng, M.D.
Medical Director, Baylor Genetics
J. Dianne Keen-Kim, Ph.D., FACMG
Senior Laboratory Director, Natera
Linyan Meng, Ph.D.
Laboratory Director, Baylor Genetics
Yang Wang, Ph.D., FACMG
Laboratory Director, Natera

Patient Information

Patient Name: Donor 8326

Test Information

Ordering Physician: [REDACTED]

**Horizon™**
Advanced carrier screening

Date Of Birth: [REDACTED]

Case File ID: [REDACTED]

Clinic Information: Fairfax Cryobank

Report Date: 03/04/2026

RECOMMENDATIONS

Individuals who would like to review their Horizon report with a Natera Laboratory Genetic Counselor may schedule a telephone genetic information session by calling 650-249-9090 or visiting naterasession.com. Clinicians with questions may contact Natera at 650-249-9090 or email support@natera.com. Individuals with positive results may wish to discuss these results with family members to allow them the option to be screened. Comprehensive genetic counseling to discuss the implications of these test results and possible associated reproductive risk is recommended.



Patient Information

Patient Name: Donor 8326

Test Information

Ordering Physician: [REDACTED]



Date Of Birth: [REDACTED]

Case File ID: [REDACTED]

Clinic Information: Fairfax Cryobank

Report Date: 03/04/2026

MUCOPOLYSACCHARIDOSIS, TYPE III A (SANFILIPPO A)**Understanding Your Horizon Carrier Screen Results****What is Mucopolysaccharidosis, Type IIIA (Sanfilippo A)?**

Mucopolysaccharidosis (MPS), Type IIIA (also called Sanfilippo A) is an inherited disorder that affects many parts of the body. Signs and symptoms of MPS, Type IIIA usually begin in early childhood and include unusual facial features, a large head size, bone and joint abnormalities, intellectual disability, behavioral problems, sleep difficulties, and coordination and movement problems. Other symptoms include recurrent respiratory and ear infections, vision problems, hearing loss, and hernias. Children lose developmental skills over time, symptoms worsen, and lifespan is shortened with death usually occurring by early adulthood. In some cases, affected individuals have been treated with or participated in clinical trials using stem cell transplantation from cord blood or bone marrow. Couples at risk of having an affected child may consider cord blood banking, as siblings have a higher chance of being a match for stem cell transplantation than a non-related individual. More information can be found at: <https://parentsguidecordblood.org/en>. Clinical trials involving potential new treatments for this condition may be available (see www.clinicaltrials.gov).

What causes Mucopolysaccharidosis, Type IIIA (Sanfilippo A)?

MPS, Type IIIA is caused by a change, or mutation, in both copies of the SGSH gene pair. These mutations cause the genes to not work properly or not work at all. When both copies of the SGSH gene do not work correctly, it leads to the symptoms described above. MPS, Type IIIA is inherited in an autosomal recessive manner. This means that, in most cases, both parents must be carriers of a mutation in one copy of the SGSH gene to have a child with MPS, Type IIIA. People who are carriers for MPS, Type IIIA are usually healthy and do not have symptoms nor do they have MPS, Type IIIA themselves. Usually a child inherits two copies of each gene, one copy from the mother and one copy from the father. If the mother and father are both carriers for MPS, Type IIIA there is a 1 in 4, or 25%, chance in each pregnancy for both partners to pass on their SGSH gene mutations to the child, who will then have this condition. Individuals found to carry more than one mutation for MPS, Type III should discuss their risk for having an affected child with their health care provider. There are many other types of Mucopolysaccharidosis, each caused by mutations in different genes. A carrier for MPS, Type IIIA is not likely to be at increased risk for having children with the other forms of MPS.

What can I do next?

You may wish to speak with a local genetic counselor about your carrier test results. A genetic counselor in your area can be located on the National Society of Genetic Counselors website (www.nsgc.org). Your siblings and other relatives are at increased risk to also have this mutation. You are encouraged to inform your family members of your test results as they may wish to consider being tested themselves. If you are pregnant, your partner can have carrier screening for MPS, Type IIIA ordered by a health care professional. If your partner is not found to be a carrier for MPS, Type IIIA, your risk of having a child with this condition is greatly reduced. Couples at risk of having a baby with MPS, Type IIIA can opt to have prenatal diagnosis done through chorionic villus sampling (CVS) or amniocentesis during pregnancy or can choose to have the baby tested after birth for this condition. If you are not yet pregnant, your partner can have carrier screening for MPS, Type IIIA ordered by a health care professional. If your partner is found to be a carrier for MPS, Type IIIA, you have several reproductive options to consider:

- Natural pregnancy with or without prenatal diagnosis of the fetus or testing the baby after birth for MPS, Type IIIA
- Preimplantation genetic diagnosis (PGD) with in vitro fertilization (IVF) to test embryos for MPS, Type IIIA
- Adoption or use of a sperm or egg donor who is not a carrier for MPS, Type IIIA

What resources are available?

- National MPS Society: <http://mpssociety.org/mps/mps-iii/>
- Prenatal diagnosis done through CVS: <http://www.marchofdimes.org/chorionic-villus-sampling.aspx>
- Prenatal diagnosis done through Amniocentesis: <http://www.marchofdimes.org/amniocentesis.aspx>
- PGD with IVF: <http://natera.com/spectrum>



Patient Information

Patient Name: Donor 8326

Test Information

Ordering Physician: [REDACTED]



Date Of Birth: [REDACTED]

Case File ID: [REDACTED]

Clinic Information: Fairfax Cryobank

Report Date: 03/04/2026

PIGN-CONGENITAL DISORDER OF GLYCOSYLATION

Understanding Your Horizon Carrier Screen Results

What does being a carrier mean?

Your result shows that you are a carrier of PIGN-congenital disorder of glycosylation (PIGN-CDG). A carrier of a genetic condition does not have the condition. Carriers also are not certain to have a child with the condition. We are all carriers of one or more genetic conditions.

Your children are not at high risk for this condition unless your partner or donor is also a carrier of PIGN-CDG. Further testing can be done to see if your partner or donor is a carrier.

What is PIGN-congenital disorder of glycosylation (PIGN-CDG)?

PIGN-CDG can affect many areas of the body, including the brain, muscles, and stomach. There are two forms of PIGN-CDG called multiple congenital anomalies-hypotonia-seizures syndrome 1 (MCAHS1) and Fryns syndrome.¹

Babies with **MCAHS1** start having symptoms soon after birth or in the first few months of life. They can be born with differences in the way their organs are formed, including their heart, stomach, intestines, and urinary tract. They also have low muscle tone and often have shaky eye movements. Children with MCAHS1 can have seizures that cannot be fully controlled with medication and intellectual disability. They are slow to sit and crawl and usually do not learn how to talk. Many children with MCAHS1 will have reflux, where food from their stomach moves back up into their food pipe (esophagus). Most people with MCAHS1 die in early childhood.^{1,2,3,4}

Fryns syndrome is a more severe form of MCAHS1. Babies with Fryns syndrome have more severe differences in the way their organs are formed than babies with MCAHS1. They can also have congenital diaphragmatic hernia (CDH). CDH is a hole in the muscle that separates the stomach and other organs from the chest (the diaphragm). CDH causes poor lung development and breathing difficulty. Most babies with Fryns syndrome die before birth or shortly after.^{1,4,5}

Rarely, children with PIGN-CDG have brain and muscle problems without CDH or other differences in the way their organs are formed.¹

Currently there is no cure for PIGN-CDG, and treatment is based on symptoms.^{1,2,3,4,5} Clinical trials involving potential new treatments for this condition could be available (see clinicaltrials.gov).

What causes PIGN-congenital disorder of glycosylation (PIGN-CDG)?

PIGN-CDG is caused by changes, or variants, in the PIGN gene. These changes make the gene not work properly. Genes are a set of instructions inside the cells of our bodies that tell our bodies how to grow and function. Everyone has two copies of the PIGN gene. Carriers of PIGN-CDG have one working copy and one nonworking copy of the gene. People with PIGN-CDG have no working copies of the gene.

PIGN-CDG is usually passed down, or inherited, from both genetic parents. We inherit one copy of the PIGN gene from each of our genetic parents. When both genetic parents are carriers, each child has a 1 in 4 (25%) chance of inheriting two nonworking genes and having PIGN-CDG. Each child also has a 1 in 2 (50%) chance of being a carrier of PIGN-CDG and a 1 in 4 (25%) chance of inheriting two working copies of the gene. This type of inheritance is called autosomal recessive inheritance.

Will my children have PIGN-congenital disorder of glycosylation (PIGN-CDG)?

If your partner or donor also has a nonworking copy of the PIGN gene, your children could have PIGN-CDG. Each child you have together would have a 1 in 4 (25%) chance of having PIGN-CDG. Each child you have together would also have a 3 in 4 (75%) chance of **not** having the condition.

If your partner or donor has PIGN carrier screening and no variants are found, the chance that your children would have PIGN-CDG is very low. No further testing would usually be needed for you, your partner or donor, or your children related to PIGN-CDG.

What can I do next?

If you want to know if your children are at risk for PIGN-CDG, your partner or donor would need to have PIGN carrier screening. If you have questions about this testing, please ask your healthcare provider or use the resources below. Many people find it helpful to speak with a genetic counselor.

If your partner or donor is found to be a PIGN-CDG carrier, your children would be at risk for having PIGN-CDG.

If you or your partner or surrogate are currently pregnant, tests called CVS (chorionic villus sampling) and amniocentesis can be done during pregnancy to find out if a baby has PIGN-CDG. These tests both have a small risk of miscarriage. Babies can also be tested for PIGN-CDG after birth instead.

If you or your partner or surrogate are not yet pregnant, you could have these options:

- natural pregnancy with CVS or amniocentesis to test for PIGN-CDG during pregnancy;
- natural pregnancy and testing the baby after birth for PIGN-CDG;
- preimplantation genetic testing (PGT-M) with in vitro fertilization (IVF) to test embryos for PIGN-CDG;



Patient Information

Patient Name: Donor 8326

Test Information

Ordering Physician: [REDACTED]



Horizon™
Advanced carrier screening

Date Of Birth: [REDACTED]

Case File ID: [REDACTED]

Clinic Information: Fairfax Cryobank

Report Date: 03/04/2026

- adoption; or
- use of a sperm or egg donor who had no variants found in PIGN carrier screening.

Where can I find more information?

- Genetic and Rare Disease Information Center rarediseases.info.nih.gov/diseases/12781/multiple-congenital-anomalies-hypotonia-seizures-syndrome
- CDH International cdhi.org
- Epilepsy Foundation epilepsy.com
- CVS marchofdimes.org/chorionic-villus-sampling
- Amniocentesis marchofdimes.org/pregnancy/amniocentesis

What does this mean for my family?

You likely got (inherited) this nonworking gene from one of your genetic parents. Your genetic siblings and other family members could also carry it. You should tell your family members about your test result so they can decide if they want carrier screening for PIGN-CDG.

References

1. Loong L et al. Biallelic variants in PIGN cause Fryns syndrome, multiple congenital anomalies-hypotonia-seizures syndrome, and neurologic phenotypes: A genotype-phenotype correlation study. *Genet Med*. 2023 Jan;25(1):37-48. doi: 10.1016/j.jim.2022.09.007. Epub 2022 Nov 2. PMID: 36322149.
2. Fleming L et al. Genotype-phenotype correlation of congenital anomalies in multiple congenital anomalies hypotonia seizures syndrome (MCAHS1)/PIGN-related epilepsy. *Am J Med Genet A*. 2016 Jan;170A(1):77-86. doi: 10.1002/ajmg.a.37369. Epub 2015 Sep 23. PMID: 26394714; PMCID: PMC4886552.
3. Maydan G et al. Multiple congenital anomalies-hypotonia-seizures syndrome is caused by a mutation in PIGN. *J Med Genet*. 2011 Jun;48(6):383-9. doi: 10.1136/jmg.2010.087114. Epub 2011 Apr 14. PMID: 21493957.
4. Paprocka J et al. Congenital Disorders of Glycosylation from a Neurological Perspective. *Brain Sci*. 2021 Jan 11;11(1):88. doi: 10.3390/brainsci11010088. PMID: 33440761; PMCID: PMC7827962.
5. Slavotinek A. Fryns Syndrome. 2007 Apr 18 [Updated 2020 Sep 17]. In: Adam MP et al, editors. *GeneReviews®* [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2024. Available from: <https://www.ncbi.nlm.nih.gov/sites/books/NBK1459/>. Accessed December 2024.



Patient Information

Patient Name: Donor 8326

Test Information

Ordering Physician: [REDACTED]



Date Of Birth: [REDACTED]

Case File ID: [REDACTED]

Clinic Information: Fairfax Cryobank

Report Date: 03/04/2026

SALLA DISEASE**Understanding Your Horizon Carrier Screen Results****What is Salla Disease?**

Salla Disease, also known as Sialic Acid Storage Disease, is an inherited disorder that mainly affects the nervous system. There are multiple forms of Salla Disease/Sialic Acid Storage Disease. The most severe form is called Infantile Free Sialic Acid Storage Disease (ISSD) and has symptoms that start in infancy or before birth. Symptoms include enlarged liver and heart, seizures, and buildup of fluid in the abdomen (ascites). Infants with this severe form usually do not survive past childhood. Infants with the mildest form of Salla Disease appear normal at birth. Then during infancy, signs and symptoms begin to appear including slowly progressing loss of skills, poor muscle tone (hypotonia) that changes with time to tight and stiff muscles (spasticity), seizures, developmental delay, intellectual disability, speech problems, coordination problems (ataxia), and slow involuntary movements (athetosis) of the arms and legs. Although symptoms vary from person to person, about two-thirds of people with Salla Disease are not able to walk. Most people with Salla Disease live into adulthood. There is also an intermediate form of the disorder. Currently there is no cure for this condition and treatment is based on the symptoms. Clinical trials involving potential new treatments for this condition may be available (see www.clinicaltrials.gov).

What causes Salla Disease?

Salla Disease is caused by a gene change, or mutation, in both copies of the SLC17A5 gene pair. These mutations cause the genes to not work properly or not work at all. When both copies of this gene do not work correctly, it leads to the symptoms described above. It is sometimes, but not always, possible to determine whether a specific mutation in the SLC17A5 will cause the severe, intermediate, or mild form of the disorder. Salla Disease is inherited in an autosomal recessive manner. This means that, in most cases, both parents must be carriers of a mutation in one copy of the SLC17A5 gene to have a child with Salla Disease. People who are carriers for Salla Disease are usually healthy and do not have symptoms nor do they have Salla Disease themselves. Usually a child inherits two copies of each gene, one copy from the mother and one copy from the father. If the mother and father are both carriers for Salla Disease, there is a 1 in 4, or 25%, chance in each pregnancy for both partners to pass on their SLC17A5 gene mutations to the child, who will then have Salla Disease. Individuals found to carry more than one mutation for Salla Disease should discuss their risk for having an affected child with their health care provider.

What can I do next?

You may wish to speak with a local genetic counselor about your carrier test results. A genetic counselor in your area can be located on the National Society of Genetic Counselors website (www.nsgc.org). Your siblings and other relatives are at increased risk to also have this mutation. You are encouraged to inform your family members of your test results as they may wish to consider being tested themselves. If you are pregnant, your partner can have carrier screening for Salla Disease ordered by a health care professional. If your partner is not found to be a carrier for Salla Disease, your risk of having a child with this condition is greatly reduced. Couples at risk of having a baby with Salla Disease can opt to have prenatal diagnosis done through chorionic villus sampling (CVS) or amniocentesis during pregnancy or can choose to have the baby tested after birth for this condition. If you are not yet pregnant, your partner can have carrier screening for Salla Disease ordered by a health care professional. If your partner is found to be a carrier for Salla Disease, you have several reproductive options to consider:

- Natural pregnancy with or without prenatal diagnosis of the fetus or testing the baby after birth for Salla Disease
- Preimplantation genetic diagnosis (PGD) with in vitro fertilization (IVF) to test embryos for Salla Disease
- Adoption or use of a sperm or egg donor who is not a carrier for Salla Disease

What resources are available?

- Genetics Home Reference: <http://ghr.nlm.nih.gov/condition/sialic-acid-storage-disease>
- GeneReviews: <https://www.ncbi.nlm.nih.gov/books/NBK1470/>
- Prenatal diagnosis done through CVS: <http://www.marchofdimes.org/chorionic-villus-sampling.aspx>
- Prenatal diagnosis done through Amniocentesis: <http://www.marchofdimes.org/amniocentesis.aspx>
- PGD with IVF: <http://www.natera.com/spectrum>



Patient Information

Patient Name: Donor 8326

Test Information

Ordering Physician: [REDACTED]



Date Of Birth: [REDACTED]

Case File ID: [REDACTED]

Clinic Information: Fairfax Cryobank

Report Date: 03/04/2026

XERODERMA PIGMENTOSUM, GROUP C**Understanding Your Horizon Carrier Screen Results****What is Xeroderma Pigmentosum, Group C?**

Xeroderma Pigmentosum, Group C is an inherited condition that causes the body to have extreme sensitivity to ultraviolet (UV) rays from sunlight. This condition affects the eyes and areas of skin exposed to the sun, and in some cases affects the brain and nervous system. Signs and symptoms of Xeroderma Pigmentosum, Group C usually appear in infancy or early childhood. Affected children can develop a severe sunburn after spending only a few minutes in the sun. By age 2, almost all children with Xeroderma Pigmentosum, Group C have freckling in sun-exposed areas, dry skin, and patchy changes in skin color. The eyes are very sensitive to the sun, leading to irritation and other eye problems that can affect vision. People with Xeroderma Pigmentosum, Group C have a greatly increased risk of developing skin cancer. Unless protected from the sun, about half of children will develop a skin cancer by age 10. Most people with Xeroderma Pigmentosum, Group C will have a number of skin cancers during their life. There may also be an increased risk for other types of cancer, including eye cancer and brain tumor, although more studies need to be done to know for sure. Lifelong medical care is important for both prevention and early detection of skin cancers. Some people with Xeroderma Pigmentosum, Group C develop neurological problems that worsen over time. These problems may include hearing loss, poor coordination, difficulty walking, movement problems, loss of intellectual abilities, difficulty swallowing and talking, and seizures. Currently there is no cure for this disorder and treatment is based on symptoms. Clinical trials involving potential new treatments for this condition may be available (see www.clinicaltrials.gov).

What causes Xeroderma Pigmentosum, Group C?

Xeroderma Pigmentosum, Group C is caused by a change, or mutation, in both copies of the XPC gene pair. These mutations cause the genes to not work properly or not work at all. The normal function of the XPC gene pair is to help repair the body's DNA. When both copies of the XPC gene do not work correctly, DNA mistakes are left uncorrected, leading to the symptoms described above. Xeroderma Pigmentosum, Group C is inherited in an autosomal recessive manner. This means that, in most cases, both parents must be carriers of a mutation in one copy of the XPC gene to have a child with Xeroderma Pigmentosum, Group C. People who are carriers for Xeroderma Pigmentosum, Group C are usually healthy and do not have symptoms, nor do they have the disorder themselves. Usually a child inherits two copies of each gene, one copy from the mother and one copy from the father. If the mother and father are both carriers for Xeroderma Pigmentosum, Group C there is a 1 in 4, or 25%, chance in each pregnancy for both partners to pass on their XPC gene mutations to the child, who will then have this disorder. Individuals found to carry more than one mutation for Xeroderma Pigmentosum, Group C should discuss their risk for having an affected child with their health care provider. There are a several forms of Xeroderma Pigmentosum, each caused by mutations in different genes. Carriers for Xeroderma Pigmentosum, Group C are not likely to be at increased risk of having a child with these other forms of this group of disorders.

What can I do next?

You may wish to speak with a local genetic counselor about your carrier test results. A genetic counselor in your area can be located on the National Society of Genetic Counselors website (www.nsgc.org). Your siblings and other relatives are at increased risk to also have this mutation. You are encouraged to inform your family members of your test results as they may wish to consider being tested themselves. If you are pregnant, your partner can have carrier screening for Xeroderma Pigmentosum, Group C ordered by a health care professional. If your partner is not found to be a carrier for the Xeroderma Pigmentosum, Group C, your risk of having an affected child is greatly reduced. Couples at risk of having a baby with Xeroderma Pigmentosum, Group C can opt to have prenatal diagnosis done through chorionic villus sampling (CVS) or amniocentesis during pregnancy or can choose to have the baby tested after birth. If you are not yet pregnant, your partner can have carrier screening for Xeroderma Pigmentosum, Group C ordered by a health care professional. If your partner is found to be a carrier for Xeroderma Pigmentosum, Group C, you have several reproductive options to consider:

- Natural pregnancy with or without prenatal diagnosis of the fetus or testing the baby after birth for Xeroderma Pigmentosum, Group C
- Preimplantation genetic diagnosis (PGD) with in vitro fertilization (IVF) to test embryos for Xeroderma Pigmentosum, Group C
- Adoption or use of a sperm or egg donor who is not a carrier for Xeroderma Pigmentosum, Group C

What resources are available?

- Genetics Home Reference: <https://ghr.nlm.nih.gov/condition/xeroderma-pigmentosum>
- Prenatal diagnosis done through CVS: <http://www.marchofdimes.org/chorionic-villus-sampling.aspx>
- Prenatal diagnosis done through Amniocentesis: <http://www.marchofdimes.org/amniocentesis.aspx> PGD with IVF: <http://www.natera.com/spectrum>



Patient Information

Patient Name: Donor 8326

Test Information

Ordering Physician: [REDACTED]

**Horizon™**
Advanced carrier screening

Date Of Birth: [REDACTED]

Case File ID: [REDACTED]

Clinic Information: Fairfax Cryobank

Report Date: 03/04/2026

VARIANT DETAILS**PIGN, c.2354G>A (p.R785H), likely pathogenic**

- The c.2354G>A (p.R785H) variant in the PIGN gene has been observed at a frequency of 0.0061% in the gnomAD v2.1.1 dataset.
- This variant has been reported in a homozygous state or in conjunction with another variant in individual(s) with multiple congenital anomalies-hypotonia-seizures syndrome 1 (PMID: 33619735).
- This variant has been reported in ClinVar [ID: 378363].

SGSH, c.220C>T (p.R74C), pathogenic

- The c.220C>T (p.R74C) variant in the SGSH gene has been observed at a frequency of 0.0192% in the gnomAD v2.1.1 dataset.
- This variant has been reported in a homozygous state or in conjunction with another variant in individual(s) with mucopolysaccharidosis, type IIIA (Sanfilippo syndrome A) (PMID: 9285796, 18407553).
- This variant has been reported in ClinVar [ID: 5108].

SLC17A5, c.1138_1139del (p.V380Sfs*8), pathogenic

- The c.1138_1139del (p.V380Sfs*8) variant in the SLC17A5 gene has been observed at a frequency of 0.0049% in the gnomAD v2.1.1 dataset.
- This variant has been reported in a homozygous state or in conjunction with another variant in individual(s) with Salla disease (PMID: 10947946).
- This premature termination variant is predicted to cause nonsense-mediated decay (NMD) in a gene where loss-of-function is a known mechanism of disease.
- This variant has been reported in ClinVar [ID: 56552].

XPC, c.342_343del (p.A116Yfs*4), pathogenic

- The c.342_343del (p.A116Yfs*4) variant in the XPC gene has been observed at a frequency of 0.0007% in the gnomAD v2.1.1 dataset.
- This variant has been observed in a homozygous state in individual(s) with Xeroderma pigmentosum group C (PMID: 23173980).
- This premature termination variant is predicted to cause nonsense-mediated decay (NMD) in a gene where loss-of-function is a known mechanism of disease.
- This variant has been described in ClinVar [ID: 553148].

Patient Information

Patient Name: Donor 8326

Test Information

Ordering Physician:



Clinic Information: Fairfax Cryobank

Date Of Birth:

Case File ID:

Report Date: 03/04/2026

DISEASES SCREENED

Below is a list of all diseases screened and the result. Certain conditions have unique patient-specific numerical values, therefore, results for those conditions are formatted differently.

Autosomal Recessive**1**17-BETA HYDROXYSTEROID DEHYDROGENASE 3 DEFICIENCY (*HSD17B3*) **negative****3**

3-BETA-HYDROXYSTEROID DEHYDROGENASE TYPE II DEFICIENCY (*HSD3B2*) **negative**
 3-HYDROXY-3-METHYLGLUTARYL-COENZYME A LYASE DEFICIENCY (*HMGCL*) **negative**
 3-HYDROXYACYL-COA DEHYDROGENASE DEFICIENCY (*HADH*) **negative**
 3-METHYLCROTONYL-CoA CARBOXYLASE 2 DEFICIENCY (*MCCC2*) **negative**
 3-PHOSPHOGLYCERATE DEHYDROGENASE DEFICIENCY (*PHGDH*) **negative**

55-ALPHA-REDUCTASE DEFICIENCY (*SRD5A2*) **negative****6**6-PYRUVYL-TETRAHYDROPTERIN SYNTHASE (*PTPS*) DEFICIENCY (*PTS*) **negative****A**

ABCA4-RELATED CONDITIONS (*ABCA4*) **negative**
 ABETALIPROTEINEMIA (*MTTP*) **negative**
 ACHONDROGENESIS, TYPE 1B (*SLC26A2*) **negative**
 ACHROMATOPSIA, CNGB3-RELATED (*CNGB3*) **negative**
 ACRODERMATITIS ENTEROPATHICA (*SLC39A4*) **negative**
 ACTION MYOCLONUS-RENAL FAILURE (AMRF) SYNDROME (*SCARB2*) **negative**
 ACUTE INFANTILE LIVER FAILURE, TRMU-RELATED (*TRMU*) **negative**
 ACYL-COA OXIDASE I DEFICIENCY (*ACOX1*) **negative**
 AICARDI-GOUTIERES SYNDROME (*SAMHD1*) **negative**
 AICARDI-GOUTIERES SYNDROME, RNASEH2A-RELATED (*RNASEH2A*) **negative**
 AICARDI-GOUTIERES SYNDROME, RNASEH2B-RELATED (*RNASEH2B*) **negative**
 AICARDI-GOUTIERES SYNDROME, RNASEH2C-RELATED (*RNASEH2C*) **negative**
 AICARDI-GOUTIERES SYNDROME, TREX1-RELATED (*TREX1*) **negative**
 ALPHA-MANNOSIDOSIS (*MAN2B1*) **negative**
 ALPHA-THALASSEMIA (*HBA1/HBA2*) **negative**
 ALPORT SYNDROME, COL4A3-RELATED (*COL4A3*) **negative**
 ALPORT SYNDROME, COL4A4-RELATED (*COL4A4*) **negative**
 ALSTROM SYNDROME (*ALMS1*) **negative**
 AMISH INFANTILE EPILEPSY SYNDROME (*ST3GAL5*) **negative**
 ANDERMANN SYNDROME (*SLC12A6*) **negative**
 ARGININE:GLYCINE AMIDINOTRANSFERASE DEFICIENCY (AGAT DEFICIENCY) (*GATM*) **negative**
 ARGININEMIA (*ARG1*) **negative**
 ARGININOSUCCINATE LYASE DEFICIENCY (*ASL*) **negative**
 AROMATASE DEFICIENCY (*CYP19A1*) **negative**
 ASPARAGINE SYNTHETASE DEFICIENCY (*ASNS*) **negative**
 ASPARTYLGLYCOSAMINURIA (*AGA*) **negative**
 ATAXIA WITH VITAMIN E DEFICIENCY (*TTPA*) **negative**
 ATAXIA-TELANGIECTASIA (*ATM*) **negative**
 ATAXIA-TELANGIECTASIA-LIKE DISORDER 1 (*MRE11*) **negative**
 ATRANSFERRINEMIA (*TF*) **negative**
 AUTISM SPECTRUM, EPILEPSY AND ARTHROGRYPOSIS (*SLC35A3*) **negative**
 AUTOIMMUNE POLYGLANDULAR SYNDROME, TYPE 1 (*AIRE*) **negative**
 AUTOSOMAL RECESSIVE CONGENITAL ICHTHYOSIS (*ARCI*), *SLC27A4*-RELATED (*SLC27A4*) **negative**
 AUTOSOMAL RECESSIVE SPASTIC ATAXIA OF CHARLEVOIX-SAGUENAY (*SACS*) **negative**

BBARDET-BIEDL SYNDROME, ARL6-RELATED (*ARL6*) **negative**

BARDET-BIEDL SYNDROME, BBS10-RELATED (*BBS10*) **negative**
 BARDET-BIEDL SYNDROME, BBS12-RELATED (*BBS12*) **negative**
 BARDET-BIEDL SYNDROME, BBS1-RELATED (*BBS1*) **negative**
 BARDET-BIEDL SYNDROME, BBS2-RELATED (*BBS2*) **negative**
 BARDET-BIEDL SYNDROME, BBS4-RELATED (*BBS4*) **negative**
 BARDET-BIEDL SYNDROME, BBS5-RELATED (*BBS5*) **negative**
 BARDET-BIEDL SYNDROME, BBS7-RELATED (*BBS7*) **negative**
 BARDET-BIEDL SYNDROME, BBS9-RELATED (*BBS9*) **negative**
 BARDET-BIEDL SYNDROME, TTC8-RELATED (*TTC8*) **negative**
 BARE LYMPHOCYTE SYNDROME, CIITA-RELATED (*CIITA*) **negative**
 BARTTER SYNDROME, BSND-RELATED (*BSND*) **negative**
 BARTTER SYNDROME, KCNJ1-RELATED (*KCNJ1*) **negative**
 BARTTER SYNDROME, SLC12A1-RELATED (*SLC12A1*) **negative**
 BATTEN DISEASE, CLN3-RELATED (*CLN3*) **negative**
 BETA-HEMOGLOBINOPATHIES (*HBB*) **negative**
 BETA-KETOTHIOLASE DEFICIENCY (*ACAT1*) **negative**
 BETA-MANNOSIDOSIS (*MANBA*) **negative**
 BETA-UREIDOPROPIONASE DEFICIENCY (*UPB1*) **negative**
 BILATERAL FRONTOPIRIETAL POLYMICROGYRIA (*GPR56*) **negative**
 BIOTINIDASE DEFICIENCY (*BTID*) **negative**
 BIOTIN-THIAMINE-RESPONSIVE BASAL GANGLIA DISEASE (BTBGD) (*SLC19A3*) **negative**
 BLOOM SYNDROME (*BLM*) **negative**
 BRITTLE CORNEA SYNDROME 1 (*ZNF469*) **negative**
 BRITTLE CORNEA SYNDROME 2 (*PRDM5*) **negative**

C

CANAVAN DISEASE (*ASPA*) **negative**
 CARBAMOYL PHOSPHATE SYNTHETASE I DEFICIENCY (*CPS1*) **negative**
 CARNITINE DEFICIENCY (*SLC22A5*) **negative**
 CARNITINE PALMITOYLTRANSFERASE IA DEFICIENCY (*CPT1A*) **negative**
 CARNITINE PALMITOYLTRANSFERASE II DEFICIENCY (*CPT2*) **negative**
 CARNITINE-ACYLCARNITINE TRANSLOCASE DEFICIENCY (*SLC25A20*) **negative**
 CARPENTER SYNDROME (*RAB23*) **negative**
 CARTILAGE-HAIR HYPOPLASIA (*RMRP*) **negative**
 CATECHOLAMINERGIC POLYMORPHIC VENTRICULAR TACHYCARDIA (*CASQ2*) **negative**
 CD59-MEDIATED HEMOLYTIC ANEMIA (*CD59*) **negative**
 CEP152-RELATED MICROCEPHALY (*CEP152*) **negative**
 CEREBRAL DYSGENESIS, NEUROPATHY, ICHTHYOSIS, AND PALMOPLANTAR KERATODERMA (CEDNIK) SYNDROME (*SNAP29*) **negative**
 CEREBROTENDINOUS XANTHOMATOSIS (*CYP27A1*) **negative**
 CHARCOT-MARIE-TOOTH DISEASE, RECESSIVE INTERMEDIATE C (*PLEKHG5*) **negative**
 CHARCOT-MARIE-TOOTH-DISEASE, TYPE 4D (*NDRG1*) **negative**
 CHEDIAK-HIGASHI SYNDROME (*LYST*) **negative**
 CHOREOACANTHOCYTOSIS (*VPS13A*) **negative**
 CHRONIC GRANULOMATOUS DISEASE, CYBA-RELATED (*CYBA*) **negative**
 CHRONIC GRANULOMATOUS DISEASE, NCF2-RELATED (*NCF2*) **negative**
 CILIOPATHIES, RPGRIP1L-RELATED (*RPGRIP1L*) **negative**
 CITRIN DEFICIENCY (*SLC25A13*) **negative**
 CITRULLINEMIA, TYPE 1 (*ASS1*) **negative**
 CLN10 DISEASE (*CTSD*) **negative**
 COHEN SYNDROME (*VPS13B*) **negative**
 COL11A2-RELATED CONDITIONS (*COL11A2*) **negative**
 COMBINED MALONIC AND METHYLMALONIC ACIDURIA (*ACSF3*) **negative**
 COMBINED OXIDATIVE PHOSPHORYLATION DEFICIENCY 1 (*GFM1*) **negative**
 COMBINED OXIDATIVE PHOSPHORYLATION DEFICIENCY 3 (*TFSM*) **negative**



Patient Information

Patient Name: Donor 8326

Test Information

Ordering Physician: [REDACTED]



Clinic Information: Fairfax Cryobank

Date Of Birth: [REDACTED]

Case File ID: [REDACTED]

Report Date: 03/04/2026

C

COMBINED PITUITARY HORMONE DEFICIENCY 1 (*POU1F1*) **negative**
COMBINED PITUITARY HORMONE DEFICIENCY-2 (*PROP1*) **negative**
CONGENITAL ADRENAL HYPERPLASIA, 11-BETA-HYDROXYLASE DEFICIENCY (*CYP11B1*) **negative**
CONGENITAL ADRENAL HYPERPLASIA, 17-ALPHA-HYDROXYLASE DEFICIENCY (*CYP17A1*) **negative**
CONGENITAL ADRENAL HYPERPLASIA, 21-HYDROXYLASE DEFICIENCY (*CYP21A2*) **negative**
CONGENITAL ADRENAL INSUFFICIENCY, *CYP11A1*-RELATED (*CYP11A1*) **negative**
CONGENITAL AMEGAKARYOCYTIC THROMBOCYTOPENIA (*MPL*) **negative**
CONGENITAL CHRONIC DIARRHEA (*DGAT1*) **negative**
CONGENITAL DISORDER OF GLYCOSYLATION TYPE 1, *ALG1*-RELATED (*ALG1*) **negative**
CONGENITAL DISORDER OF GLYCOSYLATION, TYPE 1A, *PMM2*-Related (*PMM2*) **negative**
CONGENITAL DISORDER OF GLYCOSYLATION, TYPE 1B (*MPL*) **negative**
CONGENITAL DISORDER OF GLYCOSYLATION, TYPE 1C (*ALG6*) **negative**
CONGENITAL DYSERYTHROPOIETIC ANEMIA TYPE 2 (*SEC23B*) **negative**
CONGENITAL FINNISH NEPHROSIS (*NPHS1*) **negative**
CONGENITAL HYDROCEPHALUS 1 (*CCDC88C*) **negative**
CONGENITAL HYPERINSULINISM, *KCNJ11*-Related (*KCNJ11*) **negative**
CONGENITAL INSENSITIVITY TO PAIN WITH ANHIDROSIS (*CIPA*) (*NTRK1*) **negative**
CONGENITAL MYASTHENIC SYNDROME, *CHAT*-RELATED (*CHAT*) **negative**
CONGENITAL MYASTHENIC SYNDROME, *CHRNE*-RELATED (*CHRNE*) **negative**
CONGENITAL MYASTHENIC SYNDROME, *COLQ*-RELATED (*COLQ*) **negative**
CONGENITAL MYASTHENIC SYNDROME, *DOK7*-RELATED (*DOK7*) **negative**
CONGENITAL MYASTHENIC SYNDROME, *RAPSN*-RELATED (*RAPSN*) **negative**
CONGENITAL NEPHROTIC SYNDROME, *PLCE1*-RELATED (*PLCE1*) **negative**
CONGENITAL NEUTROPENIA, *G6PC3*-RELATED (*G6PC3*) **negative**
CONGENITAL NEUTROPENIA, *HAX1*-RELATED (*HAX1*) **negative**
CONGENITAL NEUTROPENIA, *VPS45*-RELATED (*VPS45*) **negative**
CONGENITAL SECRETORY CHLORIDE DIARRHEA 1 (*SLC26A3*) **negative**
CORNEAL DYSTROPHY AND PERCEPTIVE DEAFNESS (*SLC4A11*) **negative**
CORTICOSTERONE METHYLOXIDASE DEFICIENCY (*CYP11B2*) **negative**
COSTEFF SYNDROME (3-METHYLGUTACONIC ACIDURIA, TYPE 3) (*OPA3*) **negative**
CRB1-RELATED RETINAL DYSTROPHIES (*CRB1*) **negative**
CYSTIC FIBROSIS (*CFTR*) **negative**
CYSTINOSIS (*CTNS*) **negative**
CYTOCHROME C OXIDASE DEFICIENCY, *PET100*-RELATED (*PET100*) **negative**
CYTOCHROME P450 OXIDOREDUCTASE DEFICIENCY (*POR*) **negative**

D

D-BIFUNCTIONAL PROTEIN DEFICIENCY (*HSD17B4*) **negative**
DEAFNESS, AUTOSOMAL RECESSIVE 77 (*LOXHD1*) **negative**
DIHYDROPTERIDINE REDUCTASE (*DHPR*) DEFICIENCY (*QDPR*) **negative**
DONNAI-BARROW SYNDROME (*LRP2*) **negative**
DUBIN-JOHNSON SYNDROME (*ABCC2*) **negative**
DYSKERATOSIS CONGENITA SPECTRUM DISORDERS (*TERT*) **negative**
DYSKERATOSIS CONGENITA, *RTEL1*-RELATED (*RTEL1*) **negative**
DYSTROPHIC EPIDERMOLYSIS BULLOSA, *COL7A1*-Related (*COL7A1*) **negative**

E

EARLY INFANTILE EPILEPTIC ENCEPHALOPATHY, *CAD*-RELATED (*CAD*) **negative**
EHLERS-DANLOS SYNDROME TYPE VI (*PLOD1*) **negative**
EHLERS-DANLOS SYNDROME, CLASSIC-LIKE, *TNXB*-RELATED (*TNXB*) **negative**
EHLERS-DANLOS SYNDROME, TYPE VII C (*ADAMTS2*) **negative**
ELLIS-VAN CREVELD SYNDROME, *EVC2*-RELATED (*EVC2*) **negative**
ELLIS-VAN CREVELD SYNDROME, *EVC*-RELATED (*EVC*) **negative**
ENHANCED S-CONE SYNDROME (*NR2E3*) **negative**
EPIMERASE DEFICIENCY (GALACTOSEMIA TYPE III) (*GALE*) **negative**
EPIPHYSEAL DYSPLASIA, MULTIPLE, 7/DESBUQUOIS DYSPLASIA 1 (*CANT1*) **negative**
ERCC6-RELATED DISORDERS (*ERCC6*) **negative**
ERCC8-RELATED DISORDERS (*ERCC8*) **negative**
ETHYLMALONIC ENCEPHALOPATHY (*ETHE1*) **negative**

F

FACTOR XI DEFICIENCY (*F11*) **negative**

FAMILIAL DYSAUTONOMIA (*IKBKAP*) **negative**

FAMILIAL HEMOPHAGOCYTIC LYMPHOHISTIOCYTOSIS, *PRF1*-RELATED (*PRF1*) **negative**
FAMILIAL HEMOPHAGOCYTIC LYMPHOHISTIOCYTOSIS, *STX11*-RELATED (*STX11*) **negative**
FAMILIAL HEMOPHAGOCYTIC LYMPHOHISTIOCYTOSIS, *STXBP2*-RELATED (*STXBP2*) **negative**
FAMILIAL HEMOPHAGOCYTIC LYMPHOHISTIOCYTOSIS, *UNC13D*-RELATED (*UNC13D*) **negative**
FAMILIAL HYPERCHOLESTEROLEMIA, *LDLRAP1*-RELATED (*LDLRAP1*) **negative**
FAMILIAL HYPERCHOLESTEROLEMIA, *LDLR*-RELATED (*LDLR*) **negative**
FAMILIAL HYPERINSULINISM, *ABCC8*-RELATED (*ABCC8*) **negative**
FAMILIAL NEPHROGENIC DIABETES INSIPIDUS, *AQP2*-RELATED (*AQP2*) **negative**
FANCONI ANEMIA, GROUP A (*FANCA*) **negative**
FANCONI ANEMIA, GROUP C (*FANCC*) **negative**
FANCONI ANEMIA, GROUP D2 (*FANCD2*) **negative**
FANCONI ANEMIA, GROUP E (*FANCE*) **negative**
FANCONI ANEMIA, GROUP F (*FANCF*) **negative**
FANCONI ANEMIA, GROUP G (*FANCG*) **negative**
FANCONI ANEMIA, GROUP I (*FANCI*) **negative**
FANCONI ANEMIA, GROUP J (*BRIP1*) **negative**
FANCONI ANEMIA, GROUP L (*FANCL*) **negative**
FARBER LIPOGRANULOMATOSIS (*ASAH1*) **negative**
FOVEAL HYPOPLASIA (*SLC38A8*) **negative**
FRASER SYNDROME 3, *GRIP1*-RELATED (*GRIP1*) **negative**
FRASER SYNDROME, *FRAS1*-RELATED (*FRAS1*) **negative**
FRASER SYNDROME, *FREM2*-RELATED (*FREM2*) **negative**
FRIEDREICH ATAXIA (*FXN*) **negative**
FRUCTOSE 1,6-BISPHOSPHATASE DEFICIENCY (*FBP1*) **negative**
FUCOSIDOSIS, *FUCA1*-RELATED (*FUCA1*) **negative**
FUMARASE DEFICIENCY (*FH*) **negative**

G

GABA-TRANSAMINASE DEFICIENCY (*ABAT*) **negative**
GALACTOKINASE DEFICIENCY (GALACTOSEMIA, TYPE II) (*GALK1*) **negative**
GALACTOSEMIA (*GALT*) **negative**
GALACTOSIALIDOSIS (*CTSA*) **negative**
GAUCHER DISEASE (*GBA*) **negative**
GCH1-RELATED CONDITIONS (*GCH1*) **negative**
GDF5-RELATED CONDITIONS (*GDF5*) **negative**
GERODERMA OSTEODYSPLASTICA (*GORAB*) **negative**
GITELMAN SYNDROME (*SLC12A3*) **negative**
GLANZMANN THROMBASTHENIA (*ITGB3*) **negative**
GLUTARIC ACIDEMIA, TYPE 1 (*GCDH*) **negative**
GLUTARIC ACIDEMIA, TYPE 2A (*ETFA*) **negative**
GLUTARIC ACIDEMIA, TYPE 2B (*ETFB*) **negative**
GLUTARIC ACIDEMIA, TYPE 2C (*ETFDH*) **negative**
GLUTATHIONE SYNTHETASE DEFICIENCY (*GSS*) **negative**
GLYCINE ENCEPHALOPATHY, *AMT*-RELATED (*AMT*) **negative**
GLYCINE ENCEPHALOPATHY, *GLDC*-RELATED (*GLDC*) **negative**
GLYCOGEN STORAGE DISEASE TYPE 5 (McArdle Disease) (*PYGM*) **negative**
GLYCOGEN STORAGE DISEASE TYPE IXB (*PHKB*) **negative**
GLYCOGEN STORAGE DISEASE TYPE IXC (*PHKG2*) **negative**
GLYCOGEN STORAGE DISEASE, TYPE 1a (*G6PC*) **negative**
GLYCOGEN STORAGE DISEASE, TYPE 1b (*SLC37A4*) **negative**
GLYCOGEN STORAGE DISEASE, TYPE 2 (POMPE DISEASE) (*GAA*) **negative**
GLYCOGEN STORAGE DISEASE, TYPE 3 (*AGL*) **negative**
GLYCOGEN STORAGE DISEASE, TYPE 4 (*GBE1*) **negative**
GLYCOGEN STORAGE DISEASE, TYPE 7 (*PFKM*) **negative**
GRACILE SYNDROME (*BCS1L*) **negative**
GUANIDINOACETATE METHYLTRANSFERASE DEFICIENCY (*GAMT*) **negative**

H

HARLEQUIN ICHTHYOSIS (*ABCA12*) **negative**
HEME OXYGENASE 1 DEFICIENCY (*HMOX1*) **negative**
HEMOCHROMATOSIS TYPE 2A (*HFE2*) **negative**
HEMOCHROMATOSIS, TYPE 3, *TFR2*-Related (*TFR2*) **negative**
HEPATOCEREBRAL MITOCHONDRIAL DNA DEPLETION SYNDROME, *MPV17*-RELATED (*MPV17*) **negative**



Patient Information

Patient Name: Donor 8326

Test Information

Ordering Physician: [REDACTED]



Date Of Birth: [REDACTED]

Case File ID: [REDACTED]

Clinic Information: Fairfax Cryobank

Report Date: 03/04/2026

H

HEREDITARY FRUCTOSE INTOLERANCE (ALDOB) **negative**
 HEREDITARY HEMOCHROMATOSIS TYPE 2B (HAMP) **negative**
 HEREDITARY SPASTIC PARAPARESIS, TYPE 49 (TECPR2) **negative**
 HEREDITARY SPASTIC PARAPLEGIA, CYP7B1-RELATED (CYP7B1) **negative**
 HERMANISKY-PUDLAK SYNDROME, AP3B1-RELATED (AP3B1) **negative**
 HERMANISKY-PUDLAK SYNDROME, BLOC1S3-RELATED (BLOC1S3) **negative**
 HERMANISKY-PUDLAK SYNDROME, BLOC1S6-RELATED (BLOC1S6) **negative**
 HERMANISKY-PUDLAK SYNDROME, HPS1-RELATED (HPS1) **negative**
 HERMANISKY-PUDLAK SYNDROME, HPS3-RELATED (HPS3) **negative**
 HERMANISKY-PUDLAK SYNDROME, HPS4-RELATED (HPS4) **negative**
 HERMANISKY-PUDLAK SYNDROME, HPS5-RELATED (HPS5) **negative**
 HERMANISKY-PUDLAK SYNDROME, HPS6-RELATED (HPS6) **negative**
 HOLOCARBOXYLASE SYNTHETASE DEFICIENCY (HLCS) **negative**
 HOMOCYSTINURIA AND MEGALOBlastic ANEMIA TYPE CBLG (MTR) **negative**
 HOMOCYSTINURIA DUE TO DEFICIENCY OF MTHFR (MTHFR) **negative**
 HOMOCYSTINURIA, CBS-RELATED (CBS) **negative**
 HOMOCYSTINURIA, Type cblE (MTRR) **negative**
 HYDROLETHALUS SYNDROME (HLS1) **negative**
 HYPER-IGM IMMUNODEFICIENCY (CD40) **negative**
 HYPERORNITHINEMIA-HYPERAMMONEMIA-HOMOCITRULLINURIA (HHH SYNDROME) (SLC25A15) **negative**
 HYPERPHOSPHATEMIC FAMILIAL TUMORAL CALCINOSIS, GALNT3-RELATED (GALNT3) **negative**
 HYPOMYELINATING LEUKODYSTROPHY 12 (VPS11) **negative**
 HYPOPHOSPHATASIA, ALPL-RELATED (ALPL) **negative**

I

IMERSLUND-GRASBECK SYNDROME 2 (AMN) **negative**
 IMMUNODEFICIENCY-CENTROMERIC INSTABILITY-FACIAL ANOMALIES (ICF) SYNDROME, DNMT3B-RELATED (DNMT3B) **negative**
 IMMUNODEFICIENCY-CENTROMERIC INSTABILITY-FACIAL ANOMALIES (ICF) SYNDROME, ZBTB24-RELATED (ZBTB24) **negative**
 INCLUSION BODY MYOPATHY 2 (GNE) **negative**
 INFANTILE CEREBRAL AND CEREBELLAR ATROPHY (MED17) **negative**
 INFANTILE NEPHRONOPHTHISIS (INVS) **negative**
 INFANTILE NEUROAXONAL DYSTROPHY (PLA2G6) **negative**
 ISOLATED ECTOPIA LENTIS (ADAMTSL4) **negative**
 ISOLATED SULFITE OXIDASE DEFICIENCY (SUOX) **negative**
 ISOLATED THYROID-STIMULATING HORMONE DEFICIENCY (TSHB) **negative**
 ISOVALERIC ACIDEMIA (IVD) **negative**

J

JOHANSON-BLIZZARD SYNDROME (UBR1) **negative**
 JOUBERT SYNDROME 2 / MECKEL SYNDROME 2 (TMEM216) **negative**
 JOUBERT SYNDROME AND RELATED DISORDERS (JSRD), TMEM67-RELATED (TMEM67) **negative**
 JOUBERT SYNDROME, AHI1-RELATED (AHI1) **negative**
 JOUBERT SYNDROME, ARL13B-RELATED (ARL13B) **negative**
 JOUBERT SYNDROME, B9D1-RELATED (B9D1) **negative**
 JOUBERT SYNDROME, B9D2-RELATED (B9D2) **negative**
 JOUBERT SYNDROME, C2CD3-RELATED/OROFACIODIGITAL SYNDROME 14 (C2CD3) **negative**
 JOUBERT SYNDROME, CC2D2A-RELATED/COACH SYNDROME (CC2D2A) **negative**
 JOUBERT SYNDROME, CEP104-RELATED (CEP104) **negative**
 JOUBERT SYNDROME, CEP120-RELATED/SHORT-RIB THORACIC DYSPLASIA 13 WITH OR WITHOUT POLYDACTYL (CEP120) **negative**
 JOUBERT SYNDROME, CEP41-RELATED (CEP41) **negative**
 JOUBERT SYNDROME, CPLANE1-RELATED / OROFACIODIGITAL SYNDROME 6 (CPLANE1) **negative**
 JOUBERT SYNDROME, CSPP1-RELATED (CSPP1) **negative**
 JOUBERT SYNDROME, INPP5E-RELATED (INPP5E) **negative**
 JUNCTIONAL EPIDERMOLYSIS BULLOSA, COL17A1-RELATED (COL17A1) **negative**
 JUNCTIONAL EPIDERMOLYSIS BULLOSA, ITGA6-RELATED (ITGA6) **negative**
 JUNCTIONAL EPIDERMOLYSIS BULLOSA, ITGB4-RELATED (ITGB4) **negative**
 JUNCTIONAL EPIDERMOLYSIS BULLOSA, LAMB3-RELATED (LAMB3) **negative**
 JUNCTIONAL EPIDERMOLYSIS BULLOSA, LAMC2-RELATED (LAMC2) **negative**
 JUNCTIONAL EPIDERMOLYSIS BULLOSA/LARYNGOONYCHOCUTANEOUS SYNDROME, LAMA3-RELATED (LAMA3) **negative**

K

KRABBE DISEASE (GALC) **negative**

L

LAMELLAR ICHTHYOSIS, TYPE 1 (TGM1) **negative**
 LARON SYNDROME (GHR) **negative**
 LEBER CONGENITAL AMAUROSIS 2 (RPE65) **negative**
 LEBER CONGENITAL AMAUROSIS TYPE AIPL1 (AIPL1) **negative**
 LEBER CONGENITAL AMAUROSIS TYPE GUCY2D (GUCY2D) **negative**
 LEBER CONGENITAL AMAUROSIS TYPE TULP1 (TULP1) **negative**
 LEBER CONGENITAL AMAUROSIS, IQCB1-RELATED/SENIOR-LOKEN SYNDROME 5 (IQCB1) **negative**
 LEBER CONGENITAL AMAUROSIS, TYPE CEP290 (CEP290) **negative**
 LEBER CONGENITAL AMAUROSIS, TYPE LCA5 (LCA5) **negative**
 LEBER CONGENITAL AMAUROSIS, TYPE RDH12 (RDH12) **negative**
 LEIGH SYNDROME, FRENCH-CANADIAN TYPE (LRPPRC) **negative**
 LETHAL CONGENITAL CONTRACTURE SYNDROME 1 (GLE1) **negative**
 LEUKOENCEPHALOPATHY WITH VANISHING WHITE MATTER (EIF2B5) **negative**
 LEUKOENCEPHALOPATHY WITH VANISHING WHITE MATTER, EIF2B1-RELATED (EIF2B1) **negative**
 LEUKOENCEPHALOPATHY WITH VANISHING WHITE MATTER, EIF2B2-RELATED (EIF2B2) **negative**
 LEUKOENCEPHALOPATHY WITH VANISHING WHITE MATTER, EIF2B3-RELATED (EIF2B3) **negative**
 LEUKOENCEPHALOPATHY WITH VANISHING WHITE MATTER, EIF2B4-RELATED (EIF2B4) **negative**
 LIG4 SYNDROME (LIG4) **negative**
 LIMB-GIRDLE MUSCULAR DYSTROPHY TYPE 8 (TRIM32) **negative**
 LIMB-GIRDLE MUSCULAR DYSTROPHY, TYPE 2A (CAPN3) **negative**
 LIMB-GIRDLE MUSCULAR DYSTROPHY, TYPE 2B (DYSF) **negative**
 LIMB-GIRDLE MUSCULAR DYSTROPHY, TYPE 2C (SGCG) **negative**
 LIMB-GIRDLE MUSCULAR DYSTROPHY, TYPE 2D (SGCA) **negative**
 LIMB-GIRDLE MUSCULAR DYSTROPHY, TYPE 2E (SGCB) **negative**
 LIMB-GIRDLE MUSCULAR DYSTROPHY, TYPE 2F (SGCD) **negative**
 LIMB-GIRDLE MUSCULAR DYSTROPHY, TYPE 2I (FKRP) **negative**
 LIPOAMIDE DEHYDROGENASE DEFICIENCY (DIHYDROLIPOAMIDE DEHYDROGENASE DEFICIENCY) (DLD) **negative**
 LIPOID ADRENAL HYPERPLASIA (STAR) **negative**
 LIPOPROTEIN LIPASE DEFICIENCY (LPL) **negative**
 LONG CHAIN 3-HYDROXYACYL-CoA DEHYDROGENASE DEFICIENCY (HADHA) **negative**
 LRAT-RELATED CONDITIONS (LRAT) **negative**
 LUNG DISEASE, IMMUNODEFICIENCY, AND CHROMOSOME BREAKAGE SYNDROME (LICS) (NSMCE3) **negative**
 LYSINURIC PROTEIN INTOLERANCE (SLC7A7) **negative**

M

MALONYL-CoA DECARBOXYLASE DEFICIENCY (MLYCD) **negative**
 MAPLE SYRUP URINE DISEASE, TYPE 1A (BCKDHA) **negative**
 MAPLE SYRUP URINE DISEASE, TYPE 1B (BCKDHB) **negative**
 MAPLE SYRUP URINE DISEASE, TYPE 2 (DBT) **negative**
 MCKUSICK-KAUFMAN SYNDROME (MKKS) **negative**
 MECKEL SYNDROME 7/NEPHRONOPHTHISIS 3 (NPHP3) **negative**
 MECKEL-GRUBER SYNDROME, TYPE 1 (MKS1) **negative**
 MECP-RELATED NEUROLOGIC DISORDER (MECP) **negative**
 MEDIUM CHAIN ACYL-CoA DEHYDROGENASE DEFICIENCY (ACADM) **negative**
 MEDNIK SYNDROME (AP1S1) **negative**
 MEGALENCEPHALIC LEUKOENCEPHALOPATHY WITH SUBCORTICAL CYSTS (MLC1) **negative**
 MEROSIN-DEFICIENT MUSCULAR DYSTROPHY (LAMA2) **negative**
 METABOLIC ENCEPHALOPATHY AND ARRHYTHMIAS, TANGO2-RELATED (TANGO2) **negative**
 METACHROMATIC LEUKODYSTROPHY, ARSA-RELATED (ARSA) **negative**
 METACHROMATIC LEUKODYSTROPHY, PSAP-RELATED (PSAP) **negative**
 METHYLMALONIC ACIDEMIA AND HOMOCYSTINURIA TYPE CBLF (LMBRD1) **negative**
 METHYLMALONIC ACIDEMIA, MCEE-RELATED (MCEE) **negative**
 METHYLMALONIC ACIDURIA AND HOMOCYSTINURIA, TYPE CBLF (MMACHC) **negative**
 METHYLMALONIC ACIDURIA AND HOMOCYSTINURIA, TYPE CblD (MMADHC) **negative**



Patient Information

Patient Name: Donor 8326

Test Information

Ordering Physician:



Clinic Information: Fairfax Cryobank

Date Of Birth:

Case File ID:

Report Date: 03/04/2026

M

METHYLMALONIC ACIDURIA, MMAA-RELATED (MMAA) **negative**
 METHYLMALONIC ACIDURIA, MMAB-RELATED (MMAB) **negative**
 METHYLMALONIC ACIDURIA, TYPE MUT (O) (MUT) **negative**
 MEVALONIC KINASE DEFICIENCY (MVK) **negative**
 MICROCEPHALIC OSTEODYSPLASTIC PRIMORDIAL DWARFISM TYPE II (PCNT) **negative**
 MICROPHTHALMIA / ANOPHTHALMIA, VSX2-RELATED (VSX2) **negative**
 MITOCHONDRIAL COMPLEX 1 DEFICIENCY, ACAD9-RELATED (ACAD9) **negative**
 MITOCHONDRIAL COMPLEX 1 DEFICIENCY, NDUFAF5-RELATED (NDUFAF5) **negative**
 MITOCHONDRIAL COMPLEX 1 DEFICIENCY, NDUFS6-RELATED (NDUFS6) **negative**
 MITOCHONDRIAL COMPLEX I DEFICIENCY, NUCLEAR TYPE 1 (NDUFS4) **negative**
 MITOCHONDRIAL COMPLEX I DEFICIENCY, NUCLEAR TYPE 10 (NDUFAF2) **negative**
 MITOCHONDRIAL COMPLEX I DEFICIENCY, NUCLEAR TYPE 17 (NDUFAF6) **negative**
 MITOCHONDRIAL COMPLEX I DEFICIENCY, NUCLEAR TYPE 19 (FOXRED1) **negative**
 MITOCHONDRIAL COMPLEX I DEFICIENCY, NUCLEAR TYPE 3 (NDUFS7) **negative**
 MITOCHONDRIAL COMPLEX I DEFICIENCY, NUCLEAR TYPE 4 (NDUFV1) **negative**
 MITOCHONDRIAL COMPLEX IV DEFICIENCY, NUCLEAR TYPE 2, SCO2-RELATED (SCO2) **negative**
 MITOCHONDRIAL COMPLEX IV DEFICIENCY, NUCLEAR TYPE 6 (COX15) **negative**
 MITOCHONDRIAL DNA DEPLETION SYNDROME 2 (TK2) **negative**
 MITOCHONDRIAL DNA DEPLETION SYNDROME 3 (DGUOK) **negative**
 MITOCHONDRIAL MYOPATHY AND SIDEROBLASTIC ANEMIA (MLASA1) (PUS1) **negative**
 MITOCHONDRIAL TRIFUNCTIONAL PROTEIN DEFICIENCY, HADHB-RELATED (HADHB) **negative**
 MOLYBDENUM COFACTOR DEFICIENCY TYPE B (MOCS2) **negative**
 MOLYBDENUM COFACTOR DEFICIENCY, TYPE A (MOCS1) **negative**
 MUCOLIPIDOSIS II/III A (GNPTAB) **negative**
 MUCOLIPIDOSIS III GAMMA (GNPTG) **negative**
 MUCOLIPIDOSIS, TYPE IV (MCOLN1) **negative**
 MUCOPOLYSACCHARIDOSIS, TYPE I (HURLER SYNDROME) (IDUA) **negative**
 MUCOPOLYSACCHARIDOSIS, TYPE III A (SANFILIPPO A) (SGSH) **see first page**
 MUCOPOLYSACCHARIDOSIS, TYPE III B (SANFILIPPO B) (NAGLU) **negative**
 MUCOPOLYSACCHARIDOSIS, TYPE III C (SANFILIPPO C) (HGSNAT) **negative**
 MUCOPOLYSACCHARIDOSIS, TYPE III D (SANFILIPPO D) (GNS) **negative**
 MUCOPOLYSACCHARIDOSIS, TYPE IV A (MORQUIO SYNDROME) (GALNS) **negative**
 MUCOPOLYSACCHARIDOSIS, TYPE IV B/GM1 GANGLIOSIDOSIS (GLB1) **negative**
 MUCOPOLYSACCHARIDOSIS, TYPE IX (HYAL1) **negative**
 MUCOPOLYSACCHARIDOSIS, TYPE VI (MAROTEAUX-LAMY) (ARSB) **negative**
 MUCOPOLYSACCHARIDOSIS, TYPE VII (GUSB) **negative**
 MULIBREY NANISM (TRIM37) **negative**
 MULTIPLE PTERYGIUM SYNDROME, CHRNG-RELATED/ESCOBAR SYNDROME (CHRN) **negative**
 MULTIPLE SULFATASE DEFICIENCY (SUMF1) **negative**
 MUSCLE-EYE-BRAIN DISEASE, POMGNT1-RELATED (POMGNT1) **negative**
 MUSCULAR DYSTROPHY-DYSTROGLYCANOPATHY (RXYL1) **negative**
 MUSK-RELATED CONGENITAL MYASTHENIC SYNDROME (MUSK) **negative**
 MYONEUROGASTROINTESTINAL ENCEPHALOPATHY (MNGIE) (TYMP) **negative**
 MYOTONIA CONGENITA (CLCN1) **negative**

N

N-ACETYLGLUTAMATE SYNTHASE DEFICIENCY (NAGS) **negative**
 NEMALINE MYOPATHY, NEB-RELATED (NEB) **negative**
 NEPHRONOPHTHISIS 1 (NPHP1) **negative**
 NEURONAL CEROID LIPOFUSCINOSIS, CLN5-RELATED (CLN5) **negative**
 NEURONAL CEROID LIPOFUSCINOSIS, CLN6-RELATED (CLN6) **negative**
 NEURONAL CEROID LIPOFUSCINOSIS, CLN8-RELATED (CLN8) **negative**
 NEURONAL CEROID LIPOFUSCINOSIS, MFSDB-RELATED (MFSDB) **negative**
 NEURONAL CEROID LIPOFUSCINOSIS, PPT1-RELATED (PPT1) **negative**
 NEURONAL CEROID LIPOFUSCINOSIS, TPP1-RELATED (TPP1) **negative**
 NGLY1-CONGENITAL DISORDER OF GLYCOSYLATION (NGLY1) **negative**
 NIEMANN-PICK DISEASE, TYPE C1 / D (NPC1) **negative**
 NIEMANN-PICK DISEASE, TYPE C2 (NPC2) **negative**
 NIEMANN-PICK DISEASE, TYPES A / B (SMPD1) **negative**
 NIJMEGEN BREAKAGE SYNDROME (NBN) **negative**
 NON-SYNDROMIC HEARING LOSS, GJB2-RELATED (GJB2) **negative**
 NON-SYNDROMIC HEARING LOSS, MYO15A-RELATED (MYO15A) **negative**
 NONSYNDROMIC HEARING LOSS, OTOA-RELATED (OTOA) **negative**

NONSYNDROMIC HEARING LOSS, OTOF-RELATED (OTOF) **negative**
 NONSYNDROMIC HEARING LOSS, PJVK-RELATED (PJVK) **negative**
 NONSYNDROMIC HEARING LOSS, SYNE4-RELATED (SYNE4) **negative**
 NONSYNDROMIC HEARING LOSS, TMC1-RELATED (TMC1) **negative**
 NONSYNDROMIC HEARING LOSS, TMPS3-RELATED (TMPS3) **negative**
 NONSYNDROMIC INTELLECTUAL DISABILITY (CC2D1A) **negative**
 NORMOPHOSPHATEMIC TUMORAL CALCINOSIS (SAMD9) **negative**

O

OCULOCUTANEOUS ALBINISM TYPE III (TYRP1) **negative**
 OCULOCUTANEOUS ALBINISM TYPE IV (SLC45A2) **negative**
 OCULOCUTANEOUS ALBINISM, OCA2-RELATED (OCA2) **negative**
 OCULOCUTANEOUS ALBINISM, TYPES 1A AND 1B (TYR) **negative**
 ODONTO-ONYCHO-DERMAL DYSPLASIA / SCHOPF-SCHULZ-PASSARGE SYNDROME (WNT10A) **negative**
 OMENN SYNDROME, RAG2-RELATED (RAG2) **negative**
 ORNITHINE AMINOTRANSFERASE DEFICIENCY (OAT) **negative**
 OSTEOGENESIS IMPERFECTA TYPE VII (CRTAP) **negative**
 OSTEOGENESIS IMPERFECTA TYPE VIII (P3H1) **negative**
 OSTEOGENESIS IMPERFECTA TYPE XI (FKBP10) **negative**
 OSTEOGENESIS IMPERFECTA TYPE XIII (BMP1) **negative**
 OSTEOPETROSIS, INFANTILE MALIGNANT, TCIRG1-RELATED (TCIRG1) **negative**
 OSTEOPETROSIS, OSTM1-RELATED (OSTM1) **negative**

P

PANTOTHENATE KINASE-ASSOCIATED NEURODEGENERATION (PANK2) **negative**
 PAPILLON LEFEVRE SYNDROME (CTSC) **negative**
 PARKINSON DISEASE 15 (FBXO7) **negative**
 PENDRED SYNDROME (SLC26A4) **negative**
 PERLMAN SYNDROME (DIS3L2) **negative**
 PGM3-CONGENITAL DISORDER OF GLYCOSYLATION (PGM3) **negative**
 PHENYLKETONURIA (PAH) **negative**
 PIGN-CONGENITAL DISORDER OF GLYCOSYLATION (PIGN) **see first page**
 PITUITARY HORMONE DEFICIENCY, COMBINED 3 (LHX3) **negative**
 POLG-RELATED DISORDERS (POLG) **negative**
 POLYCYSTIC KIDNEY DISEASE, AUTOSOMAL RECESSIVE (PKHD1) **negative**
 PONTOCEREBELLAR HYPOPLASIA, EXOSC3-RELATED (EXOSC3) **negative**
 PONTOCEREBELLAR HYPOPLASIA, RARS2-RELATED (RARS2) **negative**
 PONTOCEREBELLAR HYPOPLASIA, TSEN2-RELATED (TSEN2) **negative**
 PONTOCEREBELLAR HYPOPLASIA, TSEN54-RELATED (TSEN54) **negative**
 PONTOCEREBELLAR HYPOPLASIA, TYPE 1A (VRK1) **negative**
 PONTOCEREBELLAR HYPOPLASIA, TYPE 2D (SEPS3) **negative**
 PONTOCEREBELLAR HYPOPLASIA, VPS53-RELATED (VPS53) **negative**
 PRIMARY CILIARY DYSKINESIA, CCDC103-RELATED (CCDC103) **negative**
 PRIMARY CILIARY DYSKINESIA, CCDC39-RELATED (CCDC39) **negative**
 PRIMARY CILIARY DYSKINESIA, DNAH11-RELATED (DNAH11) **negative**
 PRIMARY CILIARY DYSKINESIA, DNAH5-RELATED (DNAH5) **negative**
 PRIMARY CILIARY DYSKINESIA, DNAI1-RELATED (DNAI1) **negative**
 PRIMARY CILIARY DYSKINESIA, DNAI2-RELATED (DNAI2) **negative**
 PRIMARY CONGENITAL GLAUCOMA/PETERS ANOMALY (CYP1B1) **negative**
 PRIMARY HYPEROXALURIA, TYPE 1 (AGXT) **negative**
 PRIMARY HYPEROXALURIA, TYPE 2 (GRHPR) **negative**
 PRIMARY HYPEROXALURIA, TYPE 3 (HOGA1) **negative**
 PRIMARY MICROCEPHALY 1, AUTOSOMAL RECESSIVE (MCPH1) **negative**
 PROGRESSIVE EARLY-ONSET ENCEPHALOPATHY WITH BRAIN ATROPHY AND THIN CORPUS CALLOSUM (TBCD) **negative**
 PROGRESSIVE FAMILIAL INTRAHEPATIC CHOLESTASIS, ABCB4-RELATED (ABCB4) **negative**
 PROGRESSIVE FAMILIAL INTRAHEPATIC CHOLESTASIS, TYPE 1 (PFIC1) (ATP8B1) **negative**
 PROGRESSIVE FAMILIAL INTRAHEPATIC CHOLESTASIS, TYPE 2 (ABCB11) **negative**
 PROGRESSIVE FAMILIAL INTRAHEPATIC CHOLESTASIS, TYPE 4 (PFIC4) (TJP2) **negative**
 PROGRESSIVE PSEUDORHEUMATOID DYSPLASIA (CCN6) **negative**
 PROLIDASE DEFICIENCY (PEPD) **negative**
 PROPIONIC ACIDEMIA, PCCA-RELATED (PCCA) **negative**
 PROPIONIC ACIDEMIA, PCCB-RELATED (PCCB) **negative**
 PSEUDOXANTHOMA ELASTICUM (ABCC6) **negative**



Patient Information

Patient Name: Donor 8326

Test Information

Ordering Physician: [REDACTED]



Clinic Information: Fairfax Cryobank

Date Of Birth: [REDACTED]

Case File ID: [REDACTED]

Report Date: 03/04/2026

P

PTERIN-4 ALPHA-CARBINOLAMINE DEHYDRATASE (PCD) DEFICIENCY (*PCBD1*) **negative**
PYCNODYSTOSIS (*CTSK*) **negative**
PYRIDOXAL 5'-PHOSPHATE-DEPENDENT EPILEPSY (*PNPO*) **negative**
PYRIDOXINE-DEPENDENT EPILEPSY (*ALDH7A1*) **negative**
PYRUVATE CARBOXYLASE DEFICIENCY (*PC*) **negative**
PYRUVATE DEHYDROGENASE DEFICIENCY, PDHB-RELATED (*PDHB*) **negative**

R

REFSUM DISEASE, PHYH-RELATED (*PHYH*) **negative**
RENAL TUBULAR ACIDOSIS AND DEAFNESS, ATP6V1B1-RELATED (*ATP6V1B1*) **negative**
RENAL TUBULAR ACIDOSIS, PROXIMAL, WITH OCULAR ABNORMALITIES AND MENTAL RETARDATION (*SLC4A4*) **negative**
RETINITIS PIGMENTOSA 25 (*EYS*) **negative**
RETINITIS PIGMENTOSA 26 (*CERKL*) **negative**
RETINITIS PIGMENTOSA 28 (*FAM161A*) **negative**
RETINITIS PIGMENTOSA 36 (*PRCD*) **negative**
RETINITIS PIGMENTOSA 59 (*DHDDS*) **negative**
RETINITIS PIGMENTOSA 62 (*MAK*) **negative**
RHIZOMELIC CHONDRODYSPLASIA PUNCTATA, TYPE 1 (*PEX7*) **negative**
RHIZOMELIC CHONDRODYSPLASIA PUNCTATA, TYPE 2 (*GNPAT*) **negative**
RHIZOMELIC CHONDRODYSPLASIA PUNCTATA, TYPE 3 (*AGPS*) **negative**
RLBP1-RELATED RETINOPATHY (*RLBP1*) **negative**
ROBERTS SYNDROME (*ESCO2*) **negative**
RYSR1-RELATED CONDITIONS (*RYSR1*) **negative**

S

SALLA DISEASE (*SLC17A5*) **see first page**
SANDHOFF DISEASE (*HEXB*) **negative**
SCHIMKE IMMUNOSKELETAL DYSPLASIA (*SMARCA1*) **negative**
SCHINDLER DISEASE (*NAGA*) **negative**
SEGAWA SYNDROME, TH-RELATED (*TH*) **negative**
SENIOR-LOKEN SYNDROME 4/NEPHRONOPHTHISIS 4 (*NPHP4*) **negative**
SEPIAPTERIN REDUCTASE DEFICIENCY (*SPR*) **negative**
SEVERE COMBINED IMMUNODEFICIENCY (*SCID*), CD3D-RELATED (*CD3D*) **negative**
SEVERE COMBINED IMMUNODEFICIENCY (*SCID*), CD3E-RELATED (*CD3E*) **negative**
SEVERE COMBINED IMMUNODEFICIENCY (*SCID*), FOXP1-RELATED (*FOXP1*) **negative**
SEVERE COMBINED IMMUNODEFICIENCY (*SCID*), IKBK-RELATED (*IKBK*) **negative**
SEVERE COMBINED IMMUNODEFICIENCY (*SCID*), IL7R-RELATED (*IL7R*) **negative**
SEVERE COMBINED IMMUNODEFICIENCY (*SCID*), JAK3-RELATED (*JAK3*) **negative**
SEVERE COMBINED IMMUNODEFICIENCY (*SCID*), PTPRC-RELATED (*PTPRC*) **negative**
SEVERE COMBINED IMMUNODEFICIENCY (*SCID*), RAG1-RELATED (*RAG1*) **negative**
SEVERE COMBINED IMMUNODEFICIENCY, ADA-Related (*ADA*) **negative**
SEVERE COMBINED IMMUNODEFICIENCY, TYPE ATHABASKAN (*DCLRE1C*) **negative**
SHORT-RIB THORACIC DYSPLASIA 3 WITH OR WITHOUT POLYDACTYLY (*DYNC2H1*) **negative**
SHWACHMAN-DIAMOND SYNDROME, SBDS-RELATED (*SBDS*) **negative**
SIALIDOSIS (*NEU1*) **negative**
SJOGREN-LARSSON SYNDROME (*ALDH3A2*) **negative**
SMITH-LEMLI-OPITZ SYNDROME (*DHCR7*) **negative**
SPASTIC PARAPLEGIA, TYPE 15 (*ZFYVE26*) **negative**
SPASTIC TETRAPLEGIA, THIN CORPUS CALLOSUM, AND PROGRESSIVE MICROCEPHALY (*SPATCCM*) (*SLC1A4*) **negative**
SPG11-RELATED CONDITIONS (*SPG11*) **negative**
SPINAL MUSCULAR ATROPHY (*SMN1*) **negative** *SMN1*: Two copies; g.27134T>G: absent; the absence of the g.27134T>G variant decreases the chance to be a silent (2+0) carrier.
SPINAL MUSCULAR ATROPHY WITH RESPIRATORY DISTRESS TYPE 1 (*IGHMBP2*) **negative**
SPINOCEREBELLAR ATAXIA, AUTOSOMAL RECESSIVE 10 (*ANO10*) **negative**
SPINOCEREBELLAR ATAXIA, AUTOSOMAL RECESSIVE 12 (*WWOX*) **negative**
SPONDYLOCOSTAL DYSOSTOSIS 1 (*DLL3*) **negative**
SPONDYLOTHORACIC DYSOSTOSIS, MESP2-Related (*MESP2*) **negative**
STEEL SYNDROME (*COL27A1*) **negative**
STEROID-RESISTANT NEPHROTIC SYNDROME (*NPHS2*) **negative**
STUVE-WIEDEMANN SYNDROME (*LIFR*) **negative**
SURF1-RELATED CONDITIONS (*SURF1*) **negative**

SURFACTANT DYSFUNCTION, ABCA3-RELATED (*ABCA3*) **negative****T**

TAY-SACHS DISEASE (*HEXA*) **negative**
TBCE-RELATED CONDITIONS (*TBCE*) **negative**
THIAMINE-RESPONSIVE MEGALOBlastic ANEMIA SYNDROME (*SLC19A2*) **negative**
THYROID DYSHORMONOGENESIS 1 (*SLC5A5*) **negative**
THYROID DYSHORMONOGENESIS 2A (*TPO*) **negative**
THYROID DYSHORMONOGENESIS 3 (*TG*) **negative**
THYROID DYSHORMONOGENESIS 6 (*DUOX2*) **negative**
TRANSCOBALAMIN II DEFICIENCY (*TCN2*) **negative**
TRICHOHEPATOENTERIC SYNDROME, SKIC2-RELATED (*SKIC2*) **negative**
TRICHOHEPATOENTERIC SYNDROME, TTC37-RELATED (*TTC37*) **negative**
TRICHOHYDROSTROPHY 1/XERODERMA PIGMENTOSUM, GROUP D (*ERCC2*) **negative**
TRIMETHYLAMINURIA (*FMO3*) **negative**
TRIPLE A SYNDROME (*AAA5*) **negative**
TSHR-RELATED CONDITIONS (*TSHR*) **negative**
TYROSINEMIA TYPE III (*HPD*) **negative**
TYROSINEMIA, TYPE 1 (*FAH*) **negative**
TYROSINEMIA, TYPE 2 (*TAT*) **negative**

U

USHER SYNDROME, TYPE 1B (*MYO7A*) **negative**
USHER SYNDROME, TYPE 1C (*USH1C*) **negative**
USHER SYNDROME, TYPE 1D (*CDH23*) **negative**
USHER SYNDROME, TYPE 1F (*PCDH15*) **negative**
USHER SYNDROME, TYPE 1J/DEAFNESS, AUTOSOMAL RECESSIVE, 48 (*CIB2*) **negative**
USHER SYNDROME, TYPE 2A (*USH2A*) **negative**
USHER SYNDROME, TYPE 2C (*ADGRV1*) **negative**
USHER SYNDROME, TYPE 3 (*CLRN1*) **negative**

V

VERY LONG-CHAIN ACYL-CoA DEHYDROGENASE DEFICIENCY (*ACADVL*) **negative**
VICI SYNDROME (*EPG5*) **negative**
VITAMIN D-DEPENDENT RICKETS, TYPE 1A (*CYP27B1*) **negative**
VITAMIN D-RESISTANT RICKETS TYPE 2A (*VDR*) **negative**
VLDLR-ASSOCIATED CEREBELLAR HYPOPLASIA (*VLDLR*) **negative**

W

WALKER-WARBURG SYNDROME, CRPPA-RELATED (*CRPPA*) **negative**
WALKER-WARBURG SYNDROME, FKTN-RELATED (*FKTN*) **negative**
WALKER-WARBURG SYNDROME, LARGE1-RELATED (*LARGE1*) **negative**
WALKER-WARBURG SYNDROME, POMT1-RELATED (*POMT1*) **negative**
WALKER-WARBURG SYNDROME, POMT2-RELATED (*POMT2*) **negative**
WARSAW BREAKAGE SYNDROME (*DDX11*) **negative**
WERNER SYNDROME (*WRN*) **negative**
WILSON DISEASE (*ATP7B*) **negative**
WOLCOTT-RALLISON SYNDROME (*EIF2AK3*) **negative**
WOLMAN DISEASE (*LIPA*) **negative**
WOODHOUSE-SAKATI SYNDROME (*DCAF17*) **negative**

X

XERODERMA PIGMENTOSUM VARIANT TYPE (*POLH*) **negative**
XERODERMA PIGMENTOSUM, GROUP A (*XPA*) **negative**
XERODERMA PIGMENTOSUM, GROUP C (*XPC*) **see first page**

Z

ZELLWEGER SPECTRUM DISORDER, PEX13-RELATED (*PEX13*) **negative**
ZELLWEGER SPECTRUM DISORDER, PEX16-RELATED (*PEX16*) **negative**
ZELLWEGER SPECTRUM DISORDER, PEX5-RELATED (*PEX5*) **negative**
ZELLWEGER SPECTRUM DISORDERS, PEX10-RELATED (*PEX10*) **negative**
ZELLWEGER SPECTRUM DISORDERS, PEX12-RELATED (*PEX12*) **negative**
ZELLWEGER SPECTRUM DISORDERS, PEX1-RELATED (*PEX1*) **negative**
ZELLWEGER SPECTRUM DISORDERS, PEX26-RELATED (*PEX26*) **negative**
ZELLWEGER SPECTRUM DISORDERS, PEX2-RELATED (*PEX2*) **negative**



Patient Information

Patient Name: Donor 8326

Test Information

Ordering Physician: [REDACTED]



Horizon™
Advanced carrier screening

Date Of Birth: [REDACTED]

Case File ID: [REDACTED]

Clinic Information: Fairfax Cryobank

Report Date: 03/04/2026

Z

ZELLWEGER SPECTRUM DISORDERS, PEX6-RELATED (PEX6) **negative**



Patient Information

Patient Name: Donor 8326

Test Information

Ordering Physician: [REDACTED]

**Horizon™**
Advanced carrier screening

Date Of Birth: [REDACTED]

Case File ID: [REDACTED]

Clinic Information: Fairfax Cryobank

Report Date: 03/04/2026

Testing Methodology, Limitations, and Comments:**Next-generation sequencing (NGS)**

Sequencing library prepared from genomic DNA isolated from a patient sample is enriched for targets of interest using standard hybridization capture protocols and PCR amplification (for targets specified below). NGS is then performed to achieve the standards of quality control metrics, including a minimum coverage of 99% of targeted regions at 20X sequencing depth. Sequencing data is aligned to human reference sequence, followed by deduplication, metric collection and variant calling (coding region +/- 20bp). Variants are then classified according to ACMGG/AMP standards of interpretation using publicly available databases including but not limited to ENSEMBL, HGMD Pro, ClinGen, ClinVar, 1000G, ESP and gnomAD. Variants predicted to be pathogenic or likely pathogenic for the specified diseases are reported. It should be noted that the data interpretation is based on our current understanding of the genes and variants at the time of reporting. Putative positive sequencing variants that do not meet internal quality standards or are within highly homologous regions are confirmed by Sanger sequencing or gene-specific long-range PCR as needed prior to reporting.

Copy Number Variant (CNV) analysis is limited to deletions involving two or more exons for all genes on the panel, in addition to specific known recurrent single-exon deletions. CNVs of small size may have reduced detection rate. This method does not detect gene inversions, single-exonic and sub-exonic deletions (unless otherwise specified), and duplications of all sizes (unless otherwise specified). Additionally, this method does not define the exact breakpoints of detected CNV events. Confirmation testing for copy number variation is performed by specific PCR, Multiplex Ligation-dependent Probe Amplification (MLPA), next generation sequencing, or other methodology.

This test may not detect certain variants due to local sequence characteristics, high/low genomic complexity, homologous sequence, or allele dropout (PCR-based assays). Variants within noncoding regions (promoter, 5'UTR, 3'UTR, deep intronic regions, unless otherwise specified), small deletions or insertions larger than 25bp, low-level mosaic variants, structural variants such as inversions, and/or balanced translocations may not be detected with this technology.

SPECIAL NOTES

For ABCC6, sequencing variants in exons 1-7 are not detected due to the presence of regions of high homology.

For CFTR, when the CFTR R117H variant is detected, reflex analysis of the polythymidine variations (5T, 7T and 9T) at the intron 9 branch/acceptor site of the CFTR gene will be performed. Multi-exon duplication analysis is included.

For CYP21A2, targets were enriched using long-range PCR amplification, followed by next generation sequencing. Duplication analysis will only be performed and reported when c.955C>T (p.Q319*) is detected. Sequencing and CNV analysis may have reduced sensitivity, if variants result from complex rearrangements, in trans with a gene deletion, or CYP21A2 gene duplication on one chromosome and deletion on the other chromosome. This analysis cannot detect sequencing variants located on the CYP21A2 duplicated copy.

For DDX11, sequencing variants in exons 7-11 and CNV for the entire gene are not analyzed due to high sequence homology.

For GJB2, CNV analysis of upstream deletions of GJB6-CRYL1 critical region is included.

For HBA1/HBA2, CNV analysis is offered to detect common deletions of -alpha3.7, -alpha4.2, --MED, --SEA, --FIL, --THAI, --alpha20.5, and/or HS-40. Sequencing and CNV analysis may have reduced sensitivity due to high sequence homology.

For OTOA, sequencing variants in exons 25-29 and CNV in exons 21-29 are not analyzed due to high sequence homology.

For RPGRIP1L, variants in exon 23 are not detected due to assay limitation.

For SAMD9, only p.K1495E variant will be analyzed and reported.



Patient Information

Patient Name: Donor 8326

Test Information

Ordering Physician: [REDACTED]



Date Of Birth: [REDACTED]

Case File ID: [REDACTED]

Clinic Information: Fairfax Cryobank

Report Date: 03/04/2026

Friedreich Ataxia (FXN)

The GAA repeat region of the FXN gene is assessed by trinucleotide PCR assay and capillary electrophoresis. Variances of +/-1 repeat for normal alleles and up to +/-3 repeats for premutation alleles may occur. For fully penetrant expanded alleles, the precise repeat size cannot be determined, therefore the approximate allele size is reported. Sequencing and copy number variants are analyzed by next-generation sequencing analysis.

Friedreich Ataxia Repeat Categories

Categories	GAA Repeat Sizes
Normal	<34
Premutation	34 - 65
Full	>65

Spinal Muscular Atrophy (SMN1)

The total combined copy number of SMN1 and SMN2 exon 7 is quantified based on NGS read depth. The ratio of SMN1 to SMN2 is calculated based on the read depth of a single nucleotide that distinguishes these two genes in exon 7. In addition to copy number analysis, testing for the presence or absence of a single nucleotide polymorphism (g.27134T>G in intron 7 of SMN1) associated with the presence of a SMN1 duplication allele is performed using NGS.

Ethnicity	Two SMN1 copies carrier risk before g.27134T>G testing	Carrier risk after g.27134T>G testing	
		g.27134T>G ABSENT	g.27134T>G PRESENT
Caucasian	1 in 632	1 in 769	1 in 29
Ashkenazi Jewish	1 in 350	1 in 580	LIKELY CARRIER
Asian	1 in 628	1 in 702	LIKELY CARRIER
African-American	1 in 121	1 in 396	1 in 34
Hispanic	1 in 1061	1 in 1762	1 in 140

Variant Classification

Only pathogenic or likely pathogenic variants are reported. Other variants including benign variants, likely benign variants, variants of uncertain significance, or inconclusive variants identified during this analysis may be reported in certain circumstances. Our laboratory's variant classification criteria are based on the ACMG and internal guidelines and our current understanding of the specific genes. This interpretation may change over time as more information about a gene and/or variant becomes available. Natera and its lab partner(s) may reclassify variants at certain intervals but may not release updated reports without a specific request made to Natera by the ordering provider. Natera may disclose incidental findings if deemed clinically pertinent to the test performed.

Negative Results

A negative carrier screening result reduces the risk for a patient to be a carrier of a specific disease but does not completely rule out carrier status. Please visit <https://www.natera.com/panel-option/h-all/> for a table of carrier rates, detection rates, residual risks and promised variants/exons per gene. Carrier rates before and after testing vary by ethnicity and assume a negative family history for each disease screened and the absence of clinical symptoms in the patient. Any patient with a family history for a specific genetic disease will have a higher carrier risk prior to testing and, if the disease-causing mutation in their family is not included on the test, their carrier risk would remain unchanged. Genetic counseling is recommended for patients with a family history of genetic disease so that risk figures based on actual family history can be determined and discussed along with potential implications for reproduction. Horizon carrier screening has been developed to identify the reproductive risks for monogenic inherited conditions. Even when one or both members of a couple screen negative for pathogenic variants in a specific gene, the disease risk for their offspring is not zero. There is still a low risk for the condition in their offspring due to a number of different mechanisms that are not detected by Horizon including, but not limited to, pathogenic variant(s) in the tested gene or in a different gene not included on Horizon, pathogenic variant(s) in an upstream regulator, uniparental disomy, de novo mutation(s), or digenic or polygenic inheritance.

Additional Comments

These analyses generally provide highly accurate information regarding the patient's carrier status. Despite this high level of accuracy, it should be kept in mind that there are many potential sources of diagnostic error, including misidentification of samples, polymorphisms, or other rare genetic variants that interfere with analysis. Families should understand that rare diagnostic errors may occur for these reasons.

